

POSITROIDS, KNOTS, AND q, t -CATALAN NUMBERS

PAVEL GALASHIN AND THOMAS LAM

ABSTRACT. We relate the mixed Hodge structure on the cohomology of open positroid varieties (in particular, their Betti numbers over \mathbb{C} and point counts over \mathbb{F}_q) to Khovanov–Rozansky homology of associated links. We deduce that the mixed Hodge polynomials of top-dimensional open positroid varieties are given by rational q, t -Catalan numbers. Via the curious Lefschetz property of cluster varieties, this implies the q, t -symmetry and unimodality properties of rational q, t -Catalan numbers. We show that the q, t -symmetry phenomenon is a manifestation of Koszul duality for category \mathcal{O} , and discuss relations with open Richardson varieties and extension groups of Verma modules.

CONTENTS

Introduction	1
1. Main results	3
2. Point count and the HOMFLY polynomial	13
3. Soergel bimodules, Rouquier complexes, and Khovanov–Rozansky homology	15
4. Cohomology of positroid and Richardson varieties	23
5. Equivariant derived categories of flag varieties	32
6. Proof of Proposition 5.3	35
7. Proof of Proposition 5.4	37
8. Ordinary cohomology, Koszul duality, and Verma modules	42
9. Catalan numbers associated to positroid varieties	44
References	47

INTRODUCTION

The binomial coefficients $\binom{n}{k}$ have natural q -analogs $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_q$, known as Gaussian polynomials. On the other hand, the rational Catalan numbers $C_{k,n-k} := \frac{1}{n} \binom{n}{k}$ (defined for $\gcd(k, n) = 1$) have two different well-studied q -analogs: the area generating function $\sum_{P \in \text{Dyck}_{k,n-k}} q^{\text{area}(P)}$ of rational Dyck paths [CR64], and the polynomial $\frac{1}{[n]_q} \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_q$ going back to [Mac60].

The Poincaré polynomial of the complex Grassmannian $\text{Gr}(k, n)$, and the number of points of $\text{Gr}(k, n)$ over a finite field \mathbb{F}_q , are both well known to be given by $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_q$. We give a Catalan analog of this statement by considering the top-dimensional positroid variety $\Pi_{k,n}^\circ \subset \text{Gr}(k, n)$, introduced in [KLS13] building on the results of [Pos06]. The space $\Pi_{k,n}^\circ$ is the subspace of $\text{Gr}(k, n)$ where all cyclically consecutive Plücker coordinates are non-vanishing. We show that, up to a simple factor, the mixed Hodge polynomial $\mathcal{P}(\Pi_{k,n}^\circ; q, t)$ coincides with the rational q, t -Catalan number

Date: August 12, 2023.

2010 Mathematics Subject Classification. Primary: 14M15. Secondary: 14F05, 05A15, 57K18.

Key words and phrases. Positroid varieties, q, t -Catalan numbers, HOMFLY polynomial, Khovanov–Rozansky homology, mixed Hodge structure, equivariant cohomology, Verma modules, Koszul duality.

P.G. was supported by an Alfred P. Sloan Research Fellowship and by the National Science Foundation under Grants No. DMS-1954121 and No. DMS-2046915. T.L. was supported by a von Neumann Fellowship from the Institute for Advanced Study and by grants DMS-1464693 and DMS-1953852 from the National Science Foundation.

$C_{k,n-k}(q, t)$ introduced in [LW09] in the study of Macdonald polynomials [GH96, Hag08]. It follows that the Poincaré polynomial of $\Pi_{k,n}^\circ$ equals $\sum_{P \in \text{Dyck}_{k,n-k}} q^{\text{area}(P)}$, while the point count $\#\Pi_{k,n}^\circ(\mathbb{F}_q)$ equals $\frac{1}{[n]_q} [n]_q$, both up to a simple factor.

The coincidence of the Poincaré polynomial and the point count of $\text{Gr}(k, n)$ is reflected in the purity of the mixed Hodge structure on the cohomology of $\text{Gr}(k, n)$. Purity holds for many spaces of interest in combinatorics, e.g. for complements of hyperplane arrangements. By contrast, the mixed Hodge structure on $H^\bullet(\Pi_{k,n}^\circ)$ is not pure, and simultaneously yields both of the natural q -analogs of rational Catalan numbers discussed above.

Our proof proceeds via relating both sides to Khovanov–Rozansky knot homology [KR08a, KR08b, Kho07]. Our main result connects the cohomology of arbitrary open positroid varieties, and more generally open Richardson varieties in generalized flag varieties, to knot homology.

Connections between knot invariants and Macdonald theory have received an enormous amount of attention in recent years; see e.g. [Che13, GORS14, GN15, Hag16, Mel17, ORS18, EH19, HM19]. In particular, Khovanov–Rozansky homology of torus knots and links was computed in [Mel17, Hog17, EH19, HM19]. For torus knots, the answer is given by the rational q, t -Catalan numbers.

Our main results are described in detail in the next section. We start by highlighting some consequences of our approach from several points of view.

Combinatorics. The coefficients of the Gaussian polynomial $[n]_q$ are well known to form a unimodal palindromic sequence. A geometric explanation for this phenomenon is the hard Lefschetz theorem for the cohomology of $\text{Gr}(k, n)$. It follows from the results of [LS16, GL19] that the cohomology of $\Pi_{k,n}^\circ$ satisfies the curious Lefschetz property which, combined with our main result, yields a geometric proof that $C_{k,n-k}(q, t)$ is q, t -symmetric and unimodal. Furthermore, our work produces a whole family of q, t -symmetric and unimodal polynomials, which includes $C_{k,n-k}(q, t)$ as a special case. We discuss their $q = t = 1$ specialization in Section 9 where we obtain a new combinatorial interpretation for rational Catalan numbers in terms of certain kinds of pipe dreams; see Figure 5.


Knot theory. We introduce a class of *Richardson links*, which are closures of braids of the form $\beta(w) \cdot \beta(v)^{-1}$ for pairs of permutations $v, w \in S_n$ such that $v \leq w$ in the Bruhat order. We give a geometric interpretation of the top a -degree coefficient¹ of Khovanov–Rozansky (KR) homology and of the HOMFLY polynomial [FYH⁺85, PT87] for such links. When a Richardson link is a knot, we show that the associated q, t -polynomial is q, t -symmetric. Our investigations suggest that KR homology may have hitherto unstudied unimodality and Lefschetz-type properties. Our results generalize equally well to other Dynkin types.

Representation theory. We show that the q, t -symmetry property is a consequence of the Koszul duality phenomenon [BGS96, BY13] in the derived category of the flag variety.

The computation of the extension groups $\text{Ext}^\bullet(M_v, M_w)$ between Verma modules in the principal block \mathcal{O}_0 of the Bernstein–Gelfand–Gelfand category \mathcal{O} (see e.g. [Hum08]) is a classical, still open problem. We show that these extension groups are isomorphic to knot-homology groups. Along the same vein, we show that the R -polynomials of Kazhdan and Lusztig [KL79, KL80] are certain coefficients of the HOMFLY polynomial.

Algebraic geometry. Our results provide evidence for a $P = W$ conjecture relating the weight filtration of $\Pi_{k,n}^\circ$ with the perverse filtration of the compactified Jacobian $J_{k,n-k}$; see Section 1.12.2.

¹The top a -degree coefficient encodes the zeroth Hochschild cohomology (1.17) which sometimes corresponds to the *bottom* a -degree in the literature. Our conventions are chosen so that the a -degree in KR homology matches the a -degree in HOMFLY polynomial.


 FIGURE 1. Computing the rational q, t -Catalan number $C_{3,5}(q, t)$.

1. MAIN RESULTS

We give a detailed description of our main results. The historical context and motivation for our work is delayed to Section 1.12. We give the full background on the below objects in the main body of the paper.

1.1. Rational q, t -Catalan numbers. Let a and b be coprime positive integers. The *rational q, t -Catalan number* $C_{a,b}(q, t) \in \mathbb{N}[q, t]$ was introduced by Loehr–Warrington [LW09] (see also [GM13, GM14]), generalizing the work of Garsia–Haiman [GH96]. It is defined as

$$(1.1) \quad C_{a,b}(q, t) := \sum_{P \in \text{Dyck}_{a,b}} q^{\text{area}(P)} t^{\text{div}(P)},$$

where $\text{Dyck}_{a,b}$ is the set of lattice paths P inside a rectangle of height a and width b that stay above the diagonal, $\text{area}(P)$ is the number of unit squares fully contained between P and the diagonal, and $\text{div}(P)$ is the number of pairs (h, v) satisfying the following conditions: h is a horizontal step of P , v is a vertical step of P that appears to the right of h , and there exists a line of slope a/b (parallel to the diagonal) intersecting both h and v . For example,

$$(1.2) \quad C_{3,5}(q, t) = q^4 + q^3 t + q^2 t^2 + q^2 t + q t^3 + q t^2 + t^4,$$

as shown in Figure 1.

1.2. Positroid varieties in the Grassmannian. The *Grassmannian* $\text{Gr}(k, n)$ is the space of k -dimensional linear subspaces of \mathbb{C}^n . Building on Postnikov’s cell decomposition [Pos06] of its totally nonnegative part, Knutson–Lam–Speyer [KLS13] constructed a stratification

$$(1.3) \quad \text{Gr}(k, n) = \bigsqcup_{f \in \mathbf{B}_{k,n}} \Pi_f^\circ,$$

where the (*open*) *positroid varieties* Π_f° are defined as the non-empty intersections of cyclic rotations of n Schubert cells. These varieties also arise in Poisson geometry [BGY06] and in the study of scattering amplitudes [AHBC⁺16]. Open positroid varieties are indexed by a finite set $\mathbf{B}_{k,n}$ of *bounded affine permutations*, and for $f \in \mathbf{B}_{k,n}$ the reduction of f modulo n is a permutation $\bar{f} \in S_n$. See Section 4.1 for further background.

Let $f_{k,n} \in \mathbf{B}_{k,n}$ be the bounded affine permutation given by $f_{k,n}(i) = i + k$. The positroid stratification (1.3) contains a unique open stratum, the *top-dimensional positroid variety* $\Pi_{k,n}^\circ := \Pi_{f_{k,n}}^\circ$, which can be described explicitly as

$$(1.4) \quad \Pi_{k,n}^\circ := \{V \in \text{Gr}(k, n) \mid \Delta_{1,2,\dots,k}(V), \Delta_{2,3,\dots,k+1}(V), \dots, \Delta_{n,1,\dots,k-1}(V) \neq 0\},$$

consisting of subspaces whose cyclically consecutive Plücker coordinates are non-vanishing.

For each $f \in \mathbf{B}_{k,n}$, the space Π_f° is a smooth algebraic variety. Two basic questions one can ask are:

- (1) What is the number of points in $\Pi_f^\circ(\mathbb{F}_q)$ over a finite field \mathbb{F}_q with q elements?
- (2) What are the Betti numbers of Π_f° considered as a complex manifold?

These two questions are related by the *mixed Hodge structure* [Del71] on cohomology. The cohomology ring $H^\bullet(\Pi_f^\circ) = H^\bullet(\Pi_f^\circ, \mathbb{C})$ of an open positroid variety is of Hodge–Tate type, and we have a *Deligne splitting*

$$(1.5) \quad H^k(\Pi_f^\circ, \mathbb{C}) \cong \bigoplus_{p \in \mathbb{Z}} H^{k, (p, p)}(\Pi_f^\circ, \mathbb{C}).$$

Since Π_f° is smooth, we have that $H^{k, (p, p)}$ vanishes unless $k/2 \leq p \leq k$. We view (1.5) as a bigrading on $H^\bullet(\Pi_f^\circ)$ and let $\mathcal{P}(\Pi_f^\circ; q, t)$ be the suitably renormalized (see (4.6)) Poincaré polynomial of this bigraded vector space, called the *mixed Hodge polynomial*.

We are ready to state the most important special case of our main result.

Theorem 1.1. *Assume that $\gcd(k, n) = 1$. Then*

$$(1.6) \quad \mathcal{P}(\Pi_{k,n}^\circ; q, t) = \left(q^{\frac{1}{2}} + t^{\frac{1}{2}} \right)^{n-1} C_{k, n-k}(q, t).$$

The equality (1.6) arises as a conjecture from the works [STZ17, STWZ19] and we thank Vivek Shende for drawing our attention to the conjecture; see Section 1.12.2 for further discussion. We generalize Theorem 1.1 to all positroid varieties in Theorem 1.17 below.

Let us discuss the specializations of Theorem 1.1 that give answers to Questions (1) and (2) above. Denote

$$[n]_q := 1 + q + \cdots + q^{n-1}, \quad [n]_q! := [1]_q [2]_q \cdots [n]_q, \quad \begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{[n]_q!}{[k]_q! [n-k]_q!}.$$

Corollary 1.2. *The Poincaré polynomial and point count of $\Pi_{k,n}^\circ$ are*

$$(1.7) \quad \mathcal{P}(\Pi_{k,n}^\circ; q) = (q+1)^{n-1} \cdot C_{k, n-k}(q^2, 1);$$

$$(1.8) \quad \#\Pi_{k,n}^\circ(\mathbb{F}_q) = (q-1)^{n-1} \cdot \frac{1}{[n]_q} \begin{bmatrix} n \\ k \end{bmatrix}_q.$$

Our proof of Theorem 1.1 involves a number of ingredients, including Khovanov–Rozansky knot homology and derived categories of flag varieties. The point count specialization (1.8) requires less advanced machinery and we give a quicker elementary proof in Section 2. Associating a *link* $\hat{\beta}_f$ to each positroid variety Π_f° (Section 1.5), we compare the point count $\#\Pi_f^\circ(\mathbb{F}_q)$ to the HOMFLY polynomial of $\hat{\beta}_f$ (Section 1.6). The HOMFLY polynomial is categorified by Khovanov–Rozansky knot homology, and our proof of Theorem 1.1 may be considered a “categorification” of the point count computation.

We have the following elegant but baffling corollary.

Corollary 1.3. *Let $\gcd(k, n) = 1$. Then the probability that a uniformly random k -dimensional subspace of $(\mathbb{F}_q)^n$ belongs to $\Pi_{k,n}^\circ(\mathbb{F}_q)$ is given by*

$$\text{Prob}(V \in \Pi_{k,n}^\circ(\mathbb{F}_q)) = \frac{(q-1)^n}{q^n - 1}.$$

The probability $\frac{(q-1)^n}{q^n - 1}$ does not depend on k . We do not have a direct explanation for this phenomenon.

1.3. Cluster structure and the curious Lefschetz theorem. Since the work of Scott [Sco06], positroid varieties have been expected to admit a natural *cluster algebra* [FZ02] structure arising from Postnikov diagrams. We recently proved this conjecture building on the results of [Lec16, MS17, SSBW19].

Theorem 1.4 ([GL19]). *The coordinate ring of each positroid variety Π_f° is isomorphic to the associated cluster algebra.*

This result allows one to study Π_f° as a *cluster variety*, and for such spaces the mixed Hodge structure can be explored using the machinery developed by Lam–Speyer [LS16], whose work implies the following properties of the mixed Hodge polynomials $\mathcal{P}(\Pi_f^\circ; q, t)$.

Theorem 1.5 ([LS16],[GL19]). *For each $f \in \mathbf{B}_{k,n}$, the mixed Hodge polynomial $\mathcal{P}(\Pi_f^\circ; q, t) \in \mathbb{N}[q^{\frac{1}{2}}, t^{\frac{1}{2}}]$ has the following properties:*

- (i) *q, t -symmetry: $\mathcal{P}(\Pi_f^\circ; q, t) = \mathcal{P}(\Pi_f^\circ; t, q)$;*
- (ii) *q, t -unimodality: for each d , the coefficients of $\mathcal{P}(\Pi_f^\circ; q, t)$ at $q^d t^0, q^{d-1} t^1, \dots, q^0 t^d$ form a unimodal sequence;*
- (iii) *$\mathcal{P}(\Pi_f^\circ; 1, q^2)$ equals the Poincaré polynomial of Π_f° (considered as a variety over \mathbb{C});*
- (iv) *$q^{\frac{1}{2} \dim \Pi_f^\circ} \cdot \mathcal{P}(\Pi_f^\circ; q, t)|_{t^{\frac{1}{2}} = -q^{-\frac{1}{2}}}$ equals the point count $\#\Pi_f^\circ(\mathbb{F}_q)$.*

See Example 1.9 below.

Parts (i) and (ii) are consequences of the *curious Lefschetz property*, formalized in [HRV08] and proven to hold for certain cluster varieties in [LS16]; see Section 4.3.

1.4. The Catalan variety. Let $T \cong (\mathbb{C}^*)^{n-1}$ be the group of diagonal matrices in $\mathrm{PGL}_n(\mathbb{C})$: it is the quotient of the group of diagonal $n \times n$ matrices by the group of scalar matrices. The group T acts on $\mathrm{Gr}(k, n)$ preserving the positroid stratification. For $u \in S_n$, let

$$(1.9) \quad c(u) := \text{the number of cycles of } u,$$

and let $\mathbf{B}_{k,n}^{c=1} := \{f \in \mathbf{B}_{k,n} \mid c(\bar{f}) = 1\}$. The following observation is proved in Section 4.2.

Proposition 1.6. *The action of T on Π_f° is free if and only if the permutation \bar{f} is a single cycle.*

For $f \in \mathbf{B}_{k,n}^{c=1}$, the quotient $\mathcal{X}_f^\circ := \Pi_f^\circ/T$ is again a smooth affine variety that we call a *positroid configuration space*; see also [AHLS20]. It is a cluster variety (with no frozen variables, since the T -action on the frozen variables of Π_f° is free), and thus Theorem 1.5 applies to it.

Proposition 1.7. *For $f \in \mathbf{B}_{k,n}^{c=1}$, the mixed Hodge polynomials of Π_f° and \mathcal{X}_f° are related by:*

$$(1.10) \quad \mathcal{P}(\Pi_f^\circ; q, t) = \left(q^{\frac{1}{2}} + t^{\frac{1}{2}}\right)^{n-1} \cdot \mathcal{P}(\mathcal{X}_f^\circ; q, t).$$

When $\mathrm{gcd}(k, n) = 1$, we have $f_{k,n} \in \mathbf{B}_{k,n}^{c=1}$. The quotient $\mathcal{X}_{k,n}^\circ := \Pi_{k,n}^\circ/T$ satisfies

$$(1.11) \quad \mathcal{P}(\mathcal{X}_{k,n}^\circ; q, t) = C_{k,n-k}(q, t),$$

and we refer to $\mathcal{X}_{k,n}^\circ$ as *the Catalan variety*. Let $d_{k,n} := (k-1)(n-k-1) = \dim(\mathcal{X}_{k,n}^\circ)$. We obtain the following as a consequence of Theorem 1.5.

Corollary 1.8. *Assume that $\mathrm{gcd}(k, n) = 1$. We have:*

- (i) *q, t -symmetry: $C_{k,n-k}(q, t) = C_{k,n-k}(t, q)$;*
- (ii) *q, t -unimodality: for each d , the coefficients of $C_{k,n-k}(q, t)$ at $q^d t^0, q^{d-1} t^1, \dots, q^0 t^d$ form a unimodal sequence;*
- (iii) *the Poincaré polynomial of $\mathcal{X}_{k,n}^\circ$ is given by*

$$(1.12) \quad \sum_d q^{\frac{d}{2}} \dim H^{d_{k,n}-d}(\mathcal{X}_{k,n}^\circ) = C_{k,n-k}(q, 1) = \sum_{P \in \mathrm{Dyck}_{k,n-k}} q^{\mathrm{area}(P)};$$

- (iv) *the number of \mathbb{F}_q -points of $\mathcal{X}_{k,n}^\circ$ is given by*

$$(1.13) \quad \#\mathcal{X}_{k,n}^\circ(\mathbb{F}_q) = \frac{1}{[n]_q} \begin{bmatrix} n \\ k \end{bmatrix}_q = q^{\frac{1}{2}d_{k,n}} \cdot C_{k,n-k}(q, 1/q).$$

H^k	H^0	H^1	H^2	H^3	H^4	H^5	H^6	H^7	H^8
$k - p = 0$	1	0	1	0	1	0	1	0	1
$k - p = 1$					1	0	1		

TABLE 1. The mixed Hodge table recording the dimensions of $H^{k,(p,p)}(\mathcal{X}_{3,8}^\circ)$ for the cluster algebra of type E_8 ; see [LS16, Table 5]. The dimensions agree with the coefficients of $C_{3,5}(q, t)$; see Example 1.9.




FIGURE 2. Braids and links associated to positroid varieties.

While part (i) is known, the remaining parts of Corollary 1.8 appear to be new; see Section 1.12.1. Note also that the odd Betti numbers of $\mathcal{X}_{k,n}^\circ$ vanish, a phenomenon that we do not have a direct explanation for. Parts (iii)–(iv) may be deduced directly from Corollary 1.2 using Proposition 1.6.

Example 1.9. Let $k = 3$ and $n = 8$. The coordinate ring of $\mathcal{X}_{3,8}^\circ$ is a cluster algebra of type E_8 (with no frozen variables). The associated *mixed Hodge table* is given in Table 1; see [LS16, Table 5]. The grading conventions (4.6) are chosen so that the first row contributes $q^4 + q^3t + q^2t^2 + qt^3 + t^4$ while the second row contributes $q^2t + qt^2$ to $\mathcal{P}(\mathcal{X}_{3,8}^\circ; q, t)$. Note that all odd cohomology groups vanish, which is why all monomials have integer powers of q and t . Comparing the result with (1.2), we find $\mathcal{P}(\mathcal{X}_{3,8}^\circ; q, t) = C_{3,5}(q, t)$, in agreement with Theorem 1.1.

The polynomial $C_{3,5}(q, t)$ given in (1.2) is indeed q, t -symmetric. It is also q, t -unimodal: fixing the total degree of q and t , it splits into polynomials $q^4 + q^3t + q^2t^2 + qt^3 + t^4$ and $q^2t + qt^2$, both of which have unimodal coefficient sequences, corresponding to the rows of Table 1. We also have $C_{3,5}(q, 1) = q^4 + q^3 + 2q^2 + 2q + 1$; the coefficient of $q^{d/2}$ is equal to $\dim H^{d, k, n-d}(\Pi_{3,8}^\circ)$ for each d (these coefficients are column sums in Table 1). This agrees with (1.12).

1.5. Links associated to positroid varieties. Let us say that a permutation $w \in S_n$ is k -Grassmannian if $w^{-1}(1) < w^{-1}(2) < \dots < w^{-1}(k)$ and $w^{-1}(k+1) < \dots < w^{-1}(n)$. We denote by \leq the (strong) Bruhat order on S_n . Let $\mathbf{Q}_{k,n}$ denote the set of pairs (v, w) of permutations such that $v \leq w$ and w is k -Grassmannian. The following result is well known; see Proposition 4.2.

Proposition 1.10 ([KLS13]). *There exists a bijection $(v, w) \mapsto f_{v,w}$ between $\mathbf{Q}_{k,n}$ and $\mathbf{B}_{k,n}$ such that for every $f = f_{v,w} \in \mathbf{B}_{k,n}$, we have $\bar{f} = wv^{-1}$.*

For example, when $f = f_{k,n}$, we have $v = \text{id}$ and the permutation $w = \bar{f}$ sends $i \mapsto i + k$ modulo n for all $i \in [n]$. The dimension of Π_f° equals $\ell_{v,w} := \ell(w) - \ell(v)$, where $\ell(u)$ is the number of inversions of $u \in S_n$.

The group S_n is generated by simple transpositions $s_i = (i, i+1)$ for $1 \leq i \leq n-1$. Similarly, let \mathbf{B}_n be the *braid group* on n strands, generated by $\sigma_1, \dots, \sigma_{n-1}$ with relations $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ and $\sigma_i \sigma_j = \sigma_j \sigma_i$ for $|i - j| > 1$. Connecting the corresponding endpoints of a braid β gives rise to a *link* called the *closure* $\hat{\beta}$ of β ; see Figure 2.

For each element $u \in S_n$, let $\beta(u)$ denote the corresponding braid, obtained by choosing a reduced word $u = s_{i_1} s_{i_2} \dots s_{i_{\ell(u)}}$ for u and then replacing each s_i with σ_i .

Definition 1.11. For $f = f_{v,w} \in \mathbf{B}_{k,n}$, we set

$$(1.14) \quad \beta_f := \beta(w) \cdot \beta(v)^{-1}.$$

We refer to the closure $\hat{\beta}_f$ as *the link associated to f* . See Figure 2 for an example.

The link $\hat{\beta}_f$ is a *knot* (i.e., has one connected component) if and only if $f \in \mathbf{B}_{k,n}^{c=1}$; see Proposition 1.6.

1.6. HOMFLY polynomial. The *HOMFLY polynomial* $P(L) = P(L; a, z)$ of an (oriented) link L is defined by the skein relation

$$(1.15) \quad aP(L_+) - a^{-1}P(L_-) = zP(L_0) \quad \text{and} \quad P(\bigcirc) = 1,$$

where \bigcirc denotes the unknot and L_+, L_-, L_0 are three links whose planar diagrams locally differ as follows.



Example 1.12. For $n = 2$, we may take L_+ to be the closure of σ_1 , in which case L_- is the closure of σ_1^{-1} and $L_0 = \bigcirc \bigcirc$ is the 2-component unlink. Applying (1.15), we find $P(L_0) = \frac{a-a^{-1}}{z}$.

Surprisingly, the HOMFLY polynomial computes the number of \mathbb{F}_q -points of *any* positroid variety.

Theorem 1.13. For all $f \in \mathbf{B}_{k,n}$, let $P_f^{\text{top}}(q)$ be obtained from the top a -degree term of $P(\hat{\beta}_f; a, z)$ by substituting $a := q^{-\frac{1}{2}}$ and $z := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$. Then

$$(1.16) \quad \#\Pi_f^\circ(\mathbb{F}_q) = (q-1)^{n-1} \cdot P_f^{\text{top}}(q).$$

Remark 1.14. When $\gcd(k, n) = 1$, we have $f_{k,n} \in \mathbf{B}_{k,n}^{c=1}$, and the associated knot $\hat{\beta}_{f_{k,n}}$ is the $(k, n-k)$ -torus knot; see Figure 2(right). The value of $P(\hat{\beta}_{f_{k,n}}; a, z)$ was computed in [Jon87], and its relationship with Catalan numbers was clarified in [Gor12]. Thus, (1.8) follows from Theorem 1.13 as a direct corollary.

Example 1.15. For $k = 3, n = 8$, one calculates (for instance, using Sage²) that

$$P(\hat{\beta}_{f_{k,n}}; a, z) = \frac{z^8 + 8z^6 + 21z^4 + 21z^2 + 7}{a^8} - \frac{z^6 + 7z^4 + 14z^2 + 8}{a^{10}} + \frac{z^2 + 2}{a^{12}}.$$

Substituting $a := q^{-\frac{1}{2}}$ and $z := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$ into $\frac{z^8 + 8z^6 + 21z^4 + 21z^2 + 7}{a^8}$, we get

$$P_f^{\text{top}}(q) = q^8 + q^6 + q^5 + q^4 + q^3 + q^2 + 1 = q^4 \cdot C_{3,5}(q, 1/q).$$

This agrees with (1.13) and (1.16).

1.7. Richardson varieties. Let G be a complex semisimple algebraic group of adjoint type, and choose a pair $B, B_- \subset G$ of opposite Borel subgroups. Let $T := B \cap B_-$ be the maximal torus and $W := N_G(T)/T$ the associated Weyl group. We have Bruhat decompositions $G = \bigsqcup_{w \in W} BwB = \bigsqcup_{v \in W} B_-vB$, and the intersection $BwB \cap B_-vB$ is nonempty if and only if $v \leq w$ in the Bruhat order on W . For $v \leq w$, we denote by $R_{v,w}^\circ := (BwB \cap B_-vB)/B$ an *open Richardson variety* inside the (*generalized*) *complete flag variety* G/B . The varieties $R_{v,w}^\circ$ form a stratification of G/B .

Now suppose $G = \text{PGL}_n(\mathbb{C})$. We have $W = S_n$, the subgroups $B, B_- \subset G$ consist of upper and lower triangular matrices, and $T \cong (\mathbb{C}^*)^{n-1}$ is the group of diagonal matrices modulo scalar matrices. In this case we denote the generalized flag variety G/B by $\text{Fl}(n)$. By Proposition 1.10, positroid varieties correspond to pairs $v \leq w$ of permutations such that w is k -Grassmannian. The projection map $\text{Fl}(n) \rightarrow \text{Gr}(k, n)$ restricts to an isomorphism $R_{v,w}^\circ \cong \Pi_f^\circ$ for every permutation

²<https://doc.sagemath.org/html/en/reference/knots/sage/knots/link.html>

Title	n	v	w	$\ell_{v,w}$	$\hat{\beta}_{v,w}$	$R_{v,w}^\circ$	$H^{k,(p,p)}(R_{v,w}^\circ)$	$H_{T,c}^{k,(p,p)}(R_{v,w}^\circ)$	$H^{k,(p)}(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet))$	$H^{k,(p)}(\mathbb{H}\mathbb{H}^0(F_{v,w}^\bullet))$
Unknot-I	1	id	id	0	\bigcirc	pt	$k=0, p=0$	$k=0, p=0$	$k=0, p=0$	$k=0, p=0$
Unknot-II	2	id	s_1	1	\bigcirc	$\Pi_{1,2}^\circ$	$k=0, p=0$ $k=1, p=1$	$k=1, p=0$	$k=0, p=0$ $k=-1, p=1$	$k=0, p=0$
2-cpt. unlink	2	s_1	s_1	0	\bigcirc	pt	$k=0, p=0$	$k=2p, p \in \mathbb{Z}_{\geq 0}$	$k=0, p=0$	$k=0, p \in \mathbb{Z}_{\geq 0}$
Hopf link	4	id	$f_{2,4}$	4	\bigcirc	$\Pi_{2,4}^\circ$	Table 3	$k=4, p=0$ $k=2+2p, p \in \mathbb{Z}_{>2}$	Table 3	$k=0, p=0$ $k=-2, p \in \mathbb{Z}_{>2}$
Trefoil knot	5	id	$f_{2,5}$	6	\bigcirc	$\Pi_{2,5}^\circ$	Table 3	$k=6, p=0$ $k=8, p=2$	Table 3	$k=0, p=0$ $k=-2, p=2$

TABLE 2. Summary of examples computed in Sections 3.6 and 4.6, illustrating Theorems 1.16 and 1.17. The last four columns list all values of k, p , for which the corresponding bigraded component is nonzero (in each case, it is 1-dimensional). We have $f_{2,4} = s_2 s_1 s_3 s_2$ and $f_{2,5} = s_3 s_2 s_1 s_4 s_3 s_2$.

$f = f_{v,w} \in \mathbf{B}_{k,n}$ (see Proposition 4.3). Thus, positroid varieties are special cases of Richardson varieties. One can similarly associate a braid $\beta_{v,w} := \beta(w) \cdot \beta(v)^{-1}$ to any pair $v \leq w$ in S_n and consider its closure $\hat{\beta}_{v,w}$. We refer to links of the form $\hat{\beta}_{v,w}$ as *Richardson links*.

The point count $\#R_{v,w}(\mathbb{F}_q)$ is given by the *Kazhdan–Lusztig R -polynomial* [KL79, KL80], and both the statement and the proof of Theorem 1.13 generalize to this setting (for G of arbitrary type); see Theorems 2.1 and 2.3.

1.8. Main result: ordinary cohomology. Our results for the positroid variety $\Pi_{k,n}^\circ$ are special cases of a statement which applies to open Richardson varieties in arbitrary Dynkin type. This includes all positroid varieties Π_f° for $f \in \mathbf{B}_{k,n}$, where the number of cycles $c(\bar{f})$ can be arbitrary. We start with the non-equivariant version of our result.

Let $\mathfrak{h} := \text{Lie}(T)$ be the Cartan subalgebra of $\text{Lie}(G)$ corresponding to T , and denote $R := \mathbb{C}[\mathfrak{h}] = \text{Sym}_{\mathbb{C}} \mathfrak{h}^*$. The ring R is graded so that the elements of $\mathfrak{h}^* \subset R$ have polynomial degree 2. For $G = \text{PGL}_n(\mathbb{C})$, $R = \mathbb{C}[y_1, \dots, y_{n-1}]$ is the polynomial ring. Since W is a Coxeter group, we can consider the category SBim of *Soergel bimodules* [Soe92, EMTW20]. Each object $B \in \text{SBim}$ is a graded R -bimodule, and we will be interested in its R -invariants, which by definition form the *zeroth Hochschild cohomology of B* :

$$(1.17) \quad \mathbb{H}\mathbb{H}^0(B) := \{b \in B \mid rb = br \text{ for all } r \in R\}.$$

Thus, $\mathbb{H}\mathbb{H}^0(B)$ is a graded R -module. Denote

$$(1.18) \quad \mathbb{H}\mathbb{H}_{\mathbb{C}}^0(B) := \mathbb{H}\mathbb{H}^0(B) \otimes_R \mathbb{C},$$

where $\mathbb{C} = R/(\mathfrak{h}^*)$ is the R -module on which \mathfrak{h}^* acts by 0. While the functor $\mathbb{H}\mathbb{H}^0$ involves Soergel bimodules, the functor $\mathbb{H}\mathbb{H}_{\mathbb{C}}^0$ involves *Soergel modules* instead; see Corollary 3.6.

To any element $u \in W$, Rouquier [Rou04] associates two cochain complexes $F^\bullet(u)$ and $F^\bullet(u)^{-1}$ of Soergel bimodules. For a braid $\beta_{v,w} = \beta(w) \cdot \beta(v)^{-1}$, we set $F_{v,w}^\bullet := F^\bullet(w) \otimes_R F^\bullet(v)^{-1}$. Applying the functor $\mathbb{H}\mathbb{H}_{\mathbb{C}}^0$ to each term of this complex yields a complex $\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet)$ of graded R -modules. Taking its cohomology

$$(1.19) \quad \mathbb{H}\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet) := H^\bullet(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet)),$$

we get a bigraded vector space. Explicitly, we have

$$\mathbb{H}\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet) = \bigoplus_{k,p \in \mathbb{Z}} H^{k,(p)}(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet)),$$

where $H^{k,(p)}(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet))$ is the polynomial degree $2p$ part of $H^k(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^\bullet))$.

Recall from (1.5) that we have a bigrading on $H^\bullet(R_{v,w}^\circ)$ coming from the Deligne splitting.

Theorem 1.16. *For all $v \leq w \in W$ and $k, p \in \mathbb{Z}$, we have*

$$(1.20) \quad H^{k, (p, p)}(R_{v, w}^\circ) \cong H^{-k, (p)}(\mathbb{H}\mathbb{H}_\mathbb{C}^0(F_{v, w}^\bullet)).$$

See Tables 2 and 3 for examples.

1.9. Main result: equivariant cohomology. The spaces $\mathbb{H}\mathbb{H}^0(F_{v, w}^\bullet)$ and $\mathbb{H}\mathbb{H}_\mathbb{C}^0(F_{v, w}^\bullet)$ are closely related. By Theorem 1.16, $\mathbb{H}\mathbb{H}_\mathbb{C}^0(F_{v, w}^\bullet)$ yields the cohomology of $R_{v, w}^\circ$. It turns out that $\mathbb{H}\mathbb{H}^0(F_{v, w}^\bullet)$ yields the *torus-equivariant* cohomology of $R_{v, w}^\circ$.

The algebraic torus T acts on each Richardson variety $R_{v, w}^\circ$, and thus we can consider its T -equivariant cohomology with compact support, denoted $H_{T, c}^\bullet(R_{v, w}^\circ)$. It is equipped with an action of the ring $H_T^\bullet(\text{pt}) \cong R$. Similarly to (1.5), $H_{T, c}^\bullet(R_{v, w}^\circ)$ admits a second grading via the mixed Hodge structure and is therefore a bigraded R -module.

Theorem 1.17. *For all $v \leq w \in W$, we have an isomorphism of bigraded R -modules*

$$(1.21) \quad H_{T, c}^\bullet(R_{v, w}^\circ) \cong \mathbb{H}\mathbb{H}^0(F_{v, w}^\bullet).$$

For each $k, p \in \mathbb{Z}$, it restricts to a vector space isomorphism

$$(1.22) \quad H_{T, c}^{\ell_{v, w} + 2p + k, (p, p)}(R_{v, w}^\circ) \cong H^{k, (p)}(\mathbb{H}\mathbb{H}^0(F_{v, w}^\bullet)),$$

where $\ell_{v, w} = \ell(w) - \ell(v) = \dim R_{v, w}^\circ$.

See Table 2 for examples. We explain how Theorems 1.1 and 1.16 follow from Theorem 1.17 in Sections 4.4 and 8.1, respectively.

Observe that the grading conventions in (1.20) and (1.22) are quite different. In fact, the two statements are related by an application of the q, t -symmetry (1.23), as we now explain.

1.10. Koszul duality and q, t -symmetry. For any $f \in \mathbf{B}_{k, n}^{c=1}$, the positroid variety \mathcal{X}_f° is a cluster variety [GL19], so the polynomial $\mathcal{P}(\Pi_f^\circ; q, t)$ is q, t -symmetric by Corollary 1.8(i). Richardson varieties are not yet known to admit cluster structures (see [Lec16]), and in particular, the curious Lefschetz theorem of [LS16] cannot yet be applied to conclude that $\mathcal{P}(R_{v, w}^\circ; q, t)$ is q, t -symmetric for arbitrary $v \leq w \in S_n$. In Section 8.2, we show that the q, t -symmetry phenomenon for positroid and Richardson varieties is a manifestation of *Koszul duality* for the derived category of Schubert-constructible sheaves on the flag variety [BGS96, BY13, AR16b].

Theorem 1.18. *For all $v \leq w \in W$ and $k, p \in \mathbb{Z}$, we have an isomorphism*

$$(1.23) \quad H^{k, (p, p)}(R_{v, w}^\circ, \mathbb{C}) \cong H^{\ell_{v, w} + k - 2p, (\ell_{v, w} - p, \ell_{v, w} - p)}(R_{v, w}^\circ, \mathbb{C})$$

of vector spaces. In other words, the polynomial $\mathcal{P}(R_{v, w}^\circ; q, t)$ is q, t -symmetric.

This gives a new proof of the q, t -symmetry of $C_{k, n-k}(q, t)$ for $\gcd(k, n) = 1$.

We now explain the connection to link invariants. Given a Richardson link $\hat{\beta}_{v, w}$, one can consider the bigraded vector spaces $\mathbb{H}\mathbb{H}^0(F^\bullet(\beta_{v, w}))$ and $\mathbb{H}\mathbb{H}_\mathbb{C}^0(F^\bullet(\beta_{v, w}))$, and their suitably renormalized bigraded Hilbert series, denoted $\mathcal{P}_{\text{KR}}^{\text{top}}(\beta_{v, w}; q, t)$ and $\mathcal{P}_{\text{KR}; \mathbb{C}}^{\text{top}}(\beta_{v, w}; q, t)$, respectively; see (3.15)–(3.16).

The polynomial $\mathcal{P}_{\text{KR}}^{\text{top}}(\beta_{v, w}; q, t)$ is the top a -degree coefficient (see the footnote in the introduction) of the celebrated *Khovanov–Rozansky homology* [KR08a, KR08b, Kho07] of $\hat{\beta}_{v, w}$, which is a link invariant, i.e., depends only on the closure $\hat{\beta}_{v, w}$ of $\beta_{v, w}$; see Section 3.5.

Our results imply (see Section 8.1) that when $\hat{\beta}_{v, w}$ is a knot, we have

$$(1.24) \quad \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v, w}; q, t) = \mathcal{P}_{\text{KR}; \mathbb{C}}^{\text{top}}(\beta_{v, w}; q, t) \quad \text{and}$$

$$(1.25) \quad \mathcal{P}(R_{v, w}^\circ/T; q, t) = \left(q^{\frac{1}{2}}t^{\frac{1}{2}}\right)^{\chi(\beta_{v, w})} \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v, w}; q, t),$$

where $\chi(\beta_{v,w}) := \frac{\ell_{v,w} - n + c(f)}{2}$; see (3.12). More generally, Theorem 1.16 implies that for any $v \leq w \in S_n$, one can relate $\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{v,w}; q, t)$ to $\mathcal{P}(R_{v,w}^{\circ}; q, t)$ by a simple transformation. As we show in Corollary 3.13, (1.24) holds more generally for arbitrary knots $\hat{\beta}$.

For a general link $\hat{\beta}$, the question of whether $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$ is q, t -symmetric has been a major open problem³ going back to [DGR06]. For Richardson links, we show that it also amounts to applying Koszul duality.

Corollary 1.19. *For any $v \leq w \in S_n$, we have*

$$\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{v,w}; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{v,w}; t, q).$$

Consequently, by (1.24), if $\hat{\beta}_{v,w}$ is a knot then

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; t, q).$$

1.11. Extensions of Verma modules. The above theorems have representation-theoretic applications to *Verma modules*, which are certain infinite-dimensional modules over the Lie algebra \mathfrak{g} of G . Consider the principal block \mathcal{O}_0 of the Bernstein–Gelfand–Gelfand category \mathcal{O} , and let M_w be the Verma module with highest weight $w(\rho) - \rho$, where ρ is half the sum of positive roots of the root system of \mathfrak{g} . We also denote by L_w the corresponding simple module. The graded dimensions of $\text{Ext}^{\bullet}(M_v, L_w)$ famously coincide with the coefficients of the *Kazhdan–Lusztig P-polynomials* $P_{v,w}(q)$; see e.g. [BGS96, Theorem 3.11.4]. On the other hand, computing extension groups $\text{Ext}^{\bullet}(M_v, M_w)$ is an important open problem; see e.g. [Maz07, DM17].

A *graded* version of \mathcal{O}_0 was introduced by Beilinson–Ginzburg–Soergel [BGS96]. They constructed the (essentially unique) graded lifts of Verma modules M_w (see also [Str03]), thus endowing the space $\text{Ext}^{\bullet}(M_v, M_w)$ with a second grading:

$$\text{Ext}^{\bullet}(M_v, M_w) = \bigoplus_{k,r \in \mathbb{Z}} \text{Ext}^{k,(r/2)}(M_v, M_w).$$

These Ext-groups can be related to the cohomology of open Richardson varieties using the *localization theorem* of [BB81, BK81]. In the case of Kazhdan–Lusztig polynomials, the groups $\text{Ext}^{\bullet}(M_v, L_w)$ are “pure”: the two gradings agree. On the other hand, the bigrading on $\text{Ext}^{\bullet}(M_v, M_w)$ turns out to be quite non-trivial. As a corollary to our approach, we obtain the following result.⁴

Theorem 1.20. *For all $v \leq w \in W$ and $k, r \in \mathbb{Z}$, we have*

$$\dim \text{Ext}^{k,((r-\ell_{v,w})/2)}(M_v, M_w) = \dim H^{k-r,(r/2)}(\mathbb{H}\mathbb{H}_{\mathbb{C}}^0(F_{v,w}^{\bullet})).$$

(In particular, both sides are zero for odd r .)

Thus, while $\text{Ext}^{\bullet}(M_v, L_w)$ gives the Kazhdan–Lusztig polynomials, $\text{Ext}^{\bullet}(M_v, M_w)$ gives the rational q, t -Catalan numbers for $v = \text{id}$ and $w = f_{k,n}$.

1.12. Notes. We collect the historical background and several remarks on the above results.

³At the final stages of the preparation of this manuscript, we learnt that the q, t -symmetry of $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$ for the case when $\hat{\beta}$ is a knot has been established in a very recent preprint [OR20]. (Note added in 2023: see also [GHM21].)

⁴We remark that Soergel’s original work [Soe90] directly relates Ext-groups in category \mathcal{O} and Hom-groups in SBim ; see also (3.9).

1.12.1. **Symmetry and unimodality.** The symmetry and unimodality of the Gaussian polynomial $\begin{bmatrix} n \\ k \end{bmatrix}_q$ are consequences of the hard Lefschetz theorem for $H^\bullet(\text{Gr}(k, n))$. Whereas symmetry is apparent from the combinatorial definition of $\begin{bmatrix} n \\ k \end{bmatrix}_q$ ([Sta12, Proposition 1.7.3]), unimodality is notoriously difficult to see combinatorially. Unimodality was first proven by Sylvester [Syl78], the relation to hard Lefschetz observed by Stanley [Sta84], and a combinatorial proof given by O'Hara [O'H90].

When $a = n$ and $b = n + 1$, $C_{a,b}(q, t)$ recovers the famous q, t -Catalan numbers $C_n(q, t)$ of Garsia and Haiman [GH96] studied extensively in e.g. [GH02, Hag08, Hag03, GM13, GM14]. The fact that $C_n(q, t)$ is q, t -symmetric and q, t -unimodal follows from the results of Haiman [Hai94, Hai02]. For arbitrary a, b , an explanation for the q, t -symmetry property was given by the *Rational Shuffle Conjecture* of [GN15], proved recently in [Mel16, CM18]. The specialization $q^{\frac{1}{2}dk,n} C_{k,n-k}(q, 1/q) = \frac{1}{\begin{bmatrix} n \\ k \end{bmatrix}_q} \begin{bmatrix} n \\ k \end{bmatrix}_q$ in (1.13) is also a consequence of the Rational Shuffle Conjecture. To our knowledge, the q, t -unimodality of $C_{k,n-k}(q, t)$ in Corollary 1.8(ii) is a new result. See also [Stu18, Section 2.2], which includes a specialization of our unimodality result.

1.12.2. **Compactified Jacobians and the $P = W$ conjecture.** We explain the original motivation coming from the results of [STZ17, STWZ19] that led to the statement of Theorem 1.1. The compactified Jacobian $J_{a,b}$ of the plane curve singularity $x^a = y^b$ (with $\gcd(a, b) = 1$) is a compact, singular variety with a long history of connections to Catalan theory. Beauville [Bea99] showed that the Euler characteristic of $J_{a,b}$ is the rational Catalan number $C_{a,b}$ and Piontkowski [Pio07] (see also Lusztig–Smelt [LS91]) showed that the Poincaré polynomial and point count are given by the q -analog $\sum_{P \in \text{Dyck}_{k,n-k}} q^{\text{div}(P)}$. Gorsky and Mazin [GM13, GM14] first suggested the relation between $J_{a,b}$ and q, t -Catalan numbers and since then there has been an explosion of developments relating compactified Jacobians and knot invariants; see e.g. [GORS14, CD16, GN15]. Our work provides evidence for the following conjecture, arising from the works [STZ17, STWZ19]; see [dCHM12] for the original $P = W$ conjecture.

Conjecture 1.21. *There is a deformation retraction from the torus quotient $\mathcal{X}_{k,n}^\circ$ to the compactified Jacobian $J_{k,n-k}$ sending the weight filtration of $H^\bullet(\mathcal{X}_{k,n}^\circ)$ to the perverse filtration of $H^\bullet(J_{k,n-k})$ [MS13, MY14].*

Conjecture 1.21 is motivated by the isomorphism, discovered in [STWZ19], between open positroid varieties and moduli spaces of constructible sheaves associated to Legendrian knots [STZ17]. We thank Vivek Shende for explaining a conjectural *wild* non-abelian Hodge correspondence in that setting.

More generally, when a Richardson link is algebraic (that is, arising as the link of a singularity), one may expect a statement similar to Conjecture 1.21 for the compactified Jacobian of the singularity. See Remark 4.24 for related discussion.

Remark 1.22. After discovering the proof of (1.8) via the HOMFLY polynomial, we found that it can also be deduced from the results of [STZ17, STWZ19]. Our proof is new and yields a generalization (Theorem 1.13) of (1.8) to arbitrary open positroid varieties, and more generally to open Richardson varieties in generalized flag varieties.

1.12.3. **Plabic graph links.** In Section 1.5, we associated a link $\hat{\beta}_f$ to each positroid variety Π_f° . Two other (more complicated) ways of assigning a Legendrian/transverse link to a positroid variety have appeared recently in [STWZ19, FPST17], stated in the language of Postnikov's *plabic graphs* [Pos06]. Conjecturally, the links of [STWZ19, FPST17] are isotopic to our links $\hat{\beta}_f$. We hope to return to this question in future work [GL22]; see also [CGGS21].

1.12.4. **Geometric interpretations and other Dynkin types.** A geometric interpretation of the full triply-graded KR homology was given by Webster–Williamson [WW17]. Our approach

yields a different geometric interpretation of the (doubly-graded) top a -degree part of KR homology. Our geometric interpretation in addition holds for Dynkin types outside type A. The analog of the HOMFLY polynomial in other Dynkin types (as a trace on the Hecke algebra; cf. Section 2.2) was introduced in [Gom06]; see also [Rou17]. For related discussion of knot invariants in other types, see for example [WW08, WW11]; see also [BT20, CGGS20, CP18, CP19, Mel19] for related results.

1.12.5. Odd cohomology vanishing. It follows from the results of [Mel17, HM19] that Khovanov–Rozansky homology of any (positive) torus knot or link is concentrated in even degrees. It is tempting to conjecture that the same property holds for all Richardson knots or links. However, this is not the case: see Examples 4.21, 4.22, and 4.23. See Remark 4.24 for a discussion of the closely related class of *algebraic knots*.

1.12.6. Complements of hyperplane arrangements. The top positroid variety $\Pi_{k,n}^\circ$ may be considered “the complement of a hyperplane arrangement in the Grassmannian”: it is obtained from $\text{Gr}(k, n)$ by removing n hypersurfaces, each given by a linear equation in the Plücker coordinates on $\text{Gr}(k, n)$. More general “Grassmannian hyperplane arrangements” appear naturally in the study of amplituhedra and Grassmann polytopes [Lam16, GL20].

The cohomology of complements of hyperplane arrangements in projective space is very well studied: both the Poincaré polynomial and the point count are simple specializations of the characteristic polynomial. The coincidence is a manifestation of the purity of the mixed Hodge structure [Sha93], a property that also holds for the Grassmannian $\text{Gr}(k, n)$.

1.12.7. Recurrence relations. Our results associate a q, t -polynomial to each positroid variety. One possible advantage of this approach is a recurrence for these polynomials, arising from the recurrence for positroid varieties developed by Muller–Speyer [MS17]. For instance, their results allow to compute the point counts recursively; cf. [GL22]. To compute the Poincaré or the mixed Hodge polynomials, the recurrence of [MS17] yields a long exact sequence for the cohomology. It seems plausible that in favorable cases (e.g., when the odd cohomology vanishes), this sequence may be used to calculate the mixed Hodge polynomials of special families of links as was done in [Mel17, EH19, HM19]. We remark that the latter recurrences pass through complexes of Soergel bimodules which do not come from any braids; an interesting open problem is to understand the positroid/Richardson interpretation of such complexes.

Structure of the paper. In Section 2, we study the relationship between the point count and the HOMFLY polynomial, and prove Theorem 1.13 and its generalization (Theorem 2.3) to open Richardson varieties. In Sections 3 and 4, we discuss background on KR homology and cohomology of positroid varieties, respectively. We deduce Theorem 1.1 from Theorem 1.17 in Section 4.4. In Section 5, we recast our results in the language of equivariant derived categories, and split the main result (Theorem 1.17) into two statements, Propositions 5.3 and 5.4. These statements are proved in Sections 6 and 7, respectively, thereby completing the proof of Theorem 1.17. In Section 8, we deduce the rest of our results (Theorems 1.16, 1.18 and 1.20) from Theorem 1.17. Finally, in Section 9, we study analogs of Catalan numbers associated to arbitrary positroid varieties.

Acknowledgments. We are grateful to David Speyer for his contributions to this project at the early stages. We also thank Pramod Achar, Sergey Fomin, Alexei Oblomkov, Vic Reiner, and Vivek Shende for stimulating discussions. In addition, we are grateful to Ivan Cherednik, Eugene Gorsky, Wolfgang Soergel, and Minh-Tam Trinh for their comments on the first version of this manuscript. Finally, we thank the anonymous referees for their extremely careful reading of our manuscript and for suggesting numerous improvements.

2. POINT COUNT AND THE HOMFLY POLYNOMIAL

2.1. Type A. Let $W = S_n$ and $G = \mathrm{PGL}_n(\mathbb{C})$. Recall from [KL79, Lemmas A3 and A4] that the number of \mathbb{F}_q -points of a Richardson variety $R_{v,w}^\circ$ is given by the *Kazhdan–Lusztig R -polynomial* $R_{v,w}(q)$. When $v \not\leq w$, we have $R_{v,w}(q) = 0$ and $R_{v,w}^\circ = \emptyset$, and for $v = w$, we have $R_{v,w}(q) = 1$ and $R_{v,w}^\circ = \mathrm{pt}$. For $v \leq w \in W$, $R_{v,w}(q)$ can then be computed by a recurrence relation [KL79, Section 2]:

$$(2.1) \quad R_{v,w}(q) = \begin{cases} R_{sv,sw}(q), & \text{if } sv < v \text{ and } sw < w, \\ (q-1)R_{sv,w}(q) + qR_{sv,sw}(q), & \text{if } sv > v \text{ and } sw < w. \end{cases}$$

Here, $s = s_i$ for some $1 \leq i \leq n-1$ is a simple transposition satisfying $sw < w$.

Recall from Section 1.7 that we associate an n -strand braid $\beta_{v,w} := \beta(w) \cdot \beta(v)^{-1}$ to any pair $v, w \in S_n$ of permutations. For a Laurent polynomial $P = P(a, z)$, we denote by $\deg_a^{\mathrm{top}}(P) \in \mathbb{Z}$ the maximal degree of a in P , and for $\kappa \in \mathbb{Z}$, we let $[a^\kappa]P \in \mathbb{C}[z^{\pm 1}]$ be the coefficient of a^κ in P . For $v, w \in S_n$, recall that we set $\ell_{v,w} := \ell(w) - \ell(v)$. Denote

$$(2.2) \quad \kappa_{v,w} := n - 1 - \ell_{v,w} \quad \text{and} \quad P_{v,w} = P_{v,w}(a, z) := P(\hat{\beta}_{v,w}; a, z),$$

where $P(\hat{\beta}_{v,w}; a, z)$ is the HOMFLY polynomial defined in Section 1.6. The goal of this section is to show the following strengthening of Theorem 1.13.

Theorem 2.1. *Let $v, w \in S_n$.*

- (i) *If $v \not\leq w$ then $\deg_a^{\mathrm{top}}(P_{v,w}) < \kappa_{v,w}$.*
- (ii) *If $v \leq w$ then $\deg_a^{\mathrm{top}}(P_{v,w}) = \kappa_{v,w}$.*
- (iii) *For any $v, w \in S_n$, let $P_{v,w}^{\mathrm{top}}(q)$ be obtained from $a^{\kappa_{v,w}} \cdot ([a^{\kappa_{v,w}}]P_{v,w})$ by substituting $a := q^{-\frac{1}{2}}$ and $z := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$. Then*

$$(2.3) \quad R_{v,w}(q) = (q-1)^{n-1} \cdot P_{v,w}^{\mathrm{top}}(q).$$

Proof. We start by recalling the following result, which states that the (lower) Morton–Franks–Williams inequality [Mor86, FW87] is not sharp for negative braids. It may be alternatively deduced from (2.10) below.

Lemma 2.2 ([GMM14, Proposition 2.1]). *Let $v \in S_n$ be a non-identity permutation and let $\beta := \beta(v)^{-1}$ be the associated negative braid. Then⁵*

$$(2.4) \quad \deg_a^{\mathrm{top}}(P(\hat{\beta}; a, z)) < n - 1 + \ell(v). \quad \square$$

We now prove all parts of Theorem 2.1 by induction on $\ell(w)$. Consider the base case $\ell(w) = 0$. Then (i) is the content of (2.4). For (ii), we observe that $\ell(w) = 0$ implies $v = w = \mathrm{id}$, and iterating Example 1.12, we get $P_{v,w} = \left(\frac{a-a^{-1}}{z}\right)^{n-1}$. Thus, $\deg_a^{\mathrm{top}}(P_{v,w}) = n-1 = \kappa_{v,w}$ for $v = w = \mathrm{id}$. For (iii), if $v \not\leq w$ then by (i), we get $[a^{\kappa_{v,w}}]P_{v,w} = 0$, so $P_{v,w}^{\mathrm{top}}(q) = 0$, in agreement with $R_{v,w}(q) = 0$. If $v \leq w$ then $v = w = \mathrm{id}$, $a^{\kappa_{v,w}} \cdot [a^{\kappa_{v,w}}]P_{v,w} = (a/z)^{n-1}$, so $P_{v,w}^{\mathrm{top}}(q) = (q-1)^{-(n-1)}$, in agreement with (2.3). We have shown the base case for each part.

For the induction step, suppose that $\ell(w) > 0$. Choose some $1 \leq i \leq n-1$ such that $s_i w < w$ and let $s := s_i$ and $\sigma := \sigma_i$. If $sv < v$ then the links $\hat{\beta}_{v,w}$ and $\hat{\beta}_{sv,sw}$ are isotopic since $\beta_{v,w} = \sigma \beta_{sv,sw} \sigma^{-1}$, and thus $P_{v,w} = P_{sv,sw}$. We also have $\kappa_{v,w} = \kappa_{sv,sw}$, and thus $P_{v,w}^{\mathrm{top}}(q) = P_{sv,sw}^{\mathrm{top}}(q)$. By (2.1), $R_{v,w}(q) = R_{sv,sw}(q)$. So in the case $sv < v$, the induction step holds trivially for each of the three parts.

⁵Our conventions for $P(\hat{\beta}; a, z)$ differ from those of [GMM14] by changing $a \mapsto a^{-1}$.

Assume now that we have $sw < w$ and $sv > v$. In this case, we have $\beta_{v,w} = \sigma\beta(sw)\beta(sv)^{-1}\sigma \sim \beta(sw)\beta(sv)^{-1}\sigma^2$, $\beta_{v,sw} = \beta(sw)\beta(sv)^{-1}\sigma$, and $\beta_{sv,sw} = \beta(sw)\beta(sv)^{-1}$, where \sim relates conjugate braids. Applying (1.15) with

$$L_+ := \hat{\beta}_{v,w}, \quad L_0 := \hat{\beta}_{v,sw}, \quad L_- := \hat{\beta}_{sv,sw},$$

we get $aP_{v,w} - a^{-1}P_{sv,sw} = zP_{v,sw}$, and thus

$$(2.5) \quad P_{v,w} = \frac{z}{a}P_{v,sw} + a^{-2}P_{sv,sw}.$$

Note that $\kappa_{v,sw} = \kappa_{v,w} + 1$ and $\kappa_{sv,sw} = \kappa_{v,w} + 2$. Let us show (i). We have $v \not\leq w$, $sw < w$, and $sv > v$, and thus clearly $v \not\leq sw$ and $sv \not\leq sw$. By the induction hypothesis, we have $\deg_a^{\text{top}}(P_{v,sw}) < \kappa_{v,sw}$ and $\deg_a^{\text{top}}(P_{sv,sw}) < \kappa_{sv,sw}$. By (2.5), we get $\deg_a^{\text{top}}(P_{v,w}) < \kappa_{v,w}$, finishing the proof of (i). In particular, we have shown that (2.3) holds for all $v \not\leq w$.

Now assume $v \leq w$. We show (ii) and (iii) simultaneously. By the induction hypothesis, we have $\deg_a^{\text{top}}(P_{v,sw}) \leq \kappa_{v,sw}$, $\deg_a^{\text{top}}(P_{sv,sw}) \leq \kappa_{sv,sw}$ (whether the equality holds depends on whether $v \leq sw$ and $sv \leq sw$). Thus, by (2.5), $\deg_a^{\text{top}}(P_{v,w}) \leq \kappa_{v,w}$. The links $L_0 = \hat{\beta}_{v,sw}$ and $\hat{\beta}_{sv,w}$ are isotopic since $\beta_{sv,w} = \sigma\beta(sw)\beta(sv)^{-1} \sim \beta(sw)\beta(sv)^{-1}\sigma = \beta_{v,sw}$, so $P_{v,sw} = P_{sv,w}$. Applying the map $P \mapsto a^{\kappa_{v,w}} \cdot ([a^{\kappa_{v,w}}]P)$ to both sides of (2.5) and substituting $a := q^{-\frac{1}{2}}$ and $z := q^{\frac{1}{2}} - q^{-\frac{1}{2}}$, we get

$$P_{v,w}^{\text{top}}(q) = (q-1)P_{sv,w}^{\text{top}}(q) + qP_{sv,sw}^{\text{top}}(q).$$

Combining this with the induction hypothesis and (2.1), we get $R_{v,w}(q) = (q-1)^{n-1} \cdot P_{v,w}^{\text{top}}(q)$. In particular, the coefficient of $a^{\kappa_{v,w}}$ in $P_{v,w}$ is nonzero, so $\deg_a^{\text{top}}(P_{v,w}) = \kappa_{v,w}$. Thus, we have completed the induction step for both (ii) and (iii). \square

2.2. Arbitrary type. The above connection between point counts and the HOMFLY polynomial can be generalized to arbitrary Weyl groups as follows. Let \mathcal{H} be the *Hecke algebra* of W : it is generated over $\mathbb{C}[q^{\pm 1}]$ by the elements $\{T_s\}_{s \in S}$ satisfying the braid relations as well as the Hecke relation

$$(2.6) \quad (T_s + q)(T_s - 1) = 0 \quad \text{for } s \in S.$$

The algebra \mathcal{H} admits a linear basis $\{T_w\}_{w \in W}$ indexed by the elements of W : we set $T_w := T_{s_1}T_{s_2} \cdots T_{s_{\ell(w)}}$ for any reduced word $w = s_1s_2 \cdots s_{\ell(w)}$. The *standard trace* $\epsilon : \mathcal{H} \rightarrow \mathbb{C}[q^{\pm 1}]$ is the $\mathbb{C}[q^{\pm 1}]$ -linear map defined by

$$(2.7) \quad \epsilon(T_w) := \begin{cases} 1, & \text{if } w = \text{id}; \\ 0, & \text{otherwise.} \end{cases}$$

Theorem 2.3. *For any $v, w \in W$, we have*

$$(2.8) \quad R_{v,w}(q) = q^{\ell_{v,w}} \epsilon(T_w^{-1}T_v).$$

For $W = S_n$, in view of the well-known relation between traces and the HOMFLY polynomial (going back to [Jon87]), Theorem 2.3 specializes to Theorem 2.1.

Proof. First, we state a simple consequence of (2.6): for any $v \in W$ and $s \in S$, we have

$$(2.9) \quad T_s T_v = \begin{cases} T_{sv}, & \text{if } sv > v, \\ (1-q)T_v + qT_{sv} & \text{if } sv < v. \end{cases}$$

Next, we claim that for any $u, v \in W$, we have

$$(2.10) \quad \epsilon(T_u T_v) = \begin{cases} q^{\ell(v)}, & \text{if } u = v^{-1}, \\ 0, & \text{otherwise.} \end{cases}$$

We prove (2.10) by induction on $\ell(u)$. The base case $\ell(u) = 0$ is clear. Otherwise, choose $s \in S$ such that $u = xs$ with $x < xs$. If $v < sv$ then $T_u T_v = T_x T_{sv}$ and we are done by induction. Thus, assume $v > sv$. By (2.9), we get

$$(2.11) \quad \epsilon(T_u T_v) = (1 - q)\epsilon(T_x T_v) + q\epsilon(T_x T_{sv}).$$

We have $u = v^{-1}$ if and only if $x = (sv)^{-1}$, in which case by induction we find $\epsilon(T_u T_v) = q^{\ell(v)}$. If $u \neq v^{-1}$ then the right-hand side of (2.11) is zero unless $x = v^{-1}$. But $x = v^{-1}$ contradicts our assumptions $x < xs$ and $v > sv$. This completes the proof of (2.10).

It is well known that $T_w^{-1} \in \text{Span}\{T_u\}_{u \leq w}$. Thus, by (2.10), $\epsilon(T_w^{-1} T_v) = 0$ unless $v \leq w$. We now proceed to prove (2.8) by induction on $\ell_{v,w}$. For $v = w$, the result again follows from (2.10). For $v < w$, we choose $s \in S$ such that $sw < w$ and then calculate using $T_s^{-1} = q^{-1}T_s + (1 - q^{-1})$ that

$$T_w^{-1} T_v = \begin{cases} T_{sw}^{-1} T_{sv}, & \text{if } sv < v \text{ and } sw < w, \\ q^{-1} T_{sw}^{-1} T_{sv} + (1 - q^{-1}) T_{sw}^{-1} T_v, & \text{if } sv > v \text{ and } sw < w. \end{cases}$$

Applying ϵ and multiplying both sides by $q^{\ell_{v,w}}$, the result matches perfectly with (2.1). \square

Remark 2.4. Theorem 2.3 may also be deduced from Theorem 1.17 by taking the Euler characteristic: Soergel bimodules categorify the Hecke algebra, with Rouquier complexes $F^\bullet(w)$ corresponding to the elements T_w , and the zeroth Hochschild cohomology functor HH^0 categorifies the trace ϵ .

3. SOERGEL BIMODULES, ROUQUIER COMPLEXES, AND KHOVANOV–ROZANSKY HOMOLOGY

In this section, we review Khovanov–Rozansky (KR) link homology. A friendly introduction to most of this material can be found in the excellent recent book [EMTW20].

3.1. Soergel bimodules. Let $R := \mathbb{C}[\mathfrak{h}]$ be as in Section 1.9. It is a graded ring where we set $\deg(y) = 2$ for $y \in \mathfrak{h}^*$. We refer to $\deg(y)$ as the *polynomial degree* (as opposed to the cohomological degree introduced later on). The Weyl group W acts naturally on R . Denote by I the indexing set of simple roots of R , and thus W is generated by the simple reflections $S = \{s_i\}_{i \in I}$. When $G = \text{PGL}_n(\mathbb{C})$, recall that $R = \mathbb{C}[y_1, \dots, y_{n-1}]$ is the polynomial ring and $S = \{s_1, \dots, s_{n-1}\}$ is the set of simple transpositions in $W = S_n$. The action of S_n on R is obtained by restricting the permutation action on $\mathbb{C}[x_1, x_2, \dots, x_n]$ to $R \subset \mathbb{C}[x_1, x_2, \dots, x_n]$, where we identify $y_i = x_i - x_{i+1}$ for $1 \leq i \leq n - 1$. For example,

$$(3.1) \quad s_1(y_1) = -y_1, \quad s_2(y_1) = y_1 + y_2, \quad s_3(y_1) = \dots = s_{n-1}(y_1) = y_1.$$

Soergel bimodules are special kinds of *graded R -bimodules*, that is, graded \mathbb{C} -vector spaces equipped with a left and a right graded action of R . For a graded R -bimodule $B = \bigoplus_i B^i$ and $m \in \mathbb{Z}$, we denote by $B\{m/2\} := \bigoplus_i B^{i-m}$ the polynomial grading shift by m on B . Thus, $y \in \mathfrak{h}^*$ has degree 2 as an element of R but has degree 0 as an element of $R\{-1\}$.

Let us introduce the “building blocks” of Soergel bimodules.

Definition 3.1. For $s \in S$, let $R^s := \{r \in R \mid sr = r\}$. Define

$$(3.2) \quad B_s := R \otimes_{R^s} R.$$

For a sequence $\underline{u} = (s_{i_1}, s_{i_2}, \dots, s_{i_m})$ of elements of S , let

$$(3.3) \quad B_{\underline{u}} := B_{s_{i_1}} \otimes_R B_{s_{i_2}} \otimes_R \dots \otimes_R B_{s_{i_m}} = R \otimes_{R^{s_{i_1}}} R \otimes_{R^{s_{i_2}}} \dots \otimes_{R^{s_{i_m}}} R.$$

Both B_s and $B_{\underline{u}}$ are naturally graded R -bimodules, called *Bott–Samelson bimodules*, where R acts on the leftmost and the rightmost terms of the tensor product by multiplication.

We let $\mathbb{S}\text{Bim}$ denote the *category of Soergel bimodules*. By definition, its objects are graded shifts of direct summands of Bott–Samelson bimodules $B_{\underline{u}}$, where \underline{u} runs over all finite sequences

of elements in S . The morphisms in $\mathbb{S}\text{Bim}$ are given by degree 0 maps of R -bimodules. The indecomposable objects $\{S_w\}_{w \in W}$ of $\mathbb{S}\text{Bim}$ are indexed by the elements of W : for each $w \in W$ and any reduced word \underline{w} for w , $B_{\underline{w}}$ contains a unique indecomposable summand S_w that is not contained in $B_{\underline{v}}$ for any $v < w$ and any reduced word \underline{v} for v . Up to isomorphism, the bimodule S_w depends only on w and not on the choice of \underline{w} .

3.2. Rouquier complexes. We let $\text{K}^b\mathbb{S}\text{Bim}$ denote the homotopy category of $\mathbb{S}\text{Bim}$. Its objects are bounded cochain complexes $C^\bullet = (\cdots \rightarrow C^{-1} \rightarrow \underline{C^0} \rightarrow C^1 \rightarrow \cdots)$ of Soergel bimodules, and morphisms are homotopy classes of maps of complexes. When depicting a complex, we usually omit some of the zeroes and underline the object that is in cohomological degree 0. For example, a complex with only two non-zero entries may be written as $(\underline{C^0} \rightarrow C^1)$ or as $(0 \rightarrow \underline{C^0} \rightarrow C^1)$. We denote by $[m]$ the *cohomological shift* on $\text{K}^b\mathbb{S}\text{Bim}$. It shifts each cochain complex m steps to the left: $C^\bullet[1] = (\cdots \rightarrow C^{-1} \rightarrow C^0 \rightarrow \underline{C^1} \rightarrow \cdots)$.

The tensor product $C^\bullet \otimes_R D^\bullet$ of two cochain complexes is the total complex of a double complex whose entries are $C^i \otimes_R D^j$ for $i, j \in \mathbb{Z}$. The sign of the map $C^i \otimes_R D^j \rightarrow C^i \otimes_R D^{j+1}$ is the negation of the obvious one for all even i ; the differential of the resulting total complex squares to zero.

Since (W, S) is a Coxeter system, we can consider the associated Artin braid group \mathcal{B}_W generated by $\{\sigma_i\}_{i \in I}$. The following construction is due to Rouquier [Rou04].

Definition 3.2. For $s = s_i \in S$ and $\sigma = \sigma_i$, define the *Rouquier complexes*

$$(3.4) \quad F^\bullet(\sigma) := (B_s \rightarrow \underline{R}), \quad F^\bullet(\sigma^{-1}) := (\underline{R} \rightarrow B_s\{-1\}),$$

where the first map sends $f \otimes g \mapsto fg$ and the second map sends $1 \mapsto (\alpha_s \otimes 1 + 1 \otimes \alpha_s)$. Here $\alpha_s \in \mathfrak{h}^*$ is the simple root corresponding to s . Note that both $1 \in R$ and $(\alpha_s \otimes 1 + 1 \otimes \alpha_s) \in B_s\{-1\}$ have polynomial degree 0. For a braid $\beta = \sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_m} \in \mathcal{B}_W$, we set

$$\begin{aligned} F^\bullet(\beta) &:= F^\bullet(\sigma_{i_1}) \otimes_R F^\bullet(\sigma_{i_2}) \otimes_R \cdots \otimes_R F^\bullet(\sigma_{i_m}), \\ F^\bullet(\beta^{-1}) &= F^\bullet(\beta)^{-1} := F^\bullet(\sigma_{i_m}^{-1}) \otimes_R \cdots \otimes_R F^\bullet(\sigma_{i_2}^{-1}) \otimes_R F^\bullet(\sigma_{i_1}^{-1}). \end{aligned}$$

We also let $F^\bullet(\text{id}) := (0 \rightarrow \underline{R} \rightarrow 0)$.

A priori, the complex $F^\bullet(\beta)$ depends on the choice the word $(\sigma_{i_1}, \sigma_{i_2}, \dots, \sigma_{i_m})$. However, modulo homotopy, it does not.

Proposition 3.3 ([Rou04, Section 3]). *If $\sigma_{i_1} \sigma_{i_2} \cdots \sigma_{i_m} = \sigma_{i'_1} \sigma_{i'_2} \cdots \sigma_{i'_m}$ in \mathcal{B}_W then*

$$F^\bullet(\sigma_{i_1}) \otimes_R F^\bullet(\sigma_{i_2}) \otimes_R \cdots \otimes_R F^\bullet(\sigma_{i_m}) \cong F^\bullet(\sigma_{i'_1}) \otimes_R F^\bullet(\sigma_{i'_2}) \otimes_R \cdots \otimes_R F^\bullet(\sigma_{i'_m})$$

in $\text{K}^b\mathbb{S}\text{Bim}$.

For example, one can check that $F^\bullet(\sigma_i) \otimes_R F^\bullet(\sigma_i^{-1}) \cong F^\bullet(\text{id})$. It follows that the functors $(-) \otimes_R F^\bullet(\sigma_i)$ and $(-) \otimes_R F^\bullet(\sigma_i^{-1})$ are mutually inverse biadjoint equivalences of categories: for complexes $C^\bullet, D^\bullet \in \text{K}^b\mathbb{S}\text{Bim}$, we have

$$(3.5) \quad \text{Hom}_{\text{K}^b\mathbb{S}\text{Bim}}(C^\bullet, D^\bullet \otimes_R F^\bullet(\sigma_i)) \cong \text{Hom}_{\text{K}^b\mathbb{S}\text{Bim}}(C^\bullet \otimes_R F^\bullet(\sigma_i^{-1}), D^\bullet),$$

$$(3.6) \quad \text{Hom}_{\text{K}^b\mathbb{S}\text{Bim}}(C^\bullet, D^\bullet \otimes_R F^\bullet(\sigma_i^{-1})) \cong \text{Hom}_{\text{K}^b\mathbb{S}\text{Bim}}(C^\bullet \otimes_R F^\bullet(\sigma_i), D^\bullet).$$

Proposition 3.3 allows one to define $F^\bullet(\beta) \in \text{K}^b\mathbb{S}\text{Bim}$ unambiguously for any braid $\beta \in \mathcal{B}_W$. Recall that we are interested in the braid $\beta_{v,w} = \beta(w) \cdot \beta(v)^{-1}$, which corresponds to the complex

$$(3.7) \quad F_{v,w}^\bullet := F^\bullet(\beta_{v,w}) = F^\bullet(\beta(w)) \otimes_R F^\bullet(\beta(v)^{-1}).$$

3.3. KR homology. Recall from (1.17) that the functor HH^0 sends a graded R -bimodule B to the graded R -module $HH^0(B)$ of its R -invariants. Alternatively, it can be expressed as

$$(3.8) \quad HH^0(B) = \bigoplus_{r \in \mathbb{Z}} \text{Hom}_{\mathbb{S}\text{Bim}}(R, B\{-r/2\}).$$

Remark 3.4. The R -module $HH^0(B)$ is free for any Bott-Samelson bimodule B . One can make explicit combinatorial computations with this R -module (including finding a basis and computing the maps in the Rouquier complexes) using the *diagrammatic calculus* developed by Elias–Williamson [EW16].

Remark 3.5. Higher Hochschild cohomology functors HH^h , which give the full (triply-graded) KR homology, are the right derived functors of HH^0 . They can be computed using a Koszul resolution of R ; see e.g. [Kho07, Mel17].

Applying the functor HH^0 to a complex C^\bullet of Soergel bimodules yields a complex $HH^0(C^\bullet)$ of graded R -modules. In particular, for each $k \in \mathbb{Z}$, the cohomology $H^k(HH^0(C^\bullet))$ of this complex is a graded R -module. For $r \in \mathbb{Z}$, we denote by $H^{k,(r/2)}(HH^0(C^\bullet))$ its graded piece of polynomial degree r . It is not hard to check that we have

$$(3.9) \quad H^{k,(r/2)}(HH^0(C^\bullet)) \cong \text{Hom}_{\mathbb{K}^b\mathbb{S}\text{Bim}}(R, C^\bullet[k]\{-r/2\}).$$

For any $\beta \in \mathcal{B}_W$, $F^\bullet(\beta)$ is concentrated in even polynomial degrees, and thus $H^{k,(r/2)}(HH^0(F^\bullet(\beta)))$ vanishes when r is odd. Similarly to (1.19), we denote

$$(3.10) \quad HHH^0(F^\bullet(\beta)) := \bigoplus_{k,p \in \mathbb{Z}} H^{k,(p)}(HH^0(F^\bullet(\beta))).$$

The complex $F_{v,w}^\bullet$ is concentrated in cohomological degrees $-\ell(w), -\ell(w) + 1, \dots, \ell(v)$, and thus $H^{k,(p)}(HH^0(F_{v,w}^\bullet)) = 0$ unless $-\ell(w) \leq k \leq \ell(v)$, and the index $p \in \mathbb{Z}$ is bounded from below.

The functor $HH_{\mathbb{C}}^0$ admits a similar description. Recall that \mathbb{C} is considered an R -bimodule on which \mathfrak{h}^* acts by zero on both sides. Any Soergel bimodule $B \in \mathbb{S}\text{Bim}$ gives rise to a *Soergel module* $B \otimes_R \mathbb{C}$, which is a graded R -module. We let $\mathbb{S}\text{Mod}$ denote the category of Soergel modules (with morphisms being maps of polynomial degree 0). By a result of Soergel [Soe92] (see [EMTW20, Proposition 15.27]), for any $B, B' \in \mathbb{S}\text{Bim}$, we have a natural isomorphism

$$\text{Hom}_{\mathbb{S}\text{Bim}}(B, B') \otimes_R \mathbb{C} \xrightarrow{\sim} \text{Hom}_{\mathbb{S}\text{Mod}}(B \otimes_R \mathbb{C}, B' \otimes_R \mathbb{C}).$$

Applying this to the case $B = R$, we get the following result.

Corollary 3.6. *For any Soergel bimodule $B \in \mathbb{S}\text{Bim}$, we have*

$$HH_{\mathbb{C}}^0(B) \cong \text{Hom}_{\mathbb{S}\text{Mod}}(\mathbb{C}, B \otimes_R \mathbb{C}).$$

3.4. Link components and R -module structure. For this section, we assume $W = S_n$. Let β be a braid and $u \in S_n$ the image of β . Let \tilde{R} denote the polynomial ring $\mathbb{C}[x_1, x_2, \dots, x_n]$, and let $\tilde{F}^\bullet(\beta)$ denote the Rouquier complex using \tilde{R} instead of R ; cf. (3.1). Thus, $F^\bullet(\beta)$ is a reduced version of $\tilde{F}^\bullet(\beta)$. The complex $\tilde{F}^\bullet(\beta)$ is an $\tilde{R} \otimes \tilde{R}$ module, and it is known [Ras07, GH22] that the action of $x_i \otimes 1$ is homotopic to the action of $1 \otimes x_{u(i)}$. Indeed, it follows from [GH22, Proposition 2.11 and Theorem 2.18], that there exist cochain maps ξ_i (of polynomial degree 2 and cohomological degree -1) called *dot-sliding homotopies* such that

- (1) $d\xi_i + \xi_i d = x_i \otimes 1 - 1 \otimes x_{u(i)}$ for $i = 1, 2, \dots, n$, and
- (2) $\xi_i \xi_j + \xi_j \xi_i = 0$ for $i, j = 1, 2, \dots, n$.

In $HH^0(\tilde{F}^\bullet(\beta))$, the two R -actions are equalized, so $d\xi_i + \xi_i d = x_i - x_{u(i)}$. Thus, the actions of x_i and x_j on $HH^0(\tilde{F}^\bullet(\beta))$ agree when i and j belong to the same component of the link $\hat{\beta}$. Working instead with the smaller polynomial ring $R = \mathbb{C}[x_1 - x_2, \dots, x_{n-1} - x_n] \subset \tilde{R}$, we deduce that $x_i - x_j$ acts as 0 on $HH^0(F^\bullet(\beta))$ when i, j belong to the same component of the link $\hat{\beta}$. In particular, if $\hat{\beta}$ is a knot, then the action of R on $HH^0(F^\bullet(\beta))$ factors through the natural map $R \rightarrow \mathbb{C}$.

Suppose now that $\hat{\beta}$ is a knot. Denote $z_i := x_i - x_{u(i)}$ for $i = 1, 2, \dots, n$. Thus, $R = \mathbb{C}[z_1, \dots, z_{n-1}]$. Recall that $HH^0(F^\bullet(\beta))$ is a complex of graded, free (cf. Remark 3.4) R -modules. Let $a \in HH^0(F^\bullet(\beta))$ be a nonzero element satisfying $d(a) = 0$. Using the relation $d\xi_i + \xi_i d = z_i$, one can show by induction on $k = 0, 1, \dots, n-1$ that for all $1 \leq i_1 < i_2 < \dots < i_k \leq n-1$, we have

$$(3.11) \quad d\xi_{i_1 i_2 \dots i_k}(a) = \sum_{j=1}^k (-1)^{j-1} z_{i_j} \xi_{i_1 \dots \hat{i}_j \dots i_k}(a),$$

where $\xi_{i_1 i_2 \dots i_k} := \xi_{i_1} \xi_{i_2} \dots \xi_{i_k}$, and \hat{i}_j denotes omission of ξ_{i_j} . We therefore obtain a subcomplex $K^\bullet(a)$ of $HH^0(F^\bullet(\beta))$ given by

$$\begin{aligned} R \cdot \xi_{1 2 \dots n-1}(a) &\rightarrow \dots \rightarrow \sum_{1 \leq i_1 < \dots < i_k \leq n-1} R \cdot \xi_{i_1 \dots i_k}(a) \rightarrow \dots \\ &\rightarrow \sum_{1 \leq i \leq n-1} R \cdot \xi_i(a) \rightarrow R \cdot a. \end{aligned}$$

Definition 3.7. Let $a \in HH^0(F^\bullet(\beta))$ be such that $d(a) = 0$. We say that $K^\bullet(a)$ is a *Koszul subcomplex* if the set $\{\xi_{i_1 \dots i_k}(a) \mid 0 \leq k \leq n-1, 1 \leq i_1 < \dots < i_k \leq n-1\}$ can be extended to a free R -module basis of $HH^0(F^\bullet(\beta))$.

It follows from (3.11) that we have a natural cochain map $\bigotimes_{i=1}^{n-1} (R \xrightarrow{z_i} R) \rightarrow HH^0(F^\bullet(\beta))$ with image $K^\bullet(a)$, and this map is an isomorphism when $K^\bullet(a)$ is a Koszul subcomplex.

Our next goal is to show that $HH^0(F^\bullet(\beta))$ admits a filtration by Koszul subcomplexes and contractible subcomplexes of the form $R \xrightarrow{\sim} R$.

Definition 3.8. We say that a complex (C^\bullet, d) of finite rank, free, graded R -modules *admits a \wedge -action* if there exist endomorphisms $\xi_1, \xi_2, \dots, \xi_{n-1}$ of cohomological degree -1 and polynomial degree 2 satisfying $d\xi_i + \xi_i d = z_i$ and $\xi_i \xi_j + \xi_j \xi_i = 0$, for all $i, j = 1, 2, \dots, n-1$.

We thank the anonymous referee for suggesting to us that the following statement may be deduced from the results of [GH22].

Proposition 3.9. *Suppose that (C^\bullet, d) admits a \wedge -action. Then C^\bullet has a filtration by Koszul complexes and trivial complexes $R \cong R$. That is, there exists a family of subcomplexes $0 = F_0^\bullet \subset F_1^\bullet \subset \dots \subset F_t^\bullet = C^\bullet$ such that for all $j = 1, 2, \dots, t$, $C^\bullet/F_{j-1}^\bullet$ is free, admits a \wedge -action, and $F_j^\bullet/F_{j-1}^\bullet$ is either a Koszul subcomplex of $C^\bullet/F_{j-1}^\bullet$ or a trivial subcomplex isomorphic to $R \xrightarrow{\sim} R$.*

Proof. Let $C^{k, (p/2)}$ denote the subspace of C^\bullet of cohomological degree k and polynomial degree p . Assuming C^\bullet is nonzero, let $D^\bullet \subset C^\bullet$ be sum of those nonzero pieces $C^{k, (p/2)}$ where $k + p/2$ is minimal.

Suppose $a \in D^\bullet$ satisfies $d(a) = 0$. Then for any i_1, \dots, i_k we have that $\xi_{i_1 \dots i_k}(a) \in D^\bullet$. Any linearly independent (over \mathbb{C}) elements in D^\bullet can be extended to a free R -module basis of C^\bullet . Thus, using (3.11), one can show by induction on $k = 0, 1, \dots, n-1$ that the elements $\{\xi_{i_1 \dots i_k}(a) \mid 1 \leq i_1 < \dots < i_k \leq n-1\}$ are linearly independent. It follows that $K^\bullet(a)$ is a Koszul subcomplex of C^\bullet , and furthermore that the quotient by this complex is again free and admits a \wedge -action.

Repeating this, we may assume that $d|_{D^\bullet}$ is injective. We claim that for any nonzero element $b \in D^\bullet$, $d(b)$ may be completed to a free basis of C^\bullet . Suppose that $b \in D^{k-1}$. We proceed by inverse induction on k . For the base case, if $D^k = 0$ then $d(b)$ is a \mathbb{C} -linear combination of free basis elements, so the statement follows. For the induction step, suppose that $b \in D^{k-1}$ satisfies $d(b) \in R \cdot D^k$. Write $d(b) = \sum_{i=1}^{n-1} y_i e_i$ and let i be such that $e_i \neq 0$. By the induction hypothesis, $f := d(e_i)$ may be extended to a free basis of C^{k+1} , and we find that the coefficient of $y_i f$ in $d^2(b)$ is nonzero, a contradiction. Thus, $d(b) \notin R \cdot D^k$. Comparing the polynomial degree of $d(b)$ to that of D^k , we get that $d(b)$ can be extended to an R -basis of C^k .

A subcomplex $R \xrightarrow{\sim} R$ in C^\bullet is called *splittable* if the quotient by this subcomplex again consists of free R -modules. Let k be the smallest index such that $D^k \neq 0$, and let $a_1 \in D^k$ be a nonzero element. We have shown above that $a_0 := d(a_1)$ may be completed to a free basis of C^\bullet , and thus a_1 and a_0 generate a splittable subcomplex $S^\bullet \cong (R \xrightarrow{\sim} R)$. For any $1 \leq i \leq n-1$, we have $\xi_i(a_1) \in D^{k-1} = 0$. Using $d\xi_i + \xi_i d = z_i$, we get $z_i a_1 = \xi_i a_0$. Thus, the subcomplex S^\bullet is closed under the action of ξ_1, \dots, ξ_{n-1} . Using Gaussian elimination [EMTW20, Exercise 19.12], one can check that in this case, the quotient complex C^\bullet/S^\bullet admits a \wedge -action.

Repeating the above procedure, we obtain the desired filtration. \square

Recall that the Koszul resolution of the R -module \mathbb{C} by free R -modules yields a 2^{n-1} -dimensional complex $\mathrm{Tor}_\bullet^R(\mathbb{C}, \mathbb{C}) \cong (\mathbb{C} \xrightarrow{0} \underline{\mathbb{C}})^{\otimes(n-1)}$.

Corollary 3.10. *Suppose that (C^\bullet, d) admits a \wedge -action. Let $(C_\mathbb{C}^\bullet, d_\mathbb{C})$ be the complex of \mathbb{C} -vector spaces obtained by setting $y_1 = y_2 = \dots = y_{n-1} = 0$. Thus, $C_\mathbb{C}^\bullet := C^\bullet \otimes_R \mathbb{C}$ as in (1.18). Then*

$$H^\bullet(C_\mathbb{C}^\bullet) \cong \mathrm{Tor}_\bullet^R(\mathbb{C}, H^\bullet(C^\bullet)) \cong H^\bullet(C^\bullet) \otimes (\mathbb{C} \xrightarrow{0} \underline{\mathbb{C}})^{\otimes(n-1)}$$

(as complexes of graded \mathbb{C} -vector spaces with zero differentials).

Proof. It is clear that the filtration constructed in Proposition 3.9 induces an injection $H^\bullet(F_j^\bullet/F_{j-1}^\bullet) \hookrightarrow H^\bullet(C^\bullet/F_{j-1}^\bullet)$, and a similar statement holds after setting $y_1 = y_2 = \dots = y_{n-1} = 0$. Thus, each Koszul complex $K^\bullet(a)$ appearing in the filtration contributes a 1-dimensional subcomplex to $H^\bullet(C^\bullet)$. In view of (3.11), $K^\bullet(a)$ contributes to $H^\bullet(C_\mathbb{C}^\bullet)$ a 2^{n-1} -dimensional subcomplex isomorphic to $(\mathbb{C} \xrightarrow{0} \underline{\mathbb{C}})^{\otimes(n-1)}$. \square

3.5. Link invariant. For this section, we continue to assume $W = S_n$. The above construction may be turned into a link invariant as we now explain. We follow the conventions of [HM19].

For a braid $\beta \in \mathcal{B}_{S_n} = \mathcal{B}_n$, let $e(\beta)$ denote the *exponent sum* of β :

$$\beta = \sigma_{i_1}^{\epsilon_1} \sigma_{i_2}^{\epsilon_2} \cdots \sigma_{i_m}^{\epsilon_m} \implies e(\beta) := \epsilon_1 + \epsilon_2 + \cdots + \epsilon_m.$$

Thus, $e(\beta_{v,w}) = \ell_{v,w} = \ell(w) - \ell(v)$. Next, define

$$(3.12) \quad \chi(\beta) := \frac{e(\beta) - n + c(\beta)}{2}$$

where $c(\beta)$ is the number of components of the link $\hat{\beta}_{v,w}$, which equals the number of cycles of the corresponding permutation (obtained from the group homomorphism $\mathcal{B}_{S_n} \rightarrow S_n$ sending $\sigma_i \mapsto s_i$ for each $1 \leq i \leq n-1$). It is easy to check that $\chi(\beta)$ is always an integer. Define

$$\begin{aligned} \mathcal{P}_{\mathrm{KR}}(\beta; a, q, t) &:= (1-t)^{c(\beta)-1} \left(q^{\frac{1}{2}} t^{-\frac{1}{2}} a^{-2} \right)^{\chi(\beta)} \\ &\quad \times \sum_{k,p,h \in \mathbb{Z}} (-1)^h q^{\frac{k}{2}} t^{p+\frac{k}{2}+h} a^{-2h} \dim H^{k,(p)}(\mathrm{HH}^h(F^\bullet(\beta))). \end{aligned}$$

Let $\mathcal{P}_{\mathrm{KR}}^{\mathrm{top}}(\beta; q, t)$ be its top a -degree coefficient:

$$(3.13) \quad \mathcal{P}_{\mathrm{KR}}^{\mathrm{top}}(\beta; q, t) := [a^{\mathrm{deg}_a^{\mathrm{top}}(\mathcal{P}_{\mathrm{KR}}(\beta))}] \mathcal{P}_{\mathrm{KR}}(\beta).$$

Theorem 3.11 ([Kho07]). \mathcal{P}_{KR} and $\mathcal{P}_{\text{KR}}^{\text{top}}$ are link invariants: if $\beta \in \mathcal{B}_{S_n}, \beta' \in \mathcal{B}_{S_n}$ are two braids such that the corresponding links $\hat{\beta} \cong \hat{\beta}'$ are isotopic then

$$\mathcal{P}_{\text{KR}}(\beta; a, q, t) = \mathcal{P}_{\text{KR}}(\beta'; a, q, t) \quad \text{and} \quad \mathcal{P}_{\text{KR}}^{\text{top}}(\beta; q, t) = \mathcal{P}_{\text{KR}}^{\text{top}}(\beta'; q, t)$$

Thus, it makes sense to write $\mathcal{P}_{\text{KR}}(\hat{\beta}; a, q, t) := \mathcal{P}_{\text{KR}}(\beta; a, q, t)$ and $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t) := \mathcal{P}_{\text{KR}}^{\text{top}}(\beta; q, t)$. Both $\mathcal{P}_{\text{KR}}(\hat{\beta}; a, q, t)$ and $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$ have been recently shown to be q, t -symmetric when $\hat{\beta}$ is a knot [OR20].

It is well known that $\mathcal{P}_{\text{KR}}(\hat{\beta}; a, q, t)$ specializes to the HOMFLY polynomial of $\hat{\beta}$:

$$(3.14) \quad \mathcal{P}_{\text{KR}}(\hat{\beta}) \Big|_{t^{\frac{1}{2}} = -q^{-\frac{1}{2}}} = (-1)^{\chi(\beta)} (z/a)^{c(\beta)-1} P(\hat{\beta}; a, z) \Big|_{z=q^{\frac{1}{2}}-q^{-\frac{1}{2}}}.$$

Note that $\chi(\beta)$ is *not* a link invariant, but $(-1)^{\chi(\beta)}$ and $c(\beta)$ are link invariants.

Clearly, for any braid β , we have $\deg_a^{\text{top}}(\mathcal{P}_{\text{KR}}(\beta)) \leq -2\chi(\beta)$. Let $v \leq w \in S_n$. Comparing (3.12) with (2.2), we find $\kappa_{v,w} = -2\chi(\beta_{v,w}) + c(\beta_{v,w}) - 1$, and thus the coefficient of $a^{-2\chi(\beta_{v,w})}$ in $\mathcal{P}_{\text{KR}}(\beta_{v,w})$ is nonzero, by (3.14) combined with Theorem 2.1(ii). Therefore $\deg_a^{\text{top}}(\mathcal{P}_{\text{KR}}(\beta_{v,w})) = -2\chi(\beta_{v,w})$, and we get the following result.

Proposition 3.12. For $v \leq w \in S_n$, $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}) = [a^{-2\chi(\beta_{v,w})}] \mathcal{P}_{\text{KR}}(\hat{\beta}_{v,w})$ is given by

$$(3.15) \quad \begin{aligned} \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) &= (1-t)^{c(\beta_{v,w})-1} \left(q^{\frac{1}{2}} t^{-\frac{1}{2}} \right)^{\chi(\beta_{v,w})} \\ &\quad \times \sum_{k,p \in \mathbb{Z}} q^{\frac{k}{2}} t^{p+\frac{k}{2}} \dim H^{k,(p)}(\text{HH}^0(F^\bullet(\beta_{v,w}))). \end{aligned}$$

Let us also define the analogous polynomial in the non-equivariant case (cf. Section 1.8). For $\beta \in \mathcal{B}_{S_n}$, set

$$(3.16) \quad \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) = \frac{\left(q^{\frac{1}{2}} t^{-\frac{1}{2}} \right)^{\chi(\beta)}}{\left(1 + q^{-\frac{1}{2}} t^{\frac{1}{2}} \right)^{n-c(\beta)}} \sum_{k,p \in \mathbb{Z}} q^{\frac{k}{2}} t^{p+\frac{k}{2}} \dim H^{k,(p)}(\text{HH}_{\mathbb{C}}^0(F^\bullet(\beta))).$$

The denominator $\left(1 + q^{-\frac{1}{2}} t^{\frac{1}{2}} \right)^{n-c(\beta)}$ in (3.16) is chosen in view of the discussion in Section 3.4: when β is a link with $c(\beta)$ components, a filtration analogous to the one in Proposition 3.9 would involve complexes with $2^{n-c(\beta)}$ terms.

The following result is a consequence of Corollary 3.10.

Corollary 3.13. Assume that $\hat{\beta}$ is a knot such that $\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) \neq 0$. Then

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\beta; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t).$$

In Section 8.1, we give an alternative proof for knots of the form $\hat{\beta}_{v,w}$ for $v \leq w \in S_n$.

3.6. Examples. We compute $\text{HHH}^0(F^\bullet(\beta))$ and $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$, as well as $\text{HHH}_{\mathbb{C}}^0(F^\bullet(\beta))$ and $\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t)$, for a few braids β . Throughout, we assume $G = \text{PGL}_n(\mathbb{C})$, in which case recall that $R = \mathbb{C}[y_1, \dots, y_{n-1}]$ is a polynomial ring. These examples are summarized in Table 2. We abbreviate \otimes_R by \otimes .

Example 3.14 (Unknot I). Let $n = 1$, $v = \text{id}$, $w = \text{id}$, and thus $c(\beta_{v,w}) = 1$ and $\chi(\beta_{v,w}) = 0$. We have $F^\bullet(\beta_{v,w}) = (0 \rightarrow \underline{R} \rightarrow 0)$ and $R = \mathbb{C}$. Thus, the only nonzero term is $H^{0,(0)}(\text{HH}^0(F_{v,w}^\bullet)) \cong H^{0,(0)}(\text{HH}_{\mathbb{C}}^0(F_{v,w}^\bullet)) \cong \mathbb{C}$. We have

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = 1.$$

H^k	H^0	H^1	H^2	H^3	H^4	H^k	H^0	H^1	H^2	H^3	H^4	H^5	H^6
$k-p=0$	1	3	4	3	1	$k-p=0$	1	4	7	8	7	4	1
$H^{k,(p,p)}(\Pi_{2,4}^\circ)$						$H^{k,(p,p)}(\Pi_{2,5}^\circ)$							
H^k	H^{-4}	H^{-3}	H^{-2}	H^{-1}	H^0	H^k	H^{-6}	H^{-5}	H^{-4}	H^{-3}	H^{-2}	H^{-1}	H^0
$k+p=0$	1	3	4	3	1	$k+p=0$	1	4	7	8	7	4	1
$H^{k,(p)}HH_{\mathbb{C}}^0(F^\bullet(\beta_{f_{2,4}}))$						$H^{k,(p)}HH_{\mathbb{C}}^0(F^\bullet(\beta_{f_{2,5}}))$							

TABLE 3. Comparing the mixed Hodge tables of $\Pi_{2,4}^\circ$ and $\Pi_{2,5}^\circ$ (top) to $HHH_{\mathbb{C}}^0$ of the associated links (bottom).

Note that any $(1, b)$ -torus knot is isotopic to the unknot, and we have $C_{1,b}(q, t) = 1$.

Example 3.15 (Unknot II). Let $n = 2$, $v = \text{id}$, $w = s_1$, and thus $c(\beta_{v,w}) = 1$ and $\chi(\beta_{v,w}) = 0$. We have $F_{v,w}^\bullet = (B_{s_1} \rightarrow \underline{R})$. It is easy to see that $HH^0(B_{s_1})$ is a free R -module spanned by $(y_1 \otimes 1 + 1 \otimes y_1)$, and thus $HH^0(B_{s_1}) \cong R\{1\}$, and $HH^0(F_{v,w}^\bullet) = (R\{1\} \rightarrow \underline{R})$, with the map sending $1 \mapsto 2y_1$. The only nonzero term is $H^{0,(0)}(HH^0(F_{v,w}^\bullet)) \cong \mathbb{C}$. Tensoring with \mathbb{C} , we get $HH_{\mathbb{C}}^0(F_{v,w}^\bullet) = (\mathbb{C}\{1\} \xrightarrow{0} \underline{\mathbb{C}})$, so there are two nonzero terms: $H^{0,(0)}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)) \cong H^{-1,(1)}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)) \cong \mathbb{C}$. Therefore

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = 1 \quad \text{and} \quad \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = \frac{1}{1 + q^{-\frac{1}{2}}t^{\frac{1}{2}}} \left(1 + q^{-\frac{1}{2}}t^{\frac{1}{2}}\right) = 1.$$

Let us also consider an example of $\hat{\beta}_{v,w}$ for $v \not\leq w$.

Example 3.16 (Unknot III). Let $n = 2$, $v = s_1$, $w = \text{id}$, and thus $c(wv^{-1}) = 1$ and $\chi(\beta_{v,w}) = -1$. We have $F_{v,w}^\bullet = (\underline{R} \rightarrow B_{s_1}\{-1\})$, $HH^0(F_{v,w}^\bullet) = (\underline{R} \xrightarrow{\sim} R)$, and $HH_{\mathbb{C}}^0(F_{v,w}^\bullet) = (\underline{\mathbb{C}} \xrightarrow{\sim} \mathbb{C})$ so the right-hand sides of (3.15) and (3.16) are zero:

$$[a^2]\mathcal{P}_{\text{KR}}(\hat{\beta}_{v,w}; a, q, t) = 0 \quad \text{and} \quad \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{v,w}; q, t) = 0.$$

This is consistent with the fact that $\mathcal{P}_{\text{KR}}(\hat{\beta}_{v,w})$ is a link invariant satisfying $\mathcal{P}_{\text{KR}}(\bigcirc) = 1$, therefore by (3.13) we have $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}) = 1$. For a computation of $HH^1(F^\bullet(\beta_{v,w}))$, see [Mel17].

In the next two examples, we have $c(\beta) > 1$. We start with the case of the 2-component unlink $\bigcirc \bigcirc$. It is the closure of $\text{id} \in \mathcal{B}_{S_2}$, but we consider the representative $\beta_{s_1, s_1} = \sigma_1 \sigma_1^{-1}$ instead.

Example 3.17 (2-component unlink). Let $n = 2$, $v = s_1$, $w = s_1$, and thus $c(wv^{-1}) = 2$ and $\chi(\beta_{v,w}) = 0$. We have

$$F^\bullet(\beta_{v,w}) = (B_{s_1} \rightarrow (R \oplus (B_{s_1} \otimes B_{s_1}\{-1\})) \rightarrow B_{s_1}\{-1\}).$$

We apply the well-known (see, e.g., [GKS21, Example 3.12]) Soergel bimodule isomorphism $B_{s_1} \otimes B_{s_1} \cong B_{s_1}\{1\} \oplus B_{s_1}$, sending $1 \otimes 1 \otimes 1 \mapsto (0, 1 \otimes 1)$ and $1 \otimes y_1 \otimes 1 \mapsto (1 \otimes 1, 0)$. Next, we use Gaussian elimination [EMTW20, Exercise 19.12] to obtain $F^\bullet(\beta_{v,w}) \cong F^\bullet(\text{id})$ in $\text{K}^{\text{b}}\text{SBim}$, in agreement with Proposition 3.3. We have $R = \mathbb{C}[y_1]$, $HH^0(R) \cong R$, and $HH_{\mathbb{C}}^0(R) \cong \mathbb{C}$. (More generally, recall from Remark 3.4 that the R -module $HH^0(B)$ is always free.) Therefore the only nonzero terms are $H^{0,(p)}(HH^0(F_{v,w}^\bullet)) \cong \mathbb{C}$ for $p = 0, 1, 2, \dots$, and $H^{0,(0)}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)) \cong \mathbb{C}$. We find

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w}; q, t) = (1-t)(1+t+t^2+\dots) = 1 \quad \text{and} \quad \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{v,w}; q, t) = 1.$$

The Hopf link $\hat{\beta} = \bigcirc \bigcirc$ consists of two linked unknots. It is isotopic to $\hat{\beta}_{f_{2,4}}$, as well as to the closure of $(\sigma_1)^2 \in \mathcal{B}_{S_2}$.

Example 3.18 (Hopf link). Let $n = 2$, $\beta = (\sigma_1)^2$, and thus $c(\beta) = 2$ and $\chi(\beta) = 1$. We have

$$F^\bullet(\beta) = (B_{s_1} \otimes B_{s_1} \rightarrow B_{s_1} \oplus B_{s_1} \rightarrow \underline{R}).$$

Using Gaussian elimination as in Example 3.17, we obtain

$$F^\bullet(\beta) \cong (B_{s_1}\{1\} \rightarrow B_{s_1} \rightarrow \underline{R}).$$

Here, the first map sends $1 \otimes 1 \mapsto y_1 \otimes 1 - 1 \otimes y_1$, and the second map sends $1 \otimes 1 \mapsto 1$. Taking R -invariants (cf. Remark 3.4), we find

$$(3.17) \quad HH^0(F^\bullet(\beta)) = \left(R\{2\} \xrightarrow{0} R\{1\} \xrightarrow{2y_1} \underline{R} \right).$$

We get $H^{0,(0)}(HH^0(F^\bullet(\beta))) \cong \mathbb{C}$ and $H^{-2}(HH^0(F^\bullet(\beta))) \cong R\{2\}$. In other words,

$$(3.18) \quad H^{0,(0)}(HH^0(F^\bullet(\beta))) \cong \mathbb{C}, \quad H^{-2,(p)}(HH^0(F^\bullet(\beta))) \cong \mathbb{C} \quad \text{for } p = 2, 3, 4, \dots$$

Sending $y_1 \rightarrow 0$ in (3.17), we find

$$H^{k,(p)}(HH_{\mathbb{C}}^0(F^\bullet(\beta))) \cong \begin{cases} \mathbb{C}, & \text{if } (k, p) \in \{(0, 0), (-1, 1), (-2, 2)\}; \\ 0, & \text{otherwise.} \end{cases}$$

Thus,

$$\begin{aligned} \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t) &= (1-t)q^{\frac{1}{2}}t^{-\frac{1}{2}}(1+t/q(1+t+t^2+\dots)) \\ &= q^{\frac{1}{2}}t^{-\frac{1}{2}} - q^{\frac{1}{2}}t^{\frac{1}{2}} + q^{-\frac{1}{2}}t^{\frac{1}{2}}, \\ \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) &= q^{\frac{1}{2}}t^{-\frac{1}{2}}(1+q^{-\frac{1}{2}}t^{\frac{1}{2}}+t/q) = q^{\frac{1}{2}}t^{-\frac{1}{2}} + 1 + q^{-\frac{1}{2}}t^{\frac{1}{2}}. \end{aligned}$$

Remark 3.19. Since $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$ is a link invariant and $\hat{\beta} \cong \hat{\beta}_{2,4}$ (with $\chi(\hat{\beta}) = \chi(\hat{\beta}_{2,4})$), we see that $HHH^0(F^\bullet(\beta)) \cong HHH^0(F^\bullet(\beta_{2,4}))$ as bigraded vector spaces. Using an elaborate computation, one can also check that $\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{2,4}; q, t)$. However, observe that (3.16) has $(1+q^{-\frac{1}{2}}t^{\frac{1}{2}})^{n-c(\beta)}$ in the denominator, where $n = 2$ for β and $n = 4$ for $\beta_{2,4}$. Thus, $HHH_{\mathbb{C}}^0(F^\bullet(\beta_{2,4}))$ differs from $HHH_{\mathbb{C}}^0(F^\bullet(\beta))$ by ‘‘multiplication by $(1+q^{-\frac{1}{2}}t^{\frac{1}{2}})^2$,’’ and the actual bigraded dimensions of $HHH_{\mathbb{C}}^0(F^\bullet(\beta_{2,4}))$ are given in Table 3(bottom left).

Remark 3.20. We have a resolution of \mathbb{C} by free R -modules: $0 \rightarrow R \xrightarrow{y_1} R \rightarrow \mathbb{C} \rightarrow 0$. Thus, $\text{Tor}_{\bullet}^R(\mathbb{C}, \mathbb{C}) = (\mathbb{C} \xrightarrow{0} \mathbb{C})$. Noting that $\text{Tor}_{\bullet}^R(\mathbb{C}, R) = \mathbb{C}$, we see that $HHH_{\mathbb{C}}^0(F^\bullet(\beta)) \cong \text{Tor}_{\bullet}^R(\mathbb{C}, HHH^0(F^\bullet(\beta)))$. We conjecture that this holds more generally for all links; see (8.7).

As we explained in Remark 1.14, for $k = 2$ and $n = 5$, $\hat{\beta}_{f_{k,n}}$ is the $(2, 3)$ -torus knot, which is isotopic to the trefoil knot: $\hat{\beta}_{f_{k,n}} \cong \mathcal{S}$. It can be alternatively obtained as the closure of the braid $(\sigma_1)^3 \in \mathcal{B}_2$.

Example 3.21 (Trefoil knot). Let $n = 2$, $\beta = (\sigma_1)^3$, and thus $c(\beta) = 1$ and $\chi(\beta) = 1$. We have

$$F^\bullet(\beta) = (B_{s_1}^{\otimes 3} \rightarrow 3B_{s_1}^{\otimes 2} \rightarrow 3B_{s_1} \rightarrow \underline{R}).$$

Here $3B_{s_1}^{\otimes 2}$ denotes the direct sum of three copies of $B_{s_1} \otimes B_{s_1}$, etc. Applying Gaussian elimination as in Example 3.17, we arrive at a simplified complex

$$F^\bullet(\beta) \cong (B_{s_1}\{2\} \rightarrow B_{s_1}\{1\} \rightarrow B_{s_1} \rightarrow \underline{R})$$

with the three maps given by $1 \otimes 1 \mapsto y_1 \otimes 1 + 1 \otimes y_1$, $1 \otimes 1 \mapsto y_1 \otimes 1 - 1 \otimes y_1$, and $1 \otimes 1 \mapsto 1$, respectively. Taking R -invariants, we find

$$HH^0(F^\bullet(\beta)) = \left(R\{3\} \xrightarrow{2y_1} R\{2\} \xrightarrow{0} R\{1\} \xrightarrow{2y_1} \underline{R} \right).$$

We find that the only nonzero terms are

$$H^{0,(0)}(HH^0(F^\bullet(\beta))) \cong H^{-2,(2)}(HH^0(F^\bullet(\beta))) \cong \mathbb{C}.$$

Sending $y_1 \rightarrow 0$, we also compute $H^{k,(p)}(HH^0_C(F^\bullet(\beta)))$, which leads to

$$\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) = q^{\frac{1}{2}}t^{-\frac{1}{2}} + q^{-\frac{1}{2}}t^{\frac{1}{2}}.$$

The corresponding q, t -Catalan number is $C_{2,3}(q, t) = q + t = q^{\frac{1}{2}}t^{\frac{1}{2}} \cdot \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}; q, t)$, in agreement with (4.9).

Remark 3.22. We have $\hat{\beta} \cong \hat{\beta}_{v,w}$ for $v = \text{id}, w = f_{2,5} \in S_5$ and $\chi(\beta) = \chi(\beta_{v,w}) = 1$, thus, $\mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}) = \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_{v,w})$. Similarly to Remark 3.19, we may compute that $\mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta; q, t) = \mathcal{P}_{\text{KR};\mathbb{C}}^{\text{top}}(\beta_{2,5}; q, t)$, and thus $HH^0_C(F^\bullet(\beta_{2,5}))$ is given in Table 3(bottom right).

4. COHOMOLOGY OF POSITROID AND RICHARDSON VARIETIES

We briefly review background on positroid varieties, Richardson varieties and the various versions of cohomology that we will be using.

4.1. Positroid varieties. Recall from Section 1.2 that the Grassmannian $\text{Gr}(k, n)$ is identified with the space of $k \times n$ matrices modulo row operations. Given a $k \times n$ matrix A , we let $\text{RowSpan}(A) \in \text{Gr}(k, n)$ denote its row span and A_1, A_2, \dots, A_n be its columns. We extend this to a sequence $(A_j)_{j \in \mathbb{Z}}$ by requiring

$$A_{j+n} = A_j \quad \text{for all } j \in \mathbb{Z}.$$

Definition 4.1 ([KLS13]). A bijection $f : \mathbb{Z} \rightarrow \mathbb{Z}$ is called a (k, n) -bounded affine permutation if it satisfies

- $f(j+n) = f(j) + n$ for all $j \in \mathbb{Z}$,
- $\sum_{j=1}^n (f(j) - j) = kn$, and
- $j \leq f(j) \leq j+n$ for all $j \in \mathbb{Z}$.

Alternatively, the second condition can be replaced with $k = \#\{j \in [n] \mid f(j) > n\}$.

We let $\mathbf{B}_{k,n}$ denote the (finite) set of (k, n) -bounded affine permutations. For a full rank $k \times n$ matrix A , we let $f_A : \mathbb{Z} \rightarrow \mathbb{Z}$ be given by

$$(4.1) \quad f_A(i) = \min\{j \geq i \mid A_i \in \text{Span}(A_{i+1}, A_{i+2}, \dots, A_j)\} \quad \text{for } i \in \mathbb{Z}.$$

For example, if A_i is a zero column then $f_A(i) = i$, and if A_i is not in the span of other columns then $f_A(i) = i + n$. It is known [KLS13] that f_A is a (k, n) -bounded affine permutation which depends only on the row span of A . The *positroid stratification* of $\text{Gr}(k, n)$ is given by

$$\text{Gr}(k, n) = \bigsqcup_{f \in \mathbf{B}_{k,n}} \Pi_f^\circ, \quad \text{where } \Pi_f^\circ := \{\text{RowSpan}(A) \in \text{Gr}(k, n) \mid f_A = f\}.$$

We extend any permutation $u \in S_n$ to a bijection $\tilde{u} : \mathbb{Z} \rightarrow \mathbb{Z}$ satisfying $\tilde{u}(j+n) = \tilde{u}(j) + n$ for all n . We introduce a (k, n) -bounded affine permutation $\tau_{k,n} : \mathbb{Z} \rightarrow \mathbb{Z}$, determined by

$$\tau_{k,n}(j) = \begin{cases} j+n, & \text{if } 1 \leq j \leq k; \\ j, & \text{if } k+1 \leq j \leq n. \end{cases}$$

Recall that $\mathbf{Q}_{k,n} := \{(v, w) \in S_n \times S_n \mid v \leq w \text{ and } w \text{ is } k\text{-Grassmannian}\}$. The following result explains the bijection $(v, w) \mapsto f_{v,w}$ introduced in Proposition 1.10.

Proposition 4.2 ([KLS13, Proposition 3.15]). *For every $f \in \mathbf{B}_{k,n}$, there exists a unique pair $(v, w) \in \mathbf{Q}_{k,n}$ such that $f = \tilde{w} \circ \tau_{k,n} \circ \tilde{v}^{-1}$.*

Here, \circ denotes the usual composition of bijections $\mathbb{Z} \rightarrow \mathbb{Z}$. We thus define $f_{v,w} := \tilde{w} \circ \tau_{k,n} \circ \tilde{v}^{-1}$. Furthermore, we have the following relationship between positroid and Richardson varieties.

Proposition 4.3 ([KLS13, Theorem 5.9]). *Let $G = \mathrm{PGL}_n(\mathbb{C})$. For each $f = f_{v,w} \in \mathbf{B}_{k,n}$, the natural projection map $\mathrm{Fl}(n) \rightarrow \mathrm{Gr}(k,n)$ restricts to an isomorphism $R_{v,w}^\circ \cong \Pi_f^\circ$. Thus, open positroid varieties are special cases of open Richardson varieties.*

4.2. Torus action and Richardson varieties. The goal of this section is to prove Proposition 1.6. We start by generalizing one direction to Richardson varieties of arbitrary type; the type A specialization is discussed below. Let G be a complex semisimple algebraic group of adjoint type and of rank r , and let \dot{G} denote the simply-connected group of the same Dynkin type. We use the notation $\dot{T}, \dot{B}, \dot{U}$ for the corresponding subgroups of \dot{G} .

Let us give a convenient well-known description (see [BGY06, Theorem 2.3], [Lec16, Lemma 2.2], or [GL19, Lemma 3.1]) of $R_{v,w}^\circ$ as an explicit affine variety. For $v \in W = N_{\dot{G}}(\dot{T})/\dot{T}$, let $\dot{v} \in \dot{G}$ denote an arbitrary fixed representative. For $v \leq w$, denote

$$(4.2) \quad N_{v,w}^\circ := \dot{v}\dot{U}_- \cap \dot{U}_-\dot{v} \cap \dot{B}w\dot{B}.$$

Observe that the set $\dot{B}w\dot{B}$ does not depend on the choice of a representative for w .

Lemma 4.4. *The map $g \mapsto g\dot{B}/\dot{B}$ provides an isomorphism*

$$(4.3) \quad N_{v,w}^\circ \xrightarrow{\sim} R_{v,w}^\circ. \quad \square$$

For $u \in W$, let $\mu(u)$ denote the dimension of the eigenspace with eigenvalue 1 for u acting on \mathfrak{h}^* . We say that $u \in W$ is *elliptic* if $\mu(u) = 0$.

Let P denote the weight lattice of the root system of G (that is, the character lattice of \dot{G}) and $Q \subset P$ denote the root lattice (that is, the character lattice of G). We let $\omega_1, \dots, \omega_r \in P$ denote the fundamental weights. For $u \in W$, we note that $(u - \mathrm{id})P \subseteq Q$.

For $\gamma, \delta \in P$, let $\Delta_{\gamma,\delta}$ denote the corresponding *generalized minor* [BZ97, FZ99] for \dot{G} . The condition that $g \in \dot{B}w\dot{B}$ implies that

$$(4.4) \quad \Delta_{w\omega_i, \omega_i}(g) \neq 0 \quad \text{for } i = 1, 2, \dots, r.$$

Indeed, by [FZ99, Definition 1.4], we have $\Delta_{w\omega_i, \omega_i}(g) = \Delta_{\omega_i, \omega_i}(\overline{w^{-1}g})$ for a certain representative $\overline{w^{-1}g} \in G$ of w^{-1} , so (4.4) follows from [FZ99, Proposition 2.9 and Corollary 2.5].

Proposition 4.5. *Suppose that vw^{-1} is elliptic and $(vw^{-1} - \mathrm{id})P = Q$. Then T acts freely on $R_{v,w}^\circ$ and we have a T -equivariant isomorphism*

$$R_{v,w}^\circ \cong (R_{v,w}^\circ/T) \times T.$$

Proof. Define

$$N_{v,w}^{\circ, \Delta=1} := \{x \in N_{v,w}^\circ \mid \Delta_{\omega_i, w\omega_i}(g) = 1 \quad \text{for } i = 1, 2, \dots, r.\}$$

We will show that $N_{v,w}^{\circ, \Delta=1} \cong R_{v,w}^\circ/T$ and $R_{v,w}^\circ \cong N_{v,w}^{\circ, \Delta=1} \times T$.

The action of \dot{T} on $R_{v,w}^\circ$ corresponds to the action $t \cdot g := tg\dot{v}^{-1}t^{-1}\dot{v}$ for $t \in \dot{T}$ and $g \in N_{v,w}^\circ$. Let $\chi_\omega = \Delta_{\omega, \omega} : \dot{T} \rightarrow \mathbb{C}$ denote the character corresponding to a weight $\omega \in P$. Then

$$\Delta_{w\omega_i, \omega_i}(tg\dot{v}^{-1}t^{-1}\dot{v}) = \chi_{w\omega_i}(t)\chi_{\omega_i}(\dot{v}^{-1}t^{-1}\dot{v})\Delta_{w\omega_i, \omega_i}(g).$$

Since $\chi_{\omega_i}(\dot{v}^{-1}t^{-1}\dot{v}) = \chi_{v\omega_i}(t^{-1})$, the weight of this generalized minor is $(w-v)\omega_i = -v(\mathrm{id} - v^{-1}w)\omega_i$. The condition $\mu(vw^{-1}) = 0$ implies that $\mathrm{id} - v^{-1}w$ is invertible. The condition $(vw^{-1} - \mathrm{id})P = Q$ implies that $(\mathrm{id} - v^{-1}w)\omega_1, (\mathrm{id} - v^{-1}w)\omega_2, \dots, (\mathrm{id} - v^{-1}w)\omega_r$ form a \mathbb{Z} -basis of Q . Since Q is the character lattice of T , it follows that the action of T on $N_{v,w}^\circ$ is free and that the functions $\Delta_{w\omega_i, \omega_i}$, $i = 1, 2, \dots, r$, can be simultaneously set to 1 by a unique element $t \in T$. It follows that $N_{v,w}^{\circ, \Delta=1} \cong R_{v,w}^\circ/T$ and $R_{v,w}^\circ \cong N_{v,w}^{\circ, \Delta=1} \times T$. \square

Remark 4.6. It would be interesting to classify elliptic elements $u \in W$ satisfying the condition $(u - \text{id})P = Q$, which depends only on the conjugacy class of u . It is not satisfied for all elliptic elements. For example, in type D_4 , the longest element w_0 acts by $-\text{id}$ on \mathfrak{h}^* , but we have $2P \subsetneq Q$ since $\frac{\alpha_i}{2} \notin P$ for any $i \in I$.

Proposition 4.7. *Suppose that $c \in W$ is a Coxeter element. Then c is elliptic and satisfies $(c - \text{id})P = Q$.*

Proof. We may assume that c is a standard Coxeter element. That is, $c = s_1 s_2 \cdots s_r$ where s_i are simple generators corresponding to positive simple roots $\alpha_1, \dots, \alpha_r$. Define roots $\beta_1 = \alpha_1$, $\beta_2 = s_1 \alpha_2$, \dots , $\beta_r = s_1 s_2 \cdots s_{r-1} \alpha_r$. By [LP20, Lemma 10.2], c is elliptic. By [LP20, Proposition 10.5] we have $(\text{id} - c)\omega_i = \beta_i \in \alpha_i + \sum_{j < i} \mathbb{Z}\alpha_j$. It follows that $(c - \text{id})P = \bigoplus_{i=1}^r \mathbb{Z}\alpha_i = Q$. \square

Let us now consider the type A case, where $G = \text{PGL}_n(\mathbb{C})$ and $\dot{G} = \text{SL}_n(\mathbb{C})$. For a permutation $u \in S_n$, any of the following conditions are equivalent: (i) u is a single cycle, (ii) u is an elliptic element, and (iii) u is a Coxeter element. Each generalized minor $\Delta_{\omega_i, u\omega_i} : \dot{G} \rightarrow \mathbb{C}$ is the usual matrix minor with row set $\{1, 2, \dots, i\}$ and column set $\{u(1), u(2), \dots, u(i)\}$. The representative \dot{v} in (4.2) may be chosen to be a signed permutation matrix of v . In the next result, $c(\cdot)$ denotes the number of cycles of a permutation; cf. (1.9) (as opposed the Coxeter element c considered above).

Corollary 4.8. *For all $v \leq w \in S_n$ such that $c(wv^{-1}) = 1$, the T -action on $R_{v,w}^\circ$ is free and we have a T -equivariant isomorphism*

$$R_{v,w}^\circ \cong (R_{v,w}^\circ/T) \times T.$$

Remark 4.9. When $R_{v,w}^\circ \cong \Pi_f^\circ$ (see Proposition 4.3), the functions $\Delta_{[1,i], w[1,i]}$ on $N_{v,w}^\circ \cong R_{v,w}^\circ$ coincide with the Plücker coordinates on Π_f° corresponding to the *Grassmann necklace* of f ; see the proof of [GL19, Lemma 4.7]. In particular, for $f = f_{k,n}$, these are the cyclically consecutive maximal minors as in (1.4).

Proof of Proposition 1.6. If $c(\bar{f}) = 1$ then T acts freely on Π_f° by Corollary 4.8. We prove the converse. For $f \in \mathbf{B}_{k,n}$, let us construct a particular representative $X_f^{\min} \in \Pi_f^\circ$. If $f(i) = i$ for some $i \in \mathbb{Z}$ then the corresponding column is zero, so we may assume $f(i) \neq i$ for all i . Let us write \bar{f} in cycle notation:

$$\bar{f} = (a_1 a_2 \cdots a_{m_1})(a_{m_1+1} a_{m_1+2} \cdots a_{m_2}) \cdots (a_{m_r+1} \cdots a_n)$$

so that the minimal index of each cycle comes first. We label these indices left to right: set $\lambda(a_1) := 1$, and for $i = 1, 2, \dots, n-1$, set $\lambda(a_{i+1}) = \lambda(a_i)$ if $a_i < a_{i+1}$ and they belong to the same cycle, and $\lambda(a_{i+1}) = \lambda(a_i) + 1$ otherwise. It is easy to check that $\lambda(a_n) = k$. The element X_f^{\min} is the row span of the $k \times n$ matrix $M = (m_{i,j})$ whose only nonzero entries are $m_{\lambda(a_i), a_i} = 1$. One checks using (4.1) that $X_f^{\min} \in \Pi_f^\circ$. Furthermore, rescaling all columns that belong to a single cycle of \bar{f} by the same value preserves the element X_f^{\min} . Therefore when $c(\bar{f}) > 1$, the T -action on Π_f° is not free. \square

4.3. Mixed Hodge structure. We follow the conventions of [LS16]; see [Del71, PS08] for further background. The results of [LS16] apply to *cluster varieties*. It was shown in [GL19] that open positroid varieties Π_f° , $f \in \mathbf{B}_{k,n}$ are cluster varieties. By [GL19, Lemma 3.6], setting the functions $\Delta_{w\omega_i, \omega_i}$ to 1 as we did in the proof of Proposition 4.5 corresponds to setting the frozen variables to 1 in the cluster structure on Π_f° . Thus, for $f \in \mathbf{B}_{k,n}^{c=1}$, Π_f°/T is a cluster variety with no frozen variables.

Consider a smooth complex algebraic variety Y of dimension d . By [PS08, Lemma-Definition 3.4], the cohomology $H^k(Y, \mathbb{C})$ and the compactly supported cohomology $H_c^k(Y, \mathbb{C})$ are endowed with a

Deligne splitting

$$H^k(Y, \mathbb{C}) = \bigoplus_{p,q \in \mathbb{Z}} H^{k,(p,q)}(Y, \mathbb{C}) \quad \text{and} \quad H_c^k(Y, \mathbb{C}) = \bigoplus_{p,q \in \mathbb{Z}} H_c^{k,(p,q)}(Y, \mathbb{C}).$$

This splitting is functorial and satisfies the Poincaré duality [PS08, Theorem 6.23]:

$$(4.5) \quad H^{k,(p,q)}(Y, \mathbb{C}) \cong H_c^{2d-k,(d-p,d-q)}(Y, \mathbb{C}) \quad \text{for all } k, p, q \in \mathbb{Z}.$$

We say that the cohomology of Y is of *Hodge–Tate type* if $H^{k,(p,q)}(Y, \mathbb{C}) = 0$ whenever $p \neq q$. The notions of mixed Hodge structure and Hodge–Tate type also apply to equivariant cohomology.

Definition 4.10. Let Y be a d -dimensional complex variety whose cohomology is of Hodge–Tate type. Define its *mixed Hodge polynomial* $\mathcal{P}(Y; q, t) \in \mathbb{N}[q^{\frac{1}{2}}, t^{\frac{1}{2}}]$ by

$$(4.6) \quad \mathcal{P}(Y; q, t) := \sum_{k,p \in \mathbb{Z}} q^{p-\frac{k}{2}} t^{\frac{d-k}{2}} \dim H^{k,(p,p)}(Y, \mathbb{C}).$$

We have $H^{k,(p,p)}(Y, \mathbb{C}) = 0$ for $p > k$. By convention, we set $H^{k,(r,r)}(Y, \mathbb{C}) := 0$ for $r \notin \mathbb{Z}$.

It is convenient to record the dimensions of the spaces $H^{k,(p,p)}(Y, \mathbb{C})$ in a *mixed Hodge table*: the columns are labeled by H^0, H^1, \dots, H^d , while the rows are labeled by $k - p = 0, 1, 2, \dots$. Thus, an entry in a column labeled by H^k and in a row labeled by $k - p$ encodes the dimension of $H^{k,(p,p)}(Y, \mathbb{C})$. Examples of mixed Hodge tables are given in Tables 1, 3 and [LS16, Section 6]. For instance, we see from (4.6) that the two rows of Table 1 yield

$$\mathcal{P}(\Pi_{3,8}^\circ/T; q, t) = (q^4 + q^3t + q^2t^2 + qt^3 + t^4) + (q^2t + qt^2),$$

confirming the computation in Example 1.9.

We say that a polynomial $\mathcal{P}(q, t) \in \mathbb{N}[q^{\frac{1}{2}}, t^{\frac{1}{2}}]$ is *q, t -symmetric* if $\mathcal{P}(q, t) = \mathcal{P}(t, q)$. We say that \mathcal{P} is *q, t -unimodal* if for each $a, b \in \frac{1}{2}\mathbb{Z}$, the coefficients $([q^{a-k}t^{b+k}]\mathcal{P})_{k \in \mathbb{Z}}$ form a unimodal sequence.⁶ Recall that Theorem 1.18, proved in Section 8.2, states that the polynomial $\mathcal{P}(R_{v,w}^\circ; q, t)$ is *q, t -symmetric* for all $v \leq w \in W$.

A special case of Theorem 1.18 for positroid varieties follows from the *curious Lefschetz theorem* proved in [LS16, Theorem 8.3]. We say that a $2d$ -dimensional complex algebraic variety Y of Hodge–Tate type satisfies the curious Lefschetz theorem if there is a class $\gamma \in H^{2,(2,2)}(Y, \mathbb{C})$ inducing isomorphisms

$$(4.7) \quad \smile \gamma^{d-p} : H^{p+s,(p,p)}(Y, \mathbb{C}) \cong H^{2d-p+s,(2d-p,2d-p)}(Y, \mathbb{C}).$$

As explained in [LS16], for cluster varieties, one can choose γ to be the Gekhtman–Shapiro–Vainshtein form [GSV10]. In the case that Π_f° is odd-dimensional, the product $\Pi_f^\circ \times \mathbb{C}^*$ will satisfy (4.7), and using the Künneth theorem, unimodality and symmetry for $\mathcal{P}(\Pi_f^\circ; q, t)$ can be deduced.

The curious Lefschetz theorem for positroid varieties relies on the fact that they admit cluster structures [GL19], which is not yet available for Richardson varieties. Another consequence of the curious Lefschetz property is that $\mathcal{P}(\Pi_f^\circ; q, t)$ is *q, t -unimodal*. The question of whether $\mathcal{P}(R_{v,w}^\circ; q, t)$ is *q, t -unimodal* for arbitrary $v \leq w \in W$ remains open.

4.4. Proof of Theorem 1.1 from Theorem 1.17. Let $G = \mathrm{PGL}_n(\mathbb{C})$ and $f = f_{v,w} \in \mathbf{B}_{k,n}^{c=1}$, thus, we have $R_{v,w}^\circ = \Pi_f^\circ$. We set

$$\ell_{v,w} = \ell(w) - \ell(v) = \dim(R_{v,w}^\circ) = \dim(\Pi_f^\circ), \quad d_f := \dim(\mathcal{X}_f^\circ) = \ell_{v,w} - n + 1.$$

⁶This property is sometimes called *parity unimodality* since the terms with integer degrees and with half-integer degrees are required to form separate unimodal sequences.

Since $c(f) = 1$, we have $d_f = 2\chi(\beta_f)$ by (3.12), and the T -action on Π_f° is free by Proposition 1.6. In this case, we have

$$(4.8) \quad H_{T,c}^{k+n-1,(p,p)}(\Pi_f^\circ) \cong H_c^{k,(p,p)}(\mathcal{X}_f^\circ) \cong H^{2d_f-k,(d_f-p,d_f-p)}(\mathcal{X}_f^\circ),$$

where the first isomorphism is Lemma 4.14 below, and the second isomorphism is the Poincaré duality (4.5).

Therefore (1.22) yields

$$(4.9) \quad \mathcal{P}(\mathcal{X}_f^\circ; q, t) = \left(q^{\frac{1}{2}}t^{\frac{1}{2}}\right)^{\chi(\beta_f)} \mathcal{P}_{\text{KR}}^{\text{top}}(\hat{\beta}_f; q, t).$$

This is a special case of (1.24), which is proven in a similar way in Section 8.1. In the case $f = f_{k,n}$ and $\gcd(k, n) = 1$, Remark 1.14 implies that $\hat{\beta}_f$ is a torus knot, for which the right-hand side of (4.9) was shown by Mellit [Mel17] to coincide with $C_{k,n-k}(q, t)$. Thus, Theorem 1.1 follows from Theorem 1.17. \square

4.5. Mixed Hodge structures of open Richardson varieties.

Theorem 4.11. *For any $v \leq w \in W$, the cohomology $H^\bullet(R_{v,w}^\circ, \mathbb{C})$, the compactly supported cohomology $H_c^\bullet(R_{v,w}^\circ, \mathbb{C})$, and the compactly supported T -equivariant cohomology $H_{T,c}^\bullet(R_{v,w}^\circ, \mathbb{C})$ of $R_{v,w}^\circ$ are of Hodge–Tate type.*

We have omitted equivariant cohomology $H_T^\bullet(R_{v,w}^\circ, \mathbb{C})$ from Theorem 4.11 because the statement of Poincaré duality in the equivariant setting is considerably more complicated than for ordinary cohomology.

Lemma 4.12. *Let Y be a complex algebraic variety, $U \subset Y$ an open subvariety, and $Z := Y \setminus U$. Suppose that the compactly supported cohomologies of U and Z are of Hodge–Tate type. Then the same is true for Y .*

The same statement holds for compactly supported T -equivariant cohomology with the assumption that U, Z are T -stable.

Proof. We have a Gysin long exact sequence for the triple (Y, Z, U) (see for example [PS08, (B-15)])

$$(4.10) \quad \cdots \rightarrow H_c^k(U, \mathbb{C}) \rightarrow H_c^k(Y, \mathbb{C}) \rightarrow H_c^k(Z, \mathbb{C}) \rightarrow H_c^{k+1}(U, \mathbb{C}) \rightarrow \cdots$$

the maps of which respect the mixed Hodge structure [dCM04, Theorem 4.1].

Taking the (p, q) piece of the Deligne splitting, we have

$$\cdots \rightarrow H_c^{k,(p,q)}(U, \mathbb{C}) \rightarrow H_c^{k,(p,q)}(Y, \mathbb{C}) \rightarrow H_c^{k,(p,q)}(Z, \mathbb{C}) \rightarrow \cdots$$

By assumption when $p \neq q$, we have $H_c^{k,(p,q)}(U, \mathbb{C}) = 0 = H_c^{k,(p,q)}(Z, \mathbb{C})$. Thus, $H_c^{k,(p,q)}(Y, \mathbb{C}) = 0$. The same proof applies in compactly supported equivariant cohomology. \square

Proof of Theorem 4.11. Since $R_{v,w}^\circ$ is smooth, by (4.5) it suffices to show that the compactly supported cohomology, and the compactly supported equivariant cohomology are of Hodge–Tate type.

We will prove the statement by induction on $\ell_{v,w} = \dim(R_{v,w}^\circ)$. The statement clearly holds if $\ell_{v,w} = 0$, for then $R_{v,w}^\circ$ is a point. We will use a recursion for the varieties $R_{v,w}^\circ$ from [RSW14]. By [RSW14, Lemma 4.3.1 and Proof of Proposition 4.3.6], for any open Richardson $Y = R_{v,w}^\circ$ with $w > v$, we can find a decomposition $Y = U \sqcup Z$, where $U \subset Y$ is open and $Z \subset Y$ is closed, and we have isomorphisms

$$(4.11) \quad Z \cong Y' \times \mathbb{C} \quad U \cong Y'' \times \mathbb{C}^*$$

where Y', Y'' are open Richardson varieties of lower dimension, and Z is possibly empty. Furthermore, U, Z are T -stable and the isomorphisms (4.11) are torus-equivariant, for certain linear actions of T on \mathbb{C}, \mathbb{C}^* . By the Künneth formula, the Hodge–Tate type property is preserved under products. By the inductive hypothesis, the compactly supported (equivariant) cohomology of U

and of Z (when non-empty) are therefore of Hodge–Tate type. It follows that the same statement holds for Y . \square

The following corollary of Theorem 4.11 also follows from combining [LS16, Theorem 8.3] and [GL19].

Corollary 4.13. *For all $f \in \mathbf{B}_{k,n}$, the cohomology of Π_f° is of Hodge–Tate type. If $c(\bar{f}) = 1$ then the cohomology of Π_f°/T is also of Hodge–Tate type.*

Proof. For the second statement, we note that T acts freely on Π_f° by Corollary 4.8. Thus, $H_T^\bullet(\Pi_f^\circ, \mathbb{C}) \cong H^\bullet(\Pi_f^\circ/T, \mathbb{C})$ is of Hodge–Tate type by Theorem 4.11. \square

When computing examples in the next section, we shall repeatedly use the following result.

Lemma 4.14. *Suppose that T acts freely on a complex algebraic variety Y , and let $d := \dim_{\mathbb{C}}(T)$. Then we have an isomorphism of mixed Hodge structures*

$$H_{T,c}^{k+d,(p,q)}(Y, \mathbb{C}) \cong H_c^{k,(p,q)}(Y/T, \mathbb{C}) \quad \text{for all } k, p, q \in \mathbb{Z},$$

and the action of $H_T^\bullet(\text{pt}, \mathbb{C})$ on $H_{T,c}^\bullet(Y, \mathbb{C})$ is trivial (that is, factors through the map $H_T^\bullet(\text{pt}, \mathbb{C}) \rightarrow H_T^0(\text{pt}, \mathbb{C}) \cong \mathbb{C}$).

Proof. Suppose $T = (\mathbb{C}^*)^d$ and let $E_m := (\mathbb{C}^m \setminus \{0\})^d$ be a finite-dimensional approximation to a contractible space ET where T acts freely. Then by definition

$$\begin{aligned} H_{T,c}^k(Y) &= \lim_{m \rightarrow \infty} H_c^k(Y \times_T E_m) \\ &= \lim_{m \rightarrow \infty} H_c^k(Y/T \times E_m) \\ &= \lim_{m \rightarrow \infty} \bigoplus_j H_c^{k-j}(Y/T) \otimes H_c^j(E_m) && \text{by the Künneth formula} \\ &= H_c^{k-d}(Y/T). \end{aligned}$$

We have used the fact that $H_c^\bullet(\mathbb{C}^m \setminus \{0\})$ is \mathbb{C} in dimensions 1 and $2m$ and vanishes in other dimensions. Since $H_c^1(\mathbb{C}^m \setminus \{0\})$ is of type $(0,0)$, the isomorphism is compatible with mixed Hodge structures. \square

4.6. Examples. We compute the compactly supported torus-equivariant cohomology of some open Richardson varieties, corresponding to the examples computed in Section 3.6. As in Section 3.6, we label each example by the link associated to the Richardson variety. Using Table 2, one can compare the below examples with the ones in Section 3.6 and observe that the computations agree with our main results, Theorems 1.16 and 1.17.

Recall that we say that the action of $R = \mathbb{C}[\mathfrak{h}]$ is *trivial* if \mathfrak{h}^* acts by zero. We identify R with $H_T^\bullet(\text{pt}, \mathbb{C})$.

Example 4.15 (Unknot-I). Let $n = 1$ and $v = w = \text{id}$, and thus $\ell_{v,w} = 0$. Then $R_{v,w}^\circ \cong T \cong \text{pt}$, and the T -action is free. The only nonzero terms in the cohomology are

$$(4.12) \quad H^{0,(0,0)}(\text{pt}) \cong H_c^{0,(0,0)}(\text{pt}) \cong H_{T,c}^{0,(0,0)}(\text{pt}) \cong \mathbb{C}.$$

As in Example 3.14, the $R = \mathbb{C}$ -module structure on $H_{T,c}^\bullet(\text{pt})$ is trivial (since T acts freely).

Example 4.16 (Unknot-II). Let $n = 2$, $v = \text{id}$, and $w = s_1$, and thus $\ell_{v,w} = 1$. Then $R_{v,w}^\circ \cong \Pi_{1,2}^\circ \subset \text{Gr}(1,2)$ and we have $R_{v,w}^\circ \cong T \cong \mathbb{C}^*$. The T -action is free and $R_{v,w}^\circ/T \cong \text{pt}$. Using Poincaré duality (4.5) and Lemma 4.14, we find that the only nonzero terms are

$$H^{0,(0,0)}(\mathbb{C}^*) \cong H^{1,(1,1)}(\mathbb{C}^*) \cong \mathbb{C}, \quad H_c^{2,(1,1)}(\mathbb{C}^*) \cong H_c^{1,(0,0)}(\mathbb{C}^*) \cong \mathbb{C}, \quad \text{and}$$

$$(4.13) \quad H_{\mathbb{C}^*, c}^{1, (0,0)}(\mathbb{C}^*) \cong \mathbb{C}.$$

As in Example 3.15, the $R = \mathbb{C}[y_1]$ -module structure on $H_{T,c}^\bullet(R_{v,w}^\circ)$ is trivial (since T acts freely).

Example 4.17 (2-component unlink). Let $n = 2$, $v = s_1$, and $w = s_1$, and thus $\ell_{v,w} = 0$. We have $R_{v,w}^\circ \cong \text{pt}$ and $T \cong \mathbb{C}^*$. The T -action is not free. We have already computed $H^\bullet(\text{pt})$ and $H_{T,c}^\bullet(\text{pt})$ in (4.12). The nonzero terms of $H_{T,c}^\bullet(R_{v,w}^\circ) \cong H_{\mathbb{C}^*, c}^\bullet(\text{pt})$ are given by

$$H_{T,c}^{2p, (p,p)}(\text{pt}) \cong \mathbb{C} \quad \text{for } p = 0, 1, 2, \dots$$

As in Example 3.17, we have $H_{T,c}^\bullet(R_{v,w}^\circ) \cong H_{T,c}^\bullet(\text{pt}) \cong R$ as an R -module.

Example 4.18 (Hopf link). Let $n = 4$, $v = \text{id}$, $w = f_{2,4} = s_2 s_1 s_3 s_2$, and thus $\ell_{v,w} = 4$. We have $Y := R_{v,w}^\circ \cong \Pi_{2,4}^\circ \subset \text{Gr}(2, 4)$, an open positroid variety of dimension 4. It is isomorphic to

$$Y \cong \left\{ \begin{pmatrix} 1 & 0 & a & b \\ 0 & 1 & c & d \end{pmatrix} \middle| (a, b, c, d) \in \mathbb{C}^4 : a \neq 0, d \neq 0, ad - bc \neq 0 \right\}.$$


The action of $T \cong (\mathbb{C}^*)^3$ on Y is not free: T acts by rescaling rows and columns in such a way that the first two columns form the identity matrix. There exists a two-dimensional torus $T' \subset T$ that acts freely on Y : for example, one can always rescale columns 3 and 4 uniquely (since $a, d \neq 0$) to force the minors $\Delta_{2,3}$ and $\Delta_{1,4}$ to be equal to 1:

$$Y/T' \cong \left\{ \begin{pmatrix} 1 & 0 & -1 & -y \\ 0 & 1 & x & 1 \end{pmatrix} \middle| (x, y) \in \mathbb{C}^2 : xy \neq 1 \right\}.$$

The quotient Y/T' can be identified with the two-dimensional A_1 -cluster variety (with one frozen variable; cf. [LS16, Section 6.1]). We denote it by

$$U := Y/T' \cong \{(x, y) \in \mathbb{C}^2 \mid xy \neq 1\} \subset \mathbb{C}^2.$$

The action of T/T' can be identified with the action of \mathbb{C}^* on U with $\lambda \cdot (x, y) = (\lambda x, \lambda^{-1} y)$ for $\lambda \in \mathbb{C}^*$ and $(x, y) \in \mathbb{C}^2$. Forgetting this torus action, it was shown in [LS16, Corollary 7.2] that U is homotopy equivalent⁷ to a pinched torus:



Therefore the Betti numbers of U are $(1, 1, 1)$, and moreover we have [LS16, Section 6.1]

$$H^{0, (0,0)}(U) \cong H^{1, (1,1)}(U) \cong H^{2, (2,2)}(U) \cong \mathbb{C}.$$

We have $Y \cong U \times (\mathbb{C}^*)^2$, which corresponds to multiplying the mixed Hodge polynomial of U by $(q^{\frac{1}{2}} + t^{\frac{1}{2}})^2$. The resulting mixed Hodge table of Y , whose sole row contains the coefficients of the polynomial $(q + q^{\frac{1}{2}} t^{\frac{1}{2}} + t) \cdot (q^{\frac{1}{2}} + t^{\frac{1}{2}})^2$, is given in Table 3(top left).

Let us return to computing the \mathbb{C}^* -equivariant cohomology of U . Denote $W := \mathbb{C}^2$ and let $Z := W \setminus U$ be the hyperbola $\{(x, y) \mid xy = 1\}$. We first compute the compactly supported equivariant cohomologies $H_{\mathbb{C}^*, c}^\bullet(Y)$ and $H_{\mathbb{C}^*, c}^\bullet(Z)$. Set $R := H_{\mathbb{C}^*}^\bullet(\text{pt})$.

⁷We caution the reader that neither the compactly supported cohomology nor the mixed Hodge structure are preserved by homotopy equivalences.

First, suppose that \mathbb{C}^* acts on \mathbb{C}^m linearly in any way. Then it follows directly from the definitions that $H_{\mathbb{C}^*,c}^\bullet(\mathbb{C}^m)$ is a free R -module with generator in degree $2m$. Specifically, all nonzero terms are given by

$$(4.14) \quad H_{\mathbb{C}^*,c}^{2(m+k),(m+k,m+k)}(\mathbb{C}^m) \cong \mathbb{C} \quad \text{for } k = 0, 1, 2, \dots .$$

Next, observe that the variety Z is \mathbb{C}^* -equivariantly isomorphic to the variety \mathbb{C}^* on which \mathbb{C}^* acts freely. By (4.13),

$$(4.15) \quad H_{\mathbb{C}^*,c}^{1,(0,0)}(Z) \cong \mathbb{C}.$$

Now we compute $H_{\mathbb{C}^*,c}^\bullet(U)$. The Gysin sequence (4.10) for the triple (W, Z, U) gives

$$\begin{aligned} 0 \rightarrow H_{\mathbb{C}^*,c}^0(U) \rightarrow H_{\mathbb{C}^*,c}^0(W) \rightarrow H_{\mathbb{C}^*,c}^0(Z) \\ \rightarrow H_{\mathbb{C}^*,c}^1(U) \rightarrow H_{\mathbb{C}^*,c}^1(W) \rightarrow H_{\mathbb{C}^*,c}^1(Z) \rightarrow \dots . \end{aligned}$$

Applying (4.14)–(4.15), we get

$$\begin{aligned} 0 \rightarrow H_{\mathbb{C}^*,c}^0(U) \rightarrow 0 \rightarrow 0 \rightarrow H_{\mathbb{C}^*,c}^1(U) \rightarrow 0 \rightarrow \mathbb{C} \rightarrow H_{\mathbb{C}^*,c}^2(U) \rightarrow 0 \rightarrow 0 \\ \rightarrow H_{\mathbb{C}^*,c}^3(U) \rightarrow 0 \rightarrow 0 \rightarrow H_{\mathbb{C}^*,c}^4(U) \rightarrow \mathbb{C} \rightarrow 0 \rightarrow H_{\mathbb{C}^*,c}^5(U) \rightarrow 0 \rightarrow 0 \rightarrow \dots . \end{aligned}$$

We conclude that the nonzero terms of $H_{\mathbb{C}^*,c}^\bullet(U)$ are given by

$$H_{\mathbb{C}^*,c}^{2,(0,0)}(U) \cong \mathbb{C} \quad \text{and} \quad H_{\mathbb{C}^*,c}^{4+2k,(2+k,2+k)}(U) \cong \mathbb{C} \quad \text{for } k = 0, 1, 2, \dots .$$

By the same computation as in the proof of Lemma 4.14, we have $H_{T,c}^{k+2,(p,q)}(Y) \cong H_{\mathbb{C}^*,c}^{k,(p,q)}(U)$. Thus, the nonzero terms of $H_{T,c}^\bullet(Y)$ are

$$(4.16) \quad H_{T,c}^{4,(0,0)}(Y) \cong \mathbb{C} \quad \text{and} \quad H_{T,c}^{6+2k,(2+k,2+k)}(Y) \cong \mathbb{C} \quad \text{for } k = 0, 1, 2, \dots .$$

Recall that $\ell_{v,w} = 4$. In view of (1.22), the dimensions in (4.16) match perfectly with those computed in (3.18) from the Soergel bimodule perspective.

Remark 4.19. We observe that the $R = \mathbb{C}[y_1]$ -module structure on $H_{T,c}^\bullet(Y)$ also agrees with that computed in (3.18). More generally, for $W = S_n$, Corollary 4.8 can be extended to arbitrary open Richardson varieties $R_{v,w}^\circ$: we have a subtorus $T' \subset T$ of dimension $n - c(\beta)$ acting freely on $R_{v,w}^\circ$, and $H_{T,c}^\bullet(R_{v,w}^\circ) \cong H_{T'/T',c}^\bullet(R_{v,w}^\circ/T')$. Recall from Section 3.4 that we may therefore view both sides of (1.21) as graded modules over a polynomial ring R in $c(\beta) - 1$ variables, and we expect that these R -modules are isomorphic under the grading change (1.22).

It would be interesting to combine Example 4.18 with [LS16] to obtain a description of the equivariant cohomology of more general cluster varieties.

Example 4.20 (Trefoil knot). Let $n = 5$, $v = \text{id}$, $w = f_{2,5} = s_3s_2s_1s_4s_3s_2$, and thus $\ell_{v,w} = 6$. We have $R_{v,w}^\circ \cong \Pi_{2,5}^\circ$, on which the torus $T \cong (\mathbb{C}^*)^5$ acts freely. As explained in Remark 4.9, the quotient is obtained by fixing the cyclically consecutive maximal minors to 1. An explicit parametrization can be chosen as follows:

$$\begin{aligned} \Pi_{2,5}^\circ/T \cong \left\{ \left(\begin{array}{ccccc} 1 & 0 & -1 & -y & \frac{1+y}{1-xy} \\ 0 & 1 & x & xy-1 & 1 \end{array} \right) \middle| (x, y) \in \mathbb{C}^2 : xy \neq 1 \right\} \\ \sqcup \left\{ \left(\begin{array}{ccccc} 1 & 0 & -1 & 1 & z \\ 0 & 1 & -1 & 0 & 1 \end{array} \right) \middle| z \in \mathbb{C} \right\}. \end{aligned}$$

Observe that the point count therefore equals $(q^2 - q + 1) + q = q^2 + 1 = q \cdot C_{2,3}(q, 1/q)$. The variety $\Pi_{2,5}^\circ/T$ is a two-dimensional cluster variety of type A_2 with no frozen variables. Its cohomology was computed in [LS16, Section 6.2]: the nonzero terms are

$$H^{0,(0,0)}(\Pi_{2,5}^\circ/T) \cong H^{2,(2,2)}(\Pi_{2,5}^\circ/T) \cong \mathbb{C}.$$

Multiplying the mixed Hodge polynomial by $\left(q^{\frac{1}{2}} + t^{\frac{1}{2}}\right)^4$, we see that the mixed Hodge table of $\Pi_{2,5}^\circ$ is given in Table 3(top right). By (4.5) and Lemma 4.14, we find

$$H_c^{2,(0,0)}(\Pi_{2,5}^\circ) \cong H_c^{4,(2,2)}(\Pi_{2,5}^\circ) \cong \mathbb{C} \quad \text{and} \quad H_{T,c}^{6,(0,0)}(\Pi_{2,5}^\circ) \cong H_{T,c}^{8,(2,2)}(\Pi_{2,5}^\circ) \cong \mathbb{C}.$$

As in Example 3.21, the R -module structure on $H_{T,c}^\bullet(R_{v,w}^\circ)$ is trivial (since T acts freely).

We now give three examples of Richardson and positroid knots with non-vanishing odd cohomology, as promised in Section 1.12.5. Here by a *positroid knot* we mean a knot of the form $\hat{\beta}_f$ for $f \in \mathbf{B}_{k,n}^{c=1}$.

Example 4.21 (Odd cohomology-I). Let $n = 5$, $v = s_3$, and $w = s_2s_3s_4s_3s_1s_2s_1$, and thus $\ell_{v,w} = 6$ and $c(wv^{-1}) = 1$. The Richardson knot $\beta_{v,w}$ is the *3-twist knot*, listed as 5_2 in Rolfsen's table [Rol90]. By (2.3), the point count is given by

$$\#(R_{v,w}^\circ/T)(\mathbb{F}_q) = q^2 - q + 1.$$

The appearance of $-q$ implies that the cohomology of $R_{v,w}^\circ/T$ *cannot* be concentrated in even degrees.

The following two examples of *positroid* knots were discovered by David Speyer jointly with the second author.

Example 4.22 (Odd cohomology-II). Let $k = 7$, $n = 14$, and let $f = f_{v,w} \in \mathbf{B}_{k,n}^{c=1}$ and $v \leq w \in S_n$ be given by

$$\begin{aligned} f &= [3, 8, 9, 16, 7, 14, 15, 20, 12, 18, 13, 24, 19, 25], \\ v &= (2, 1, 8, 4, 6, 3, 11, 9, 10, 5, 7, 13, 14, 12), \\ w &= (8, 9, 10, 1, 11, 12, 2, 3, 13, 4, 14, 5, 6, 7). \end{aligned}$$

Here v and w are given in one-line notation, and $f = [f(1), f(2), \dots, f(n)]$ is given in *window notation*. The permutation w is 7-Grassmannian, and we have $c(f) = 1$, $\ell(v) = 19$, $\ell(w) = 40$, and $d_f = 8$. The mixed Hodge table of \mathcal{X}_f° is given by

H^k	H^0	H^1	H^2	H^3	H^4	H^5	H^6	H^7	H^8
$k - p = 0$	1	0	1	0	1	0	1	0	1
$k - p = 1$					1	1	1		

In particular, H^5 is non-vanishing.

Example 4.23 (Odd cohomology-III). Let $k = 7$, $n = 14$, and let $f = f_{v,w} \in \mathbf{B}_{k,n}^{c=1}$ and $v \leq w \in S_n$ be given by

$$\begin{aligned} f &= [7, 4, 15, 13, 11, 8, 19, 16, 14, 12, 23, 20, 17, 24], \\ v &= (1, 3, 6, 9, 2, 5, 8, 12, 4, 7, 11, 14, 10, 13), \\ w &= (8, 9, 1, 10, 11, 12, 2, 3, 13, 14, 4, 5, 6, 7). \end{aligned}$$

Similarly, w is 7-Grassmannian, $c(\bar{f}) = 1$, $\ell(v) = 19$, $\ell(w) = 40$, and $d_f = 8$. One can easily compute (e.g. using Theorem 1.13) that the point count is given by

$$\#\mathcal{X}_f^\circ(\mathbb{F}_q) = q^8 + q^6 + 3q^5 - q^4 + 3q^3 + q^2 + 1.$$

Similarly to Example 4.21, this polynomial has a negative term, and therefore the odd cohomology does not vanish.

Remark 4.24. Ivan Cherednik has suggested to us that one might expect odd cohomology vanishing for algebraic knots; see [Pio07] and [Che20, Section 3.4]. The knot 5_2 in Example 4.21 is not algebraic. We thank the anonymous referee for pointing out to us that the positroid knot in Example 4.22 is also not algebraic.

5. EQUIVARIANT DERIVED CATEGORIES OF FLAG VARIETIES

Rather than working with mixed Hodge structures, our proof will be mostly stated in the language of weights [Del80] on étale cohomology.⁸ We assume that the reader is familiar with derived categories of ℓ -adic sheaves in the equivariant setting; see e.g. [BBD82, BL94, LO08] for relevant background.

5.1. Conventions. Fix a prime power q . We shall consider schemes over the finite field \mathbb{F}_q and its algebraic closure $\overline{\mathbb{F}}_q$. For an \mathbb{F}_q -scheme Y , let $Y_{\overline{\mathbb{F}}_q} := Y \times_{\mathrm{Spec}(\mathbb{F}_q)} \mathrm{Spec}(\overline{\mathbb{F}}_q)$ denote the base change to an $\overline{\mathbb{F}}_q$ -scheme. We have the Frobenius automorphism $\mathrm{Fr} : Y_{\overline{\mathbb{F}}_q} \rightarrow Y_{\overline{\mathbb{F}}_q}$ whose fixed points are exactly the points of $Y_{\overline{\mathbb{F}}_q}$ defined over \mathbb{F}_q .

Fix a prime number ℓ different from the characteristic of \mathbb{F}_q and let $\mathbb{k} := \overline{\mathbb{Q}}_\ell$. Let H be an algebraic group acting on an \mathbb{F}_q -variety Y . We consider the bounded derived category $D_{(H)}^b(Y, \mathbb{k})$ of mixed H -constructible (that is, constructible along H -orbits) \mathbb{k} -sheaves on Y , as well as the corresponding category $D_{(H)}^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$ of H -constructible \mathbb{k} -sheaves on $Y_{\overline{\mathbb{F}}_q}$; see [BBD82]. We also let $D_H^b(Y, \mathbb{k})$ and $D_H^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$ denote the corresponding H -equivariant bounded derived categories as in [BL94] (see also [WW] for a discussion of mixed derived categories in the equivariant setting). There are functors $\mathrm{For} : D_H^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k}) \rightarrow D_{(H)}^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$ and $\mathrm{For} : D_H^b(Y, \mathbb{k}) \rightarrow D_{(H)}^b(Y, \mathbb{k})$ forgetting the equivariant structure. There are also functors $\omega : D_{(H)}^b(Y, \mathbb{k}) \rightarrow D_{(H)}^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$ and $\omega : D_H^b(Y, \mathbb{k}) \rightarrow D_H^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$ obtained by extension of scalars from \mathbb{F}_q to $\overline{\mathbb{F}}_q$.

The language of algebraic stacks [LO08] allows us to switch between ordinary and equivariant derived categories at our convenience. For example, $D^b([Y/H], \mathbb{k}) \cong D_H^b(Y, \mathbb{k})$.

We denote by $[m]$ the cohomological shift m steps to the left in a derived category as in Section 3.2. For $\mathcal{F}, \mathcal{G} \in D_H^b(Y, \mathbb{k})$ and $k \in \mathbb{Z}$, let

$$(5.1) \quad \mathrm{Ext}^k(\mathcal{F}, \mathcal{G}) = \mathrm{Ext}_Y^k(\mathcal{F}, \mathcal{G}) := \mathrm{Hom}_{D_H^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})}(\omega\mathcal{F}, \omega\mathcal{G}[k]) \quad \text{for } k \in \mathbb{Z}.$$

The space $\mathrm{Hom}_{D_H^b(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})}(\omega\mathcal{F}, \omega\mathcal{G}[k])$ has a natural action of the Frobenius Fr . Therefore $\mathrm{Ext}^\bullet(\mathcal{F}, \mathcal{G})$ is a graded $H_H^\bullet(\mathrm{pt}, \mathbb{k})$ -module equipped with an action of Fr , or in other words, an $(H_H^\bullet(\mathrm{pt}, \mathbb{k}), \mathrm{Fr})$ -module. The actual extension groups in $D_H^b(Y, \mathbb{k})$ are denoted by $\mathrm{ext}^k(\mathcal{F}, \mathcal{G})$, and are related to $\mathrm{Ext}^k(\mathcal{F}, \mathcal{G})$ by the exact sequence [BBD82, (5.1.2.5)]

$$(5.2) \quad 0 \rightarrow \mathrm{Ext}^{i-1}(\mathcal{F}, \mathcal{G})_{\mathrm{Fr}} \rightarrow \mathrm{ext}^i(\mathcal{F}, \mathcal{G}) \rightarrow \mathrm{Ext}^i(\mathcal{F}, \mathcal{G})^{\mathrm{Fr}} \rightarrow 0,$$

where $\mathrm{Ext}^i(\mathcal{F}, \mathcal{G})^{\mathrm{Fr}}$ and $\mathrm{Ext}^{i-1}(\mathcal{F}, \mathcal{G})_{\mathrm{Fr}}$ denote Frobenius invariants and coinvariants, respectively. We denote by $\mathrm{hom}(\mathcal{F}, \mathcal{G}) = \mathrm{ext}^0(\mathcal{F}, \mathcal{G}) = \mathrm{Hom}_{D_H^b(Y, \mathbb{k})}(\mathcal{F}, \mathcal{G})$ the Hom groups in $D_H^b(Y, \mathbb{k})$.

We fix an isomorphism $\mathbb{k} \cong \mathbb{C}$ and denote by $|\lambda|$ the norm of $\lambda \in \mathbb{k}$ considered as an element of \mathbb{C} . If M is an Fr -module, then the *weights* of Fr on M are the real numbers $2 \log(\lambda) / \log(q)$ for λ an eigenvalue of the action of Fr . All weights we consider will be integers: the cohomology sheaves of an object $\mathcal{F} \in D_{(H)}^b(Y, \mathbb{k})$ are required to have punctual integer weights; see [BBD82, 5.1.5]. We fix a square root $(1/2)$ of the Tate twist, and for $\mathcal{F} \in D_H^b(Y, \mathbb{k})$ and $r \in \mathbb{Z}$, we denote by $\mathcal{F}(r/2)$ the corresponding Tate twist of \mathcal{F} .

Recall [BBD82, 5.1] that $\mathcal{F} \in D_H^b(Y, \mathbb{k})$ has weights $\leq r$ if for each i the sheaf $H^i(\mathcal{F})$ has mixed punctual weights $\leq r + i$. We say that $\mathcal{F} \in D_H^b(Y, \mathbb{k})$ has weights $\geq r$ if the Verdier dual $\mathbb{D}\mathcal{F}$ has mixed punctual weights $\leq -r$. Finally, we say that $\mathcal{F} \in D_H^b(Y, \mathbb{k})$ is *pure of weight* r if it has weights $\leq r$ and weights $\geq r$. If \mathcal{F} is pure of weight r then $\mathcal{F}[1]$ is pure of weight $r + 1$, while $\mathcal{F}(1/2)$ is pure of weight $r - 1$.

⁸As pointed out to us by Wolfgang Soergel, our results could also be formulated using the language of equivariant mixed Tate motives [SVW18].

For an integer $r \in \mathbb{Z}$, we denote by $\text{Ext}^{k, (r/2)}(\mathcal{F}, \mathcal{G}) \subset \text{Ext}^k(\mathcal{F}, \mathcal{G})$ the generalized eigenspace for Fr of weight r . Thus,

$$(5.3) \quad \text{Ext}(\mathcal{F}, \mathcal{G}) = \bigoplus_{k, r \in \mathbb{Z}} \text{Ext}^{k, (r/2)}(\mathcal{F}, \mathcal{G}).$$

For all $\mathcal{F}, \mathcal{G} \in \text{D}_H^b(Y, \mathbb{k})$ and $k, k', r, r' \in \mathbb{Z}$, we have

$$(5.4) \quad \begin{aligned} \text{Ext}^{k, (r/2)}(\mathcal{F}[-k'](-r'/2), \mathcal{G}) &\cong \text{Ext}^{k, (r/2)}(\mathcal{F}, \mathcal{G}[k'](r'/2)) \\ &\cong \text{Ext}^{k+k', ((r+r')/2)}(\mathcal{F}, \mathcal{G}). \end{aligned}$$

5.2. Equivariant cohomology. For $\mathcal{F} \in \text{D}_H^b(Y, \mathbb{k})$, the equivariant hypercohomology $\mathbb{H}_H^\bullet(\mathcal{F}) = \mathbb{H}_H^\bullet(Y, \mathcal{F})$ is defined by

$$(5.5) \quad \mathbb{H}_H^\bullet(\mathcal{F}) := \text{Ext}^\bullet(\mathbb{k}_Y, \mathcal{F}).$$

In particular, we have $\mathbb{H}_H^{k, (r/2)}(\mathcal{F}[k'](r'/2)) = \mathbb{H}_H^{k+k', ((r+r')/2)}(\mathcal{F})$.

Let $\pi : Y \rightarrow \text{pt} := \text{Spec}(\mathbb{F}_q)$ be the projection to a point. By definition, the H -equivariant cohomology $H_H^\bullet(Y, \mathbb{k})$ and the compactly supported H -equivariant cohomology $H_{H,c}^\bullet(Y, \mathbb{k})$ of Y is given by

$$(5.6) \quad H_H^\bullet(Y, \mathbb{k}) := \mathbb{H}_H^\bullet(Y, \mathbb{k}_Y) = \mathbb{H}_H^\bullet(\text{pt}, \pi_* \mathbb{k}_Y) \quad \text{and} \quad H_{H,c}^\bullet(Y, \mathbb{k}) := \mathbb{H}_H^\bullet(\text{pt}, \pi! \mathbb{k}_Y).$$

Both $H_H^\bullet(Y, \mathbb{k})$ and $H_{H,c}^\bullet(Y, \mathbb{k})$ are graded $(H_H^\bullet(\text{pt}, \mathbb{k}), \text{Fr})$ -modules.

5.3. Flag varieties. We fix a semisimple algebraic group G , split over \mathbb{F}_q , and a maximal torus and a Borel subgroup $T \subset B \subset G$. Let $X = G/B$ be the flag variety over \mathbb{F}_q and let $X_{\overline{\mathbb{F}}_q} := X \times_{\text{Spec}(\mathbb{F}_q)} \text{Spec}(\overline{\mathbb{F}}_q)$ be obtained by extending scalars. The variety X is stratified by B -orbits $\mathring{X}_w := BwB/B$ (known as *Schubert cells*):

$$X = \bigsqcup_{w \in W} \mathring{X}_w.$$

Let $R = H_B^\bullet(\text{pt}, \mathbb{k}) \cong \mathbb{k}[\mathfrak{h}]$.

Remark 5.1. We switch from working over $R = \mathbb{C}[\mathfrak{h}]$ to $R = \mathbb{k}[\mathfrak{h}]$. The results in Section 3 do not depend on the field as long as it is of characteristic zero. Therefore on the Soergel bimodule side, one can freely switch between working over \mathbb{C} and over \mathbb{k} .

For $w \in W$, we let $i_w : \mathring{X}_w \rightarrow X$ be the inclusion map. Introduce the *standard* and *costandard sheaves* $\Delta_w, \nabla_w \in \text{D}_B^b(X, \mathbb{k})$ defined by

$$\Delta_w := i_{w,*} \mathbb{k}_{\mathring{X}_w}^\circ[\ell(w)](\ell(w)/2) \quad \text{and} \quad \nabla_w := i_{w,*} \mathbb{k}_{\mathring{X}_w}^\circ[\ell(w)](\ell(w)/2).$$

Here, $\mathbb{k}_{\mathring{X}_w}^\circ$ denotes the constant sheaf on \mathring{X}_w . The *intersection cohomology sheaves* $\mathcal{IC}_w \in \text{D}_B^b(X, \mathbb{k})$ are defined using the intermediate extension functor $i_{w,!*}$:

$$\mathcal{IC}_w := i_{w,!*} \mathbb{k}_{\mathring{X}_w}^\circ[\ell(w)](\ell(w)/2).$$

Since we have $\text{D}_B^b(X, \mathbb{k}) \cong \text{D}_{B \times B}^b(G, \mathbb{k})$, the equivariant cohomology $\mathbb{H}_B^\bullet(\mathcal{F})$ is a graded $(R \otimes R, \text{Fr})$ -module. Furthermore, there is a restriction functor $\text{Res}_{T,B} : \text{D}_B^b(X, \mathbb{k}) \rightarrow \text{D}_T^b(X, \mathbb{k})$ (see [BL94]), and we also have the hypercohomology functor $\mathbb{H}_T^\bullet : \text{D}_T^b(X, \mathbb{k}) \rightarrow H_T^\bullet(\text{pt}) - \text{mod}$. It is well known that $H_B^\bullet(\text{pt}, \mathbb{k}) \cong R \cong H_T^\bullet(\text{pt}, \mathbb{k})$, and furthermore we have

$$(5.7) \quad \text{Res}_{T,B} : \text{D}_B^b(\text{pt}, \mathbb{k}) \cong \text{D}_T^b(\text{pt}, \mathbb{k}) \quad \text{and} \quad \mathbb{H}_B^\bullet(X, \mathcal{F}) \cong \mathbb{H}_T^\bullet(X, \text{Res}_{T,B} \mathcal{F})$$

for $\mathcal{F} \in \text{D}_B^b(X, \mathbb{k})$.

5.4. Equivariant cohomology of open Richardson varieties. We split the proof of our main result into two parts. We will focus on the equivariant case (Theorem 1.17). The proof of its non-equivariant version (Theorem 1.16) will follow as a byproduct, and will be discussed in Section 8.1.

Recall from Section 1.9 that our goal is to show the following result.

Theorem 5.2. *For all $v \leq w \in W$, we have an isomorphism of bigraded R -modules*

$$HHH^0(F_{v,w}^\bullet) \cong H_{T,c}^\bullet(R_{v,w}^\circ).$$

For all $k, p \in \mathbb{Z}$, it restricts to an isomorphism

$$H^{k,(p)}(HH^0(F_{v,w}^\bullet)) \cong H_{T,c}^{\ell_{v,w}+2p+k,(p,p)}(R_{v,w}^\circ)$$

of vector spaces.

We will accomplish this in two steps. The first one is an equivariant version of [RSW14, Proposition 4.2.1].

Proposition 5.3. *For all $v \leq w \in W$, we have an isomorphism of bigraded R -modules*

$$H_{T,c}^\bullet(R_{v,w}^\circ) \cong \text{Ext}^\bullet(\Delta_v, \Delta_w).$$

For all $m, r \in \mathbb{Z}$, it restricts to an isomorphism

$$H_{T,c}^{\ell_{v,w}+m,(r/2,r/2)}(R_{v,w}^\circ) \cong \text{Ext}^{m,((r-\ell_{v,w})/2)}(\Delta_v, \Delta_w)$$

of vector spaces. (In particular, both sides are zero for odd r .)

The second step passes through the *mixed equivariant derived category* of [AR16a, AR16b] and involves the *degrading functor* of [Bei87] (see also [Rid13]).

Proposition 5.4. *For all $v \leq w \in W$, we have an isomorphism of bigraded R -modules*

$$HHH^0(F_{v,w}^\bullet) \cong \text{Ext}^\bullet(\Delta_v, \Delta_w).$$

For all $k, r \in \mathbb{Z}$, it restricts to an isomorphism

$$(5.8) \quad H^{k,(r/2)}(HH^0(F_{v,w}^\bullet)) \cong \text{Ext}^{r+k,((r-\ell_{v,w})/2)}(\Delta_v, \Delta_w)$$

of vector spaces. (In particular, both sides are zero for odd r .)

5.5. From weights to the mixed Hodge numbers. We briefly explain the standard relation between the mixed Hodge structure of the complex variety $Y_{\mathbb{C}} = (R_{v,w}^\circ)_{\mathbb{C}}$ and the étale cohomology of the variety $Y_{\overline{\mathbb{F}}_q} = (R_{v,w}^\circ)_{\overline{\mathbb{F}}_q}$. First, by the comparison theorem ([Mil13, Theorem 21.1]) between Betti cohomology and étale cohomology of $Y_{\mathbb{C}}$, we have an isomorphism $H_{Betti}^k(Y_{\mathbb{C}}, \mathbb{k}) \cong H_{et}^k(Y_{\mathbb{C}}, \mathbb{k})$ which preserves the weight filtration of both sides. Next, we find a discrete valuation ring $S \subset \mathbb{C}$ with residue field $\overline{\mathbb{F}}_q$ and construct the Richardson variety Y_S over S . Then $Y_{\mathbb{C}}$ and $Y_{\overline{\mathbb{F}}_q}$ are obtained from Y_S via base change, and we obtain isomorphisms between the étale cohomologies $H^\bullet(Y_{\mathbb{C}}, \mathbb{k}) \cong H^\bullet(Y_S, \mathbb{k}) \cong H^\bullet(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$, compatibly with the weight filtrations; see [Mil13, Section 20]. For $H^\bullet(Y_{\overline{\mathbb{F}}_q}, \mathbb{k})$, the weight filtration is obtained by taking sums of generalized eigenspaces of the Frobenius Fr . Finally, the cohomology of $Y_{\mathbb{C}}$ is of Hodge–Tate type, so the weight filtration is simply given by $W^{2r}(H^k(Y_{\mathbb{C}}, \mathbb{C})) = \bigoplus_{p \leq r} H^{k,(p,p)}(Y_{\mathbb{C}}, \mathbb{C})$. (All these statements hold also equivariantly.) To sum up:

Proposition 5.5. *For all $v \leq w \in W$ and $k, p \in \mathbb{Z}$, we have the equalities*

$$\begin{aligned} \dim_{\mathbb{C}} H^{k,(p,p)}((R_{v,w}^\circ)_{\mathbb{C}}, \mathbb{C}) &= \dim_{\mathbb{k}} H^{k,(p)}((R_{v,w}^\circ)_{\overline{\mathbb{F}}_q}, \mathbb{k}), \\ \dim_{\mathbb{C}} H_{T,c}^{k,(p,p)}((R_{v,w}^\circ)_{\mathbb{C}}, \mathbb{C}) &= \dim_{\mathbb{k}} H_{T,c}^{k,(p)}((R_{v,w}^\circ)_{\overline{\mathbb{F}}_q}, \mathbb{k}) \end{aligned}$$

where $H_{T,c}^{k,(p)}((R_{v,w}^\circ)_{\overline{\mathbb{F}}_q}, \mathbb{k}) = \mathbb{H}_T^{k,(p)}(\text{pt}, \pi_1 \underline{\mathbb{k}}_{R_{v,w}^\circ}) = \text{Ext}^{k,(p)}(\underline{\mathbb{k}}_{\text{pt}}, \pi_1 \underline{\mathbb{k}}_{R_{v,w}^\circ})$ as in (5.6).

See also [RSW14, Remark 7.1.4] where a comparison between derived categories of flag varieties is given.

6. PROOF OF PROPOSITION 5.3

We follow the steps in the proof of [RSW14, Proposition 4.2.1]. Using (5.4) and the adjunction $(i_{v,!}, i_v^!)$, we find

$$\begin{aligned} \mathrm{Ext}^{m, ((r-\ell_{v,w})/2)}(\Delta_v, \Delta_w) &\cong \mathrm{Ext}^{m, ((r-\ell_{v,w})/2)}(i_{v,!} \underline{\mathbb{k}}_{\mathring{X}_v}^\circ, i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ [\ell_{v,w}] (\ell_{v,w}/2)) \\ &\cong \mathrm{Ext}^{m+\ell_{v,w}, (r/2)}(i_{v,!} \underline{\mathbb{k}}_{\mathring{X}_v}^\circ, i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ) \\ &\cong \mathrm{Ext}^{m+\ell_{v,w}, (r/2)}(\underline{\mathbb{k}}_{\mathring{X}_v}^\circ, i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ). \end{aligned}$$

Note that $i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ \in D_B^b(\mathring{X}_v, \mathbb{k})$. By (5.5) and (5.7), we have

$$\mathrm{Ext}^\bullet(\underline{\mathbb{k}}_{\mathring{X}_v}^\circ, i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ) \cong \mathbb{H}_B^\bullet(\mathring{X}_v, i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ) \cong \mathbb{H}_T^\bullet(\mathring{X}_v, i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ).$$

We now switch to working with T -equivariant derived categories. First, we state a T -equivariant version of [Soe89, Proposition 1]. Let Z be a T -variety and let $q : X \hookrightarrow Z$ be an inclusion of a closed T -subvariety. An action $\mathbb{G}_m \times Z \rightarrow Z$ *contracts* Z to X if there is a commutative diagram

$$\begin{array}{ccc} \mathbb{G}_m \times Z & \xrightarrow{\mathrm{act}} & Z \\ \downarrow j & & \parallel \\ \mathbb{A}^1 \times Z & \xrightarrow{\mathrm{act}_0} & Z \\ \uparrow i & & \uparrow q \\ Z & \xrightarrow{p} & X \end{array}$$

where i, j are the obvious maps and $p \circ q = \mathrm{id}$, and all arrows are T -equivariant, with T acting trivially on \mathbb{G}_m and \mathbb{A}^1 . Let $\pi : \mathbb{G}_m \times Z \rightarrow Z$ be the projection and suppose that $\mathcal{F} \in D_T^b(Z)$ is \mathbb{G}_m -equivariant, i.e. satisfies $\mathrm{act}^*(\mathcal{F}) \cong \pi^*(\mathcal{F}) \in D_T^b(\mathbb{G}_m \times Z)$. There is a natural morphism $q^! \rightarrow p_!$ of functors (obtained by composing the adjunction morphism $q_! q^! \rightarrow \mathrm{id}$ with $p_!$) and we have the following.

Proposition 6.1 (cf. [Soe89, Proposition 1]). *The map $q^! \mathcal{F} \rightarrow p_! \mathcal{F}$ is an isomorphism.*

Let $a : \mathring{X}_v \rightarrow \mathrm{pt}$ and $b : R_{v,w}^\circ \rightarrow \mathrm{pt}$ be the projections (cf. Figure 3(right)). Our next goal is to manipulate the object $i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ$, in order to establish the following result.

Lemma 6.2. *We have $i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ \cong a^* b_! \underline{\mathbb{k}}_{R_{v,w}^\circ}$ in $D_T^b(\mathring{X}_v)$.*

Proof. Recall that $\mathring{X}_w := (BwB)/B$. Denote $\mathring{X}_v^- := (B_-vB)/B$, and thus $R_{v,w}^\circ = \mathring{X}_w \cap \mathring{X}_v^-$. Let $X_w := \bigsqcup_{u \leq w} \mathring{X}_u$ be the closure of \mathring{X}_w , and denote $R_{v,w}^\circ := X_w \cap \mathring{X}_v^-$. A diagram of the various inclusions between the spaces $\mathring{X}_w, \mathring{X}_v, X_w$, and $X = G/B$ is given in Figure 3(left). With this notation, we have isomorphisms

$$i_v^! i_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ \cong i_{v,w}^! \bar{i}_w^! \bar{j}_w^! \underline{\mathbb{k}}_{\mathring{X}_w}^\circ \cong i_{v,w}^! j_{w,!} \underline{\mathbb{k}}_{\mathring{X}_w}^\circ.$$

The first isomorphism follows from the usual composition rules for sheaf operations; see e.g. [BL94, Section 1.4.2]. The second isomorphism follows from $\bar{i}_w^! \bar{j}_w^! \cong \mathrm{id}$.

Consider the commutative diagram in Figure 3(middle). The map k is given by $k(zB/B) = (vB/B, zB/B)$ and π is the obvious projection map. The map j'_w has two components: the first




FIGURE 3. Three commutative diagrams from [RSW14].

one is the inclusion $R_{v,w}^\circ \hookrightarrow R_{v,\bar{w}}^\circ$, and the second one is the identity map $\mathring{X}_v \xrightarrow{\sim} \mathring{X}_v$. The map $j_w : \mathring{X}_w \hookrightarrow X_w$ is the inclusion map as above. It remains to define the maps j and j' . They have been considered in [KL80, Section 1.4]; see for example [FS00, KQY13] for further details. Observe that $R_{v,w}^\circ \subset R_{v,\bar{w}}^\circ \subset \mathring{X}_v^-$. The maps j, j' are the restrictions of a map $\phi_v : \mathring{X}_v^- \times \mathring{X}_v \hookrightarrow X$ defined as follows. Recall that $U \subset B$ and $U_- \subset B_-$ are the unipotent radicals of B and B_- . Any element of \mathring{X}_v^- can be written uniquely as xvB/B for an element $x \in U_- \cap vU_-v^{-1}$. (These objects do not depend on the choice of representative \dot{v} of v .) Similarly, any element of \mathring{X}_v can be written uniquely as yvB/B for an element $y \in U \cap vU_-v^{-1}$. We then define

$$\phi_v(xvB/B, yvB/B) := yxvB/B.$$

It is not hard to see (using *Gaussian decomposition*) that the map ϕ_v is injective and yields an isomorphism $\phi_v : \mathring{X}_v^- \times \mathring{X}_v \xrightarrow{\sim} vB_-B/B$. Moreover, if $xvB/B \in \mathring{X}_u$ for some $u \in W$, then we have $yxvB/B \in \mathring{X}_u$ since $y \in U \subset B$. Thus, ϕ_v restricts to an inclusion $R_{v,u}^\circ \times \mathring{X}_v \hookrightarrow \mathring{X}_u$ for each $u \in W$. The map j' is this inclusion for the special case $u := w$. The map j is the restriction of ϕ_v to the union of $R_{v,u}^\circ \times \mathring{X}_v$ over all $u \in W$ satisfying $v \leq u \leq w$.

The torus T acts on each space in each commutative diagram in Figure 3. The action on the direct products $R_{v,w}^\circ \times \mathring{X}_v$ and $R_{v,\bar{w}}^\circ \times \mathring{X}_v$ is given by $t \cdot (aB/B, bB/B) = (taB/B, tbB/B)$ for $t \in T$. Notice that conjugation by t preserves each of the subgroups U , U_- , and vU_-v^{-1} . Therefore for $x \in U_- \cap vU_-v^{-1}$, we have $txvB/B = txt^{-1}vB/B$, where $txt^{-1} \in U_- \cap vU_-v^{-1}$, and similarly for $y \in U \cap vU_-v^{-1}$. Thus, the map ϕ_v is T -equivariant:

$$\phi_v(txvB/B, tyvB/B) = tyxvB/B.$$

We conclude that all maps in Figure 3 are T -equivariant.

The maps j, j', j_w, j'_w in Figure 3(middle) form a Cartesian square. We get the following isomorphisms:

$$i_{v,w}^! j_w^! \mathbb{k}_{\mathring{X}_w} \cong k^! j'^! j_w^! \mathbb{k}_{\mathring{X}_w} \cong k^! j^* j_w^! \mathbb{k}_{\mathring{X}_w} \cong k^! j_w^! j'^* \mathbb{k}_{\mathring{X}_w} \cong k^! j_w^! \mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v}.$$

The first and the last isomorphisms are trivial. The second isomorphism follows from the fact that j is an open embedding [BL94, Section 1.4.5]. The third isomorphism is the base change theorem [BL94, Section 1.4.6].

We now apply Proposition 6.1 with $q = k$, and $p = \pi$, and $\mathcal{F} = j_w^! \mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v}$. The \mathbb{G}_m -action is the composition of the T -action with the cocharacter $\chi_\rho^\vee : \mathbb{G}_m \rightarrow T$, where ρ is a strictly dominant coweight satisfying $\langle \rho, \alpha \rangle > 0$ for any positive root α . Since ρ is strictly dominant, this \mathbb{G}_m -action extends to an \mathbb{A}^1 -action by the same argument as in [GKL19, Section 8.2]. The \mathbb{A}^1 -action is obviously compatible with the T -action, and thus Proposition 6.1 applies.

We obtain

$$k^! j_w^! \mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v} \cong \pi^! j_w^! \mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v}.$$

Applying base change to the Cartesian diagram in Figure 3(right), we get

$$\pi!j_w^!\mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v} \cong (\pi \circ j_w^!)!\mathbb{k}_{R_{v,w}^\circ \times \mathring{X}_v} \cong (\pi \circ j_w^!)!r^*\mathbb{k}_{R_{v,w}^\circ} \cong a^*b!\mathbb{k}_{R_{v,w}^\circ}. \quad \square$$

So far we have constructed an isomorphism

$$(6.1) \quad \begin{aligned} \mathrm{Ext}^{m,((r-\ell_{v,w})/2)}(\Delta_v, \Delta_w) &\cong \mathbb{H}_T^{m+\ell_{v,w},(r/2)}(\mathring{X}_v, a^*b!\mathbb{k}_{R_{v,w}^\circ}) \\ &\cong \mathbb{H}_T^{m+\ell_{v,w},(r/2)}(\mathrm{pt}, a_*a^*b!\mathbb{k}_{R_{v,w}^\circ}). \end{aligned}$$

Since \mathring{X}_v is an affine space that is T -equivariantly homotopy-equivalent to a point (cf. [Soe89, Lemma 1]), the adjunction (a^*, a_*) induces an isomorphism of functors $a_*a^* \rightarrow \mathrm{id}$, and thus

$$(6.2) \quad \mathbb{H}_T^{m+\ell_{v,w},(r/2)}(\mathrm{pt}, a_*a^*b!\mathbb{k}_{R_{v,w}^\circ}) \cong \mathbb{H}_T^{m+\ell_{v,w},(r/2)}(\mathrm{pt}, b!\mathbb{k}_{R_{v,w}^\circ}),$$

which equals to $H_{T,c}^{m+\ell_{v,w},(r/2,r/2)}(R_{v,w}^\circ)$ by (5.6). All the isomorphisms are natural (coming from sheaf operations) and thus (6.2) is compatible with the action of R on both sides. This completes the proof of Proposition 5.3. \square

7. PROOF OF PROPOSITION 5.4

Recall from Remark 5.1 that we switch to working with Soergel bimodules over \mathbb{k} , so that e.g. $R = \mathbb{k}[\mathfrak{h}]$. By definition, given a (graded) Soergel bimodule $B = \bigoplus_r B^r$, Fr acts diagonally on each B^r by multiplication by q^r .

Equivariant derived categories are identified with categories of dg-modules in the work of Bernstein-Lunts [BL94]. By using the formalism of Yun [BY13, Appendix B], we avoid explicit mention of dg-modules in the situation of interest to us.

An (R, Fr) -module is an R -module M equipped with an action of $\mathbb{Z}\mathrm{Fr}$, such that $\mathrm{Fr}(r \cdot x) = \mathrm{Fr}(r) \cdot \mathrm{Fr}(x)$ for $x \in M$ and $r \in R$. The twist functor $\{m/2\}$ sends a module M to the module $M\{m/2\}$ where the action of Fr has been multiplied by $q^{m/2}$. Let $\mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr})$ denote the full triangulated subcategory of the derived category of (R, Fr) -modules generated by half-integer twists of R . According to [BY13, Corollary B.4.1], we have an equivalence of triangulated categories $\mathrm{D}_B^b(\mathrm{pt}, \mathbb{k}) \cong \mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr})$. Similarly, we define $\mathrm{D}_{\mathrm{perf}}(R \otimes R, \mathrm{Fr})$ and have an equivalence $\mathrm{D}_{B \times B}^b(\mathrm{pt}, \mathbb{k}) \cong \mathrm{D}_{\mathrm{perf}}(R \otimes R, \mathrm{Fr})$.

Recall that $\mathrm{Hom}_{\mathbb{S}\mathrm{Bim}}$ includes only bimodule morphisms of degree zero. We let

$$\mathrm{Hom}_{R \otimes R}(B, B') := \bigoplus_{r \in \mathbb{Z}} \mathrm{Hom}_{\mathbb{S}\mathrm{Bim}}(B, B'\{-r/2\})$$

denote the space of morphisms of arbitrary degree. Thus, $\mathrm{Hom}_{R \otimes R}(B, B')$ is an (R, Fr) -module. Given a complex $C^\bullet \in \mathrm{K}^b\mathbb{S}\mathrm{Bim}$, we regard $\mathrm{Hom}_{R \otimes R}(R, C^\bullet)$ (obtained by applying $\mathrm{Hom}_{R \otimes R}(R, -)$ term-wise) as a complex of (R, Fr) -modules, treated as an element of $\mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr})$.

Now, for $\mathcal{F}, \mathcal{G} \in \mathrm{D}_B^b(X, \mathbb{k})$, let $R\mathcal{H}om(\mathcal{F}, \mathcal{G}) \in \mathrm{D}_B^b(X, \mathbb{k})$ denote the internal derived hom. With $\pi : G \rightarrow \mathrm{pt}$, we define

$$\mathrm{RHom}(\mathcal{F}, \mathcal{G}) = \mathrm{RHom}_X(\mathcal{F}, \mathcal{G}) := \pi_* R\mathcal{H}om(\mathcal{F}, \mathcal{G}) \in \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k}).$$

Thus, $\mathbb{H}_B^\bullet(\mathrm{RHom}(\mathcal{F}, \mathcal{G})) = \mathrm{Ext}^\bullet(\mathcal{F}, \mathcal{G})$. We shall establish the following strengthening of Proposition 5.4.

Proposition 7.1. *We have*

$$\mathrm{RHom}_X(\Delta_v(-\ell(v)/2), \Delta_w(-\ell(w)/2)) \cong \mathrm{Hom}_{R \otimes R}(R, F_{v,w}^\bullet)$$

inside $\mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr})$.

Proposition 5.4 follows from Proposition 7.1 by taking cohomology $\mathbb{H}_B : D_B^b(\text{pt}, \mathbb{k}) \rightarrow (R_{\text{gr}}, \text{Fr}) - \text{mod}$, where $(R_{\text{gr}}, \text{Fr}) - \text{mod}$ denotes (cohomologically) graded R -modules equipped with a Fr-action. The cohomological degree $r + k$ on the right-hand side of (5.8) appears since the functor $\mathbb{H} : D_{\text{perf}}(R, \text{Fr}) \rightarrow (R_{\text{gr}}, \text{Fr}) - \text{mod}$ sends the sum of the two gradings to the cohomological one; see [BY13, Corollary B.4.1(1)].

7.1. Realization functors. We record two results on realization functors taken from [AMRW19]. See also [Bei87, Rid13]. For a definition of a filtered version of a triangulated category, see [Bei87, Definition A.1].

Proposition 7.2 ([AMRW19, Proposition 2.2]). *Let \mathcal{T} be a triangulated category that admits a filtered version $\tilde{\mathcal{T}}$ and let $\mathcal{A} \subset \mathcal{T}$ be a full additive subcategory that admits no negative self-extensions. Then there is a functor of triangulated categories*

$$\text{real} : K^b \mathcal{A} \rightarrow \mathcal{T}$$

whose restriction to \mathcal{A} is the inclusion functor.

Proposition 7.3 ([AMRW19, Proposition 2.3]). *Let \mathcal{T}_1 and \mathcal{T}_2 be triangulated categories admitting a filtered version, and let $\mathcal{A}_1 \subset \mathcal{T}_1$ and $\mathcal{A}_2 \subset \mathcal{T}_2$ be two full additive subcategories admitting no negative self-extensions. Let $F : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ be a triangulated functor that restricts to an additive functor $F_0 : \mathcal{A}_1 \rightarrow \mathcal{A}_2$. If F lifts to a functor $\tilde{F} : \tilde{\mathcal{T}}_1 \rightarrow \tilde{\mathcal{T}}_2$, then the following diagram commutes up to natural isomorphism:*

$$\begin{array}{ccc} K^b \mathcal{A}_1 & \xrightarrow{\text{real}} & \mathcal{T}_1 \\ K^b F_0 \downarrow & & \downarrow F \\ K^b \mathcal{A}_2 & \xrightarrow{\text{real}} & \mathcal{T}_2 \end{array}$$

7.2. Semisimple complexes. Let $\text{Semis}_B(X) \subset D_B^b(X, \mathbb{k})$ denote the additive subcategory generated by semisimple complexes pure of weight 0. Thus, an object of $\text{Semis}_B(X)$ is a direct sum of the twisted intersection cohomology sheaves $\mathcal{IC}_w[n](n/2)$ for $w \in W$ and $n \in \mathbb{Z}$.

Recall from Section 3.1 that $S_w \subset B_{\underline{w}}$ denotes the indecomposable Soergel bimodule indexed by $w \in W$.

Lemma 7.4. *For $\mathcal{F}, \mathcal{G} \in \text{Semis}_B(X)$, the Ext-group $\text{Ext}^i(\mathcal{F}, \mathcal{G})$ is pure of weight i for all $i \in \mathbb{Z}$.*

Proof. Follows from [BY13, Lemma 3.1.5]. \square

Proposition 7.5. *The hypercohomology functor induces an equivalence of additive categories*

$$\mathbb{H}_B : \text{Semis}_B(X) \rightarrow \mathbb{S}\text{Bim},$$

enriched over $R \otimes R$, and sending \mathcal{IC}_w to the shifted Soergel bimodule $S_w\{-\ell(w)\}$ and the twist $[n](n/2)$ to the change of grading $\{-n/2\}$.

Proof. By a well-known result of Soergel [Soe07] (see also [EW16, Section 1.3] or [EMTW20, Section 16.1]), we have $\mathbb{H}_B(\mathcal{IC}_w) \cong S_w\{-\ell(w)/2\}$ as a graded $R \otimes R$ -module. By Lemma 7.4, the cohomological and weight gradings on $\mathbb{H}_B(\mathcal{IC}_w)$ agree, and furthermore $\mathcal{IC}_w[n](n/2)$ is sent to $S_w\{-(\ell(w) + n)/2\}$. The result can then be deduced from [BY13, Proposition 3.1.6]. \square

The following result is well-known; see [AR13] and [BY13, Lemma B.1.1].

Lemma 7.6. *For $\mathcal{F}, \mathcal{G} \in \text{Semis}_B(X)$, the action of Fr on $\text{Ext}^\bullet(\mathcal{F}, \mathcal{G})$ is semisimple. Furthermore, we have*

$$(7.1) \quad \text{hom}_{D_B^b(X, \mathbb{k})}(\mathcal{F}, \mathcal{G}) \cong \text{Ext}^0(\mathcal{F}, \mathcal{G}) \cong \text{Ext}^{0, (0)}(\mathcal{F}, \mathcal{G})$$

and $\text{ext}_{D_B^b(X, \mathbb{k})}^i(\mathcal{F}, \mathcal{G}) = 0$ for $i < 0$.

7.3. The mixed derived category. Following [AR16b], we define the *mixed derived category*

$$D_B^{\text{mix}}(X) := K^b(\text{Semis}_B(X))$$

to be the homotopy category of cochain complexes in $\text{Semis}_B(X)$. Define the *Tate twist* of $D_B^{\text{mix}}(X)$ by

$$\langle n \rangle := \{-n/2\}[-n],$$

where $[n] : K^b(\text{Semis}_B(X)) \rightarrow K^b(\text{Semis}_B(X))$ is the cohomological shift functor, and $\{n/2\} : \text{Semis}_B(X) \rightarrow \text{Semis}_B(X)$ is the autoequivalence $\mathcal{F} \mapsto \mathcal{F}[-n](-n/2)$.

By Proposition 7.5, we have an equivalence of triangulated categories

$$K^b\mathbb{H}_B : D_B^{\text{mix}}(X) \rightarrow K^b\text{SBim}.$$

Theorem 7.7. *There exists a triangulated realization functor*

$$\text{real} : D_B^{\text{mix}}(X) \rightarrow D_B^b(X, \mathbb{k}),$$

restricting to the inclusion on $\text{Semis}_B(X)$, sending

$$[n] \rightarrow [n], \quad \langle n \rangle \rightarrow (n/2), \quad \{n/2\} \rightarrow [-n](-n/2),$$

such that the composition

$$\kappa := \omega \circ \text{real}$$

satisfies

$$(7.2) \quad \text{Hom}_{D_B^{\text{mix}}(X)}(\mathcal{F}, \mathcal{G}\langle n \rangle) \cong \text{Ext}^{0, (n/2)}(\text{real}\mathcal{F}, \text{real}\mathcal{G}),$$

$$(7.3) \quad \bigoplus_{n \in \mathbb{Z}} \text{Hom}_{D_B^{\text{mix}}(X)}(\mathcal{F}, \mathcal{G}\langle n \rangle) \cong \text{Ext}^0(\text{real}\mathcal{F}, \text{real}\mathcal{G}) = \text{Hom}_{D_B^b(X, \mathbb{k})}(\kappa\mathcal{F}, \kappa\mathcal{G}).$$

Furthermore, all functors are compatible with the $R \otimes R$ action on the corresponding Ext^\bullet -groups.

Proof. By Lemma 7.6, $\text{Semis}_B(G/B)$ has no negative self-extensions. Since $D_B^b(X, \mathbb{k})$ admits a filtered version (see [Bei87]), we may apply Proposition 7.2 to obtain a realization functor

$$\text{real} : D_B^{\text{mix}}(X) \rightarrow D_B^b(X, \mathbb{k})$$

that restricts to the inclusion on $\text{Semis}_B(G/B)$. For $\mathcal{F}, \mathcal{G} \in \text{Semis}_B(X)$, the isomorphism (7.3) follows from (7.1) while (7.2) follows from (7.1) and (7.3), since $\langle n \rangle : D_B^{\text{mix}}(X) \rightarrow D_B^{\text{mix}}(X)$ is sent to $(n/2) : D_B^b(X, \mathbb{k}) \rightarrow D_B^b(X, \mathbb{k})$ by the realization functor real . For $\mathcal{F}, \mathcal{G} \in K^b\text{Semis}_B(X)$, (7.2)–(7.3) are proven by double induction on the lengths of chain complexes representing \mathcal{F}, \mathcal{G} . See [Rid13, Section 4.1] for a detailed argument. \square

7.4. Standard objects and Rouquier complexes.

Proposition 7.8. *The composition $\text{real} \circ \mathbb{H}_B^{-1} : K^b\text{SBim} \rightarrow D_B^b(X, \mathbb{k})$ takes $F^\bullet(w)$ to $\Delta_w(-\ell(w)/2)$ and $F^\bullet(v)^{-1}$ to $\nabla_v(\ell(v)/2)$.*

Note that Proposition 5.4 follows nearly immediately from Proposition 7.8 and (7.2)–(7.3): since the realization functor sends $[k]\{-r/2\}$ to $[k+r](r/2)$, we have

$$\begin{aligned} H^{k, (r/2)}(HH^0(F_{v,w}^\bullet)) &\cong \text{Hom}_{K^b\text{SBim}}(R, F_{v,w}^\bullet[k]\{-r/2\}) \\ &\cong \text{Ext}^{r+k, (r/2)}(\Delta_e, \Delta_w(-\ell(w)/2) \star \nabla_v(\ell(v)/2)) \\ &\cong \text{Ext}^{r+k, (r/2)}(\Delta_v(-\ell(v)/2), \Delta_w(-\ell(w)/2)) \\ &\cong \text{Ext}^{r+k, ((r-\ell_{v,w})/2)}(\Delta_v, \Delta_w). \end{aligned}$$

The second isomorphism above is obtained from the adjointness of the convolution functors $(-)\star \Delta_v(-\ell(v)/2)$ and $(-)\star \nabla_v(\ell(v)/2)$ to be presently explained; see Lemma 7.10 for a stronger statement, and see also (3.5)–(3.6) for a similar statement on the Soergel bimodule side.

Proof of Proposition 7.8. We prove the claim for $F^\bullet(w)$. The claim for $F^\bullet(v)^{-1}$ is similar.

There is a monoidal structure $\star : D_B^b(X, \mathbb{k}) \times D_B^b(X, \mathbb{k}) \rightarrow D_B^b(X, \mathbb{k})$ obtained by convolution; see for example [BY13, Section 3.2] or [AR16b, Section 4.3]. By [BY13, Proposition 3.2.5], the additive subcategory $\text{Semis}_B(X) \subset D_B^b(X, \mathbb{k})$ is preserved by convolution. According to [BY13, Proposition 3.2.1], convolution \star is sent by \mathbb{H}_B to the tensor product operation on $\mathbb{S}\text{Bim}$. Note that the *derived* tensor product in [BY13] reduces to the tensor product on $\mathbb{S}\text{Bim}$ since all Soergel bimodules are free as left (or right) R -modules.

For a simple generator $s \in S$, let $\pi_s : X = G/B \rightarrow G/P_s$ denote the projection to the partial flag variety, where $P_s \supset B$ denotes a minimal parabolic subgroup, and let $\theta_s : D_B^b(X, \mathbb{k}) \rightarrow D_B^b(X, \mathbb{k})$ denote the composition $\theta = \pi_s^* \pi_{s,*}$. By [BY13, Lemma 3.2.7] (see also [AR16b, Lemma 4.3]), we have a natural isomorphism of functors $\theta_s \cong (-) \star \mathcal{IC}_s[-1](-1/2)$, and θ_s restricts to an endofunctor $\theta_s : \text{Semis}_B(X) \rightarrow \text{Semis}_B(X)$. It is well-known [Soe90, Korollar 2] that the equivalence $\mathbb{H}_B : \text{Semis}_B(X) \rightarrow \mathbb{S}\text{Bim}$ takes the functor θ_s to the functor $R \otimes_{R^s} (-)$.

Now, there is a natural morphism of functors $\theta_s \rightarrow \text{id}$ arising from the adjunction of π_s^* and $\pi_{s,*}$. The map $B_s \rightarrow R$ in (3.4) arises by an analogous adjunction; see [Rou04, Section 3]. Now, $\mathcal{IC}_e \cong \Delta_e$ and $\mathbb{H}_B(\mathcal{IC}_e) = R$, and the morphism $\theta_s \rightarrow \text{id}$ applied to \mathcal{IC}_e fits into the distinguished triangle

$$\mathcal{IC}_s[-1](-1/2) \rightarrow \mathcal{IC}_e \rightarrow \Delta_s(-1/2)$$

in $D_B^b(X, \mathbb{k})$; see for example [BY13, (C.4)] or [AR16b, Lemma 4.1]. It follows that we have

$$(7.4) \quad \text{real} \circ \mathbb{H}_B^{-1}(F^\bullet(s)) = \Delta_s(-1/2).$$

See also [Rou04, Proposition 5.3]. This establishes Proposition 7.8 in the case $\ell(w) = 1$. We then obtain a natural isomorphism of functors

$$(7.5) \quad ((-) \star \Delta_s(-1/2)) \circ \text{real} \circ \mathbb{H}_B^{-1} \cong \text{real} \circ \mathbb{H}_B^{-1} \circ ((-) \otimes F(s))$$

from $\text{K}^b \mathbb{S}\text{Bim}$ to $D_B^b(X, \mathbb{k})$.

On the other hand, it is known ([BY13, Lemma 3.2.2]) that if $w = s_1 s_2 \cdots s_\ell$ is a reduced decomposition then

$$\Delta_w \cong \Delta_{s_1} \star \Delta_{s_2} \star \cdots \star \Delta_{s_\ell}$$

is in $D_B^b(X, \mathbb{k})$. Combining (7.4) with (7.5), we find

$$\begin{aligned} \text{real} \circ \mathbb{H}_B^{-1}(F^\bullet(w)) &\cong \text{real} \circ \mathbb{H}_B^{-1}(F^\bullet(s_1) \otimes \cdots \otimes F^\bullet(s_\ell)) \\ &\cong \Delta_{s_1}(-1/2) \star \cdots \star \Delta_{s_\ell}(-1/2) \cong \Delta_w(-\ell(w)/2). \end{aligned} \quad \square$$

7.5. Proof of Proposition 7.1.

7.5.1. A functor $F : D^b(Y, \mathbb{k}) \rightarrow D^b(Z, \mathbb{k})$ is called *geometric* ([AR13, Definition 6.6]) if there is a natural transformation

$$\text{RHom}_Y(\mathcal{F}, \mathcal{G}) \rightarrow \text{RHom}_Z(F\mathcal{F}, F\mathcal{G})$$

for $\mathcal{F}, \mathcal{G} \in D^b(Y, \mathbb{k})$. We shall apply the notion of a geometric functor for B -equivariant derived categories.

Lemma 7.9. *The endofunctors $(-) \star \Delta_s(-1/2)$, $(-) \star \nabla_s(1/2) : D_B^b(X, \mathbb{k}) \rightarrow D_B^b(X, \mathbb{k})$ are geometric.*

Proof. This is stated for the affine Grassmannian case in [AR13, Proposition 12.2]. The same proof applies in the flag variety case. \square

Lemma 7.10. *We have*

$$\text{RHom}(\Delta_v, \Delta_w) = \text{RHom}(\Delta_e, \Delta_w \star \nabla_v)$$

inside $D_B^b(\text{pt}, \mathbb{k})$.

Proof. By Lemma 7.9, for $\mathcal{F}, \mathcal{G} \in D_B^b(X, k)$. we have a map

$$(7.6) \quad \mathrm{RHom}(\mathcal{F}, \mathcal{G}) \rightarrow \mathrm{RHom}(\mathcal{F} \star \nabla_s, \mathcal{G} \star \nabla_s)$$

and a map

$$(7.7) \quad \mathrm{RHom}(\mathcal{F} \star \nabla_s, \mathcal{G} \star \nabla_s) \rightarrow \mathrm{RHom}(\mathcal{F} \star \nabla_s \star \Delta_s, \mathcal{G} \star \nabla_s \star \Delta_s)$$

inside $D_B^b(\mathrm{pt}, \mathbb{k})$. Convolution is associative and $\nabla_s \star \Delta_s \cong \Delta_e$ ([AR16b, Proposition 4.4]), and also $(-) \star \Delta_e$ is the identity functor. So composing (7.6) and (7.7), we get an automorphism $\mathrm{RHom}(\mathcal{F}, \mathcal{G}) \cong \mathrm{RHom}(\mathcal{F}, \mathcal{G})$ inside $D_B^b(\mathrm{pt}, \mathbb{k})$. It follows that $\mathrm{RHom}(\mathcal{F}, \mathcal{G}) \cong \mathrm{RHom}(\mathcal{F} \star \nabla_s, \mathcal{G} \star \nabla_s)$. Choosing $\mathcal{F} = \Delta_v$ and $\mathcal{G} = \Delta_w$ and repeatedly applying this we obtain the required statement. \square

We remark that we could have defined $\mathrm{RHom}(\Delta_v, \Delta_w)$ and $\mathrm{RHom}(\Delta_e, \Delta_w \star \nabla_v)$ as objects in $D_{B \times B}^b(\mathrm{pt}, \mathbb{k})$, but Lemma 7.10 would not hold. The functors $(-) \star \Delta_s$ and $(-) \star \nabla_s$ “commute” with only one of the B -actions.

7.5.2. Let $\iota_e : B \hookrightarrow G$ be the inclusion and $\pi : G/B \rightarrow \{e\} = \mathrm{pt}$ be the projection. These maps are B -equivariant. Now, the sheaf Δ_e is supported on a single point $\{e\} \subset G/B$. Thus, for any $\mathcal{F} \in D_B^b(X, \mathbb{k})$, the object $\mathrm{R}\mathcal{H}om(\Delta_e, \mathcal{F}) \in D_B^b(X, \mathbb{k})$ is also supported on $\{e\}$. The pullback ι_e^* and pushforward $\iota_{e,*}$ are inverse equivalences of categories between $D_B^b(\{e\}, \mathbb{k})$ and the full subcategory of $D_B^b(X, \mathbb{k})$ consisting of objects whose cohomology sheaves are supported on $\{e\}$. Inside $D_B^b(\{e\}, \mathbb{k}) = D_B^b(\mathrm{pt}, \mathbb{k})$, we thus have

$$(7.8) \quad \begin{aligned} \mathrm{RHom}_X(\Delta_e, \mathcal{F}) &\cong \pi_* \mathrm{R}\mathcal{H}om_X(\Delta_e, \mathcal{F}) \\ &\cong \iota_e^* \mathrm{R}\mathcal{H}om_X(\Delta_e, \mathcal{F}) = \mathrm{R}\mathcal{H}om_{\{e\}}(\mathbb{k}, \iota_e^* \mathcal{F}) = \iota_e^* \mathcal{F}. \end{aligned}$$

7.5.3. The equivalence $D_B^b(\mathrm{pt}, \mathbb{k}) \cong D_{\mathrm{perf}}(R, \mathrm{Fr})$ (resp., $D_{B \times B}^b(\mathrm{pt}, \mathbb{k}) \cong D_{\mathrm{perf}}(R \otimes R, \mathrm{Fr})$) sends objects pure of weight 0 to objects in $D_{\mathrm{perf}}(R, \mathrm{Fr})$ (resp., $D_{\mathrm{perf}}(R \otimes R, \mathrm{Fr})$) concentrated in cohomological degree 0. For $\mathcal{F} \in \mathrm{Semis}_B(X)$, we thus view the Soergel bimodule $\mathbb{H}_B(\mathcal{F})$ as sitting inside $D_{\mathrm{perf}}(R \otimes R, \mathrm{Fr})$ in cohomological degree 0, that is, as an object in $\mathrm{Mod}(R \otimes R, \mathrm{Fr})$.

Lemma 7.11. *For $\mathcal{F} \in \mathrm{Semis}_B(G/B)$, we have an isomorphism*

$$\iota_e^* \mathcal{F} \cong \mathrm{Hom}_{R \otimes R}(R, \mathbb{H}_B(\mathcal{F}))$$

inside $\mathrm{Mod}(R, \mathrm{Fr})$.

Proof. For $\mathcal{F} \in \mathrm{Semis}_B(G/B)$, we have that $\iota_e^* \mathcal{F} = \iota_e^! \mathcal{F}$ is again pure of weight 0. Thus, $\iota_e^* \mathcal{F} \in D_B^b(\mathrm{pt}, \mathbb{k})$ can be identified with an element of $\mathrm{Mod}(R, \mathrm{Fr})$. By [BY13, Proposition 3.1.6] and (7.8), we have

$$\mathrm{Hom}_{R \otimes R}(R, \mathbb{H}_B(\mathcal{F})) \cong \mathbb{H}_B(\mathrm{RHom}(\Delta_e, \mathcal{F})) \cong \mathbb{H}_B(\iota_e^* \mathcal{F}).$$

(Since $\iota_e^* \mathcal{F}$ is pure of weight 0, $\mathbb{H}_B(\iota_e^* \mathcal{F})$ is simply the corresponding object in $\mathrm{Mod}(R, \mathrm{Fr})$.) We conclude that $\iota_e^* \mathcal{F} \cong \mathrm{Hom}_{R \otimes R}(R, \mathbb{H}_B(\mathcal{F}))$ inside $\mathrm{Mod}(R, \mathrm{Fr})$ and the result follows. \square

We remark that $\mathrm{Hom}_{R \otimes R}(R, \mathbb{H}_B(\mathcal{F}))$ is free as an R -module; cf. [BY13, Lemma 3.1.5]. See also Remark 3.4.

7.5.4. Recall that we have a realization functor $\mathrm{real} : \mathrm{K}^b \mathrm{Semis}_B(X) \rightarrow D_B^b(X, \mathbb{k})$. We now have two functors from $\mathrm{K}^b \mathrm{Semis}_B(X)$ to $D_B^b(\mathrm{pt}, \mathbb{k})$. The functor

$$(7.9) \quad \iota_e^* \circ \mathrm{real} : \mathrm{K}^b \mathrm{Semis}_B(X) \rightarrow D_B^b(X, \mathbb{k}) \rightarrow D_B^b(\mathrm{pt}, \mathbb{k})$$

and the functor

$$(7.10) \quad \mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -) \circ \mathrm{K}^b \mathbb{H}_B : \mathrm{K}^b \mathrm{Semis}_B(X) \rightarrow \mathrm{K}^b \mathrm{SBim} \rightarrow D_B^b(\mathrm{pt}, \mathbb{k}).$$

We explain the last functor $\mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -) : \mathrm{K}^b \mathrm{SBim} \rightarrow D_B^b(\mathrm{pt}, \mathbb{k})$. Let $\mathrm{Free}(R, \mathrm{Fr})$ denote the category of finitely generated free R -modules equipped with an action of Fr . The functor

$\mathrm{Hom}_{R \otimes R}(R, -)$ takes $\mathbb{S}\mathrm{Bim}$ to $\mathrm{Free}(R, \mathrm{Fr})$, and $\mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -)$ takes $\mathrm{K}^b \mathbb{S}\mathrm{Bim}$ to $\mathrm{K}^b \mathrm{Free}(R, \mathrm{Fr})$. We have an inclusion $\mathrm{Free}(R, \mathrm{Fr}) \rightarrow \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k})$. Applying Proposition 7.2, we obtain a triangulated functor $\mathrm{real} : \mathrm{K}^b(\mathrm{Free}(R, \mathrm{Fr})) \rightarrow \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k})$. Composing $\mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -)$ with real we obtain $\mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -) : \mathrm{K}^b \mathbb{S}\mathrm{Bim} \rightarrow \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k})$.

By Lemma 7.11, the two triangulated functors (7.9) and (7.10) agree on the subcategory $\mathrm{Semis}_B(X) \subset \mathrm{K}^b \mathrm{Semis}_B(X)$, sending $\mathrm{Semis}_B(X)$ to $\mathrm{Free}(R, \mathrm{Fr}) \subset \mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr}) \cong \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k})$. Denoting this restriction by $L : \mathrm{Semis}_B(X) \rightarrow \mathrm{Free}(R, \mathrm{Fr})$, we apply Proposition 7.3 to deduce that both triangulated functors are isomorphic to

$$\mathrm{real} \circ \mathrm{K}^b L : \mathrm{K}^b \mathrm{Semis}_B(X) \rightarrow \mathrm{K}^b \mathrm{Free}(R, \mathrm{Fr}) \rightarrow \mathrm{D}_{\mathrm{perf}}(R, \mathrm{Fr}) \cong \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k}).$$

Thus,

$$(7.11) \quad \iota_e^* \circ \mathrm{real} \cong \mathrm{K}^b \mathrm{Hom}_{R \otimes R}(R, -) \circ \mathrm{K}^b \mathbb{H}_B.$$

Remark 7.12. The essential image of the realization functor $\mathrm{K}^b \mathrm{Free}(R, \mathrm{Fr}) \rightarrow \mathrm{D}_{\mathrm{perf}}^b(R, \mathrm{Fr})$ is a subcategory of $\mathrm{D}_{\mathrm{perf}}^b(R, \mathrm{Fr})$ equivalent to the infinitesimal extension of $\mathrm{K}^b \mathrm{Free}(R, \mathrm{Fr})$, in the sense of [AR13].

7.5.5. **Conclusion.** By (7.11) and Proposition 7.8, we have

$$\iota_e^*(\Delta_w(-\ell(w)/2) \star \nabla_v(\ell(v)/2)) \cong \mathrm{Hom}_{R \otimes R}(R, F_{v,w}^\bullet)$$

inside $\mathrm{D}_{\mathrm{perf}}^b(R, \mathrm{Fr})$. By Lemma 7.10 and (7.8), we find

$$\iota_e^*(\Delta_w(-\ell(w)/2) \star \nabla_v(\ell(v)/2)) \cong \mathrm{RHom}_X(\Delta_v(-\ell(v)/2), \Delta_w(-\ell(w)/2)).$$

This finishes the proof of Proposition 7.1, as well as of Proposition 5.4 and Theorem 1.17.

8. ORDINARY COHOMOLOGY, KOSZUL DUALITY, AND VERMA MODULES

The goal of this section is to prove Theorems 1.16, 1.18 and 1.20.

8.1. **Ordinary cohomology.** We have a forgetful functor $\mathrm{For} : \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k}) \rightarrow \mathrm{D}^b(\mathrm{pt}, \mathbb{k})$, and a commutative diagram (see [BL94] and [BY13, Proposition B.3.1])

$$(8.1) \quad \begin{array}{ccc} \mathrm{D}_B^b(\mathrm{pt}, \mathbb{k}) & \xrightarrow{\mathrm{For}} & \mathrm{D}^b(\mathrm{pt}, \mathbb{k}) \\ \cong \downarrow & & \downarrow \cong \\ \mathrm{D}_{\mathrm{perf}}^b(R, \mathrm{Fr}) & \xrightarrow{\otimes_R^L \mathbb{k}} & \mathrm{D}^b(\mathbb{k}, \mathrm{Fr}). \end{array}$$

Here, $\mathrm{D}^b(\mathbb{k}, \mathrm{Fr})$ is the derived category of finite-dimensional \mathbb{k} -vector spaces equipped with an Fr -action with integer weights. Applying For to Proposition 7.1, we obtain the following.

Proposition 8.1. *We have*

$$\mathrm{RHom}_{\mathrm{D}^b(X, \mathbb{k})}(\Delta_v^{(B)}(-\ell(v)/2), \Delta_w^{(B)}(-\ell(w)/2)) \cong \mathrm{Hom}_{R \otimes R}(R, F_{v,w}^\bullet) \otimes_R \mathbb{k}$$

inside $\mathrm{D}^b(\mathbb{k}, \mathrm{Fr})$. □

Here, $\Delta_v^{(B)} = \mathrm{For}(\Delta_v) \in \mathrm{D}_{(B)}^b(X, \mathbb{k})$ denotes the ordinary standard object in the Borel-constructible derived category, and the derived tensor product $\otimes_R^L \mathbb{k}$ is replaced by the usual tensor product since $\mathrm{Hom}_{R \otimes R}(R, F_{v,w}^\bullet)$ is free as an R -module. Taking the hypercohomology of both sides of Proposition 8.1, we obtain the following.

Corollary 8.2. *For all $v \leq w \in W$, and all $k, r \in \mathbb{Z}$, we have an isomorphism*

$$(8.2) \quad \mathrm{Ext}^{r+k, ((r-\ell_{v,w})/2)}(\Delta_v^{(B)}, \Delta_w^{(B)}) \cong H^{k, (r/2)}(\mathrm{HH}_{\mathbb{k}}^0(F_{v,w}^\bullet))$$

of \mathbb{k} -vector spaces. (In particular, both sides are zero for odd r .)

Here we set $HH_{\mathbb{k}}^0(B) := HH^0(B) \otimes_R \mathbb{k}$, similarly to (1.18). For the shift in cohomological degree, see the discussion after Proposition 7.1.

Proof of Theorem 1.16. The non-equivariant version of Proposition 5.3 is given in [RSW14, Proposition 4.2.1]. Similarly to (6.1)–(6.2), we get

$$\mathrm{Ext}^{m, ((r-\ell_{v,w})/2)}(\Delta_v^{(B)}, \Delta_w^{(B)}) \cong H_c^{m+\ell_{v,w}, (r/2)}(R_{v,w}^\circ, \mathbb{k}) \quad \text{for all } m, r \in \mathbb{Z}.$$

Poincaré duality (4.5) allows one to translate the compactly supported cohomology into the ordinary cohomology:

$$(8.3) \quad \mathrm{Ext}^{m, ((r-\ell_{v,w})/2)}(\Delta_v^{(B)}, \Delta_w^{(B)}) \cong H^{\ell_{v,w}-m, (\ell_{v,w}-r/2)}(R_{v,w}^\circ, \mathbb{k}).$$

Combining (8.2)–(8.3) with Koszul duality (1.23) proved in the next section, and switching from working over \mathbb{k} to working over \mathbb{C} via Remark 5.1 and Proposition 5.5, we get

$$\begin{aligned} \dim_{\mathbb{C}} H^{k, (r/2)}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)) &= \dim_{\mathbb{C}} H^{\ell_{v,w}-k-r, (\ell_{v,w}-r/2, \ell_{v,w}-r/2)}(R_{v,w}^\circ, \mathbb{C}) \\ &= \dim_{\mathbb{C}} H^{-k, (r/2, r/2)}(R_{v,w}^\circ, \mathbb{C}). \end{aligned} \quad \square$$

Proof of (1.24). By the Künneth formula and Corollary 4.8, we have

$$H^\bullet(R_{v,w}^\circ) \cong H^\bullet(R_{v,w}^\circ/T) \otimes_{\mathbb{C}} H^\bullet(T).$$

The space $H^\bullet(T)$ is 2^{n-1} -dimensional, and the mixed Hodge polynomials are related as

$$(8.4) \quad \mathcal{P}(R_{v,w}^\circ; q, t) = \left(q^{\frac{1}{2}} + t^{\frac{1}{2}}\right)^{n-1} \cdot \mathcal{P}(R_{v,w}^\circ/T; q, t).$$

This implies (1.10). Next, we claim

$$(8.5) \quad \mathcal{P}(R_{v,w}^\circ/T; q, t) = \left(q^{\frac{1}{2}} t^{\frac{1}{2}}\right)^{\chi(\beta_{v,w})} \mathcal{P}_{\mathrm{KR}; \mathbb{C}}^{\mathrm{top}}(\hat{\beta}_{v,w}; q, t).$$

First, combining Theorem 1.16 with Koszul duality (1.23) (to be proved below in Section 8.2), we find

$$(8.6) \quad H^{k, (p,p)}(R_{v,w}^\circ, \mathbb{C}) \cong H^{-\ell_{v,w}-k+2p, (\ell_{v,w}-p)}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)).$$

Setting $k' := -\ell_{v,w} - k + 2p$ and $p' := \ell_{v,w} - p$, we find $p = \ell_{v,w} - p'$ and $k = \ell_{v,w} - k' - 2p'$. Plugging this into (4.6) and applying (8.6), we get

$$\begin{aligned} \mathcal{P}(R_{v,w}^\circ; q, t) &= \sum_{k,p \in \mathbb{Z}} q^{p-\frac{k}{2}} t^{\frac{\ell_{v,w}-k}{2}} \dim H^{k, (p,p)}(R_{v,w}^\circ, \mathbb{C}) \\ &= \sum_{k', p' \in \mathbb{Z}} q^{\frac{\ell_{v,w}+k'}{2}} t^{p'+\frac{k'}{2}} H^{k', (p')}(HH_{\mathbb{C}}^0(F_{v,w}^\bullet)). \end{aligned}$$

On the other hand, rewriting (3.16), we see that the right-hand side of (8.5) is given by

$$\left(q^{\frac{1}{2}} + t^{\frac{1}{2}}\right)^{1-n} \sum_{k', p' \in \mathbb{Z}} q^{\frac{\ell_{v,w}+k'}{2}} t^{p'+\frac{k'}{2}} \dim H^{k', (p')}(HH_{\mathbb{C}}^0(F^\bullet(\beta))).$$

Together with (8.4), this finishes the proof of (8.5). Finally, (1.24) follows by comparing (8.5) with (the Richardson version of) (4.9). \square

Since the R -action on $H_{T,c}^\bullet(R_{v,w}^\circ)$ is trivial (that is, \mathfrak{h}^* acts by zero), by Theorem 1.17, the R -action on $HHH^0(F_{v,w}^\bullet)$ is also trivial. Alternatively, for $W = S_n$ and any knot $\hat{\beta}$, the R -action on $HHH^0(F^\bullet(\beta))$ is trivial by Corollary 3.10. It thus follows that we have an isomorphism of bigraded \mathbb{C} -modules

$$(8.7) \quad HHH_{\mathbb{C}}^0(F^\bullet(\beta)) \cong \mathrm{Tor}_{\bullet}^R(\mathbb{C}, HHH^0(F^\bullet(\beta))).$$

We conjecture that (8.7) holds for all W and all $\beta \in \mathcal{B}_W$. This would follow from (8.1) if $HH^0(F^\bullet(\beta))$ and $HHH^0(F^\bullet(\beta))$ were known to be equivalent in $D_{\text{perf}}^b(R, \text{Fr})$.

8.2. Koszul duality and q, t -symmetry. We prove Theorem 1.18. By [BY13, Equation (5.2), Theorem 5.3.1, and Remark 5.3.2], for $k, r \in \mathbb{Z}$ and $v \leq w \in W$, we get an isomorphism

$$\text{Ext}^{k, (r/2)}(\Delta_v^{(B)}, \Delta_w^{(B)}) \cong \text{Ext}^{k-r, (-r/2)}(\Delta_{v^{-1}}^{(B)}, \Delta_{w^{-1}}^{(B)})$$

of vector spaces. By (8.3) and Proposition 5.5, this implies

$$(8.8) \quad H^{k, (r/2, r/2)}(R_{v,w}^\circ, \mathbb{C}) \cong H^{\ell_{v,w}+k-r, (\ell_{v,w}-r/2, \ell_{v,w}-r/2)}(R_{v^{-1}, w^{-1}}^\circ, \mathbb{C}).$$

The only difference between (8.8) and the desired result (1.23) is the appearance of v^{-1} and w^{-1} on the right-hand side. In fact, it is not hard to see that the Richardson varieties $R_{v,w}^\circ$ and $R_{v^{-1}, w^{-1}}^\circ$ are isomorphic. Indeed, recall from Lemma 4.4 that we have an isomorphism $N_{v,w}^\circ \cong R_{v,w}^\circ$. The map $g \mapsto g^{-1}$ restricts to an isomorphism $N_{v,w}^\circ \cong N_{v^{-1}, w^{-1}}^\circ$ (choosing \dot{v}^{-1} as the representative for v^{-1}). By (4.3), we get an isomorphism $R_{v,w}^\circ \xrightarrow{\sim} R_{v^{-1}, w^{-1}}^\circ$. \square

8.3. Extensions of Verma modules. We prove Theorem 1.20. First, we explain the bigrading on $\text{Ext}^\bullet(M_v, M_w)$. Out of the several equivalent descriptions listed in [BGS96], the most convenient one for us is given in [BGS96, Section 4.4]: the bigraded vector spaces $\text{Ext}^\bullet(M_v, M_w)$ and $\text{Ext}^\bullet(\Delta_v^{(B)}, \Delta_w^{(B)})$ are isomorphic (after changing the coefficients from \mathbb{C} to \mathbb{k}), and the bigrading on $\text{Ext}^\bullet(M_v, M_w)$ comes from the bigrading on $\text{Ext}^\bullet(\Delta_v^{(B)}, \Delta_w^{(B)})$ via Frobenius weights (5.3):

$$\text{Ext}^{k, (r/2)}(M_v, M_w) := \text{Ext}^{k, (r/2)}(\Delta_v^{(B)}, \Delta_w^{(B)}).$$

See also [RSW14, Equation (1.1.1)].

The result follows by combining (8.3) with Koszul duality (1.23). \square

9. CATALAN NUMBERS ASSOCIATED TO POSITROID VARIETIES

Our results give an embedding of the rational q, t -Catalan numbers $C_{k, n-k}(q, t)$ into a family of q, t -polynomials $\mathcal{P}(\mathcal{X}_f^\circ; q, t) \in \mathbb{N}[q^{\frac{1}{2}}, t^{\frac{1}{2}}]$ (all of which are q, t -symmetric and q, t -unimodal), indexed by $f \in \mathbf{B}_{k, n}^{c=1}$. The goal of this section is to give a combinatorial interpretation for a specialization of $\mathcal{P}(\mathcal{X}_f^\circ; q, t)$.

Definition 9.1. For $f \in \mathbf{B}_{k, n}^{c=1}$, define the f -Catalan number $C_f \in \mathbb{Z}$ as the specialization

$$C_f := \mathcal{P}(\mathcal{X}_f^\circ; q, t) \Big|_{q^{\frac{1}{2}}=1, t^{\frac{1}{2}}=-1}.$$

Alternatively, C_f is the $q = 1$ specialization of the point count polynomial $\#\mathcal{X}_f^\circ(\mathbb{F}_q)$, and we also have $C_f = P_f^{\text{top}}(1)$, where the polynomial $P_f^{\text{top}}(q)$ is defined in Theorem 1.13.

In particular, $C_{f_{k, n}} = C_{k, n-k}(1, 1) = \#\text{Dyck}_{k, (n-k)}$ is the usual rational Catalan number when $\gcd(k, n) = 1$.

Recall from Proposition 1.10 that each $f = f_{v, w} \in \mathbf{B}_{k, n}$ corresponds to a pair $v \leq w \in S_n$ such that w is k -Grassmannian. The set of k -Grassmannian permutations in S_n is well known to be in bijection with the set of Young diagrams that fit inside a $k \times (n-k)$ -rectangle. Let λ be such a Young diagram. We are going to consider fillings of boxes of λ with *crossings* $\begin{array}{c} \oplus \\ \oplus \end{array}$ and *elbows* $\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}$. An example is given in Figure 4. Each such filling D gives rise to a permutation u_D , obtained as follows. Consider paths labeled by $1, 2, \dots, n$ entering from the southeast boundary of λ , where the labels increase in the northeast direction. The paths follow crossings and elbows until they exit through the northwest boundary of λ . Recording the positions of outgoing edges, one obtains the permutation u_D (cf. Figure 4).




FIGURE 4. For the two fillings on the left, we have $\lambda = (5, 4, 2)$ and $f = wv^{-1} = 35148276$. The two fillings on the right do not satisfy the distinguished condition: the specific elbow violating the condition is shaded in red.




FIGURE 5. The sets $\text{Deo}_{f_{k,n}}^{\max}$ and $\text{Dyck}_{k,n-k}$ have the same cardinality by Proposition 9.5. Compare with Figure 1.

Definition 9.2. Let λ be a Young diagram fitting in a $k \times (n - k)$ -rectangle. A *Deogram* (short for *Deodhar diagram*⁹) of shape λ is a filling D of the boxes of λ with crossings and elbows satisfying the following *distinguished condition* [Deo85]: for any elbow in D , the label of its bottom-left path is less than the label of its top-right path. In other words, once two paths have crossed an odd number of times, they cannot form an elbow. See Figure 4.

For example, any filling that consists either entirely of crossings or entirely of elbows satisfies the distinguished condition. Observe that when a Deogram D of shape λ consists entirely of crossings, the permutation $u_D = w$ indeed is k -Grassmannian: we have $w^{-1}(1) < w^{-1}(2) < \dots < w^{-1}(k)$ and $w^{-1}(k + 1) < \dots < w^{-1}(n)$. We denote this correspondence by $\lambda_w := \lambda$.

Definition 9.3. Let $f = f_{v,w} \in \mathbf{B}_{k,n}$. An f -Deogram is a Deogram D of shape λ_w satisfying $u_D = v$. A *maximal f -Deogram* is an f -Deogram with the maximal possible number of crossings among all f -Deograms.

We denote by Deo_f (resp., Deo_f^{\max}) the set of all (resp., maximal) f -Deograms.

Remark 9.4. It is easy to see that any f -Deogram must have at least $n - c(\bar{f})$ elbows. One can also check that for each $f \in \mathbf{B}_{k,n}$, there exist f -Deograms with exactly $n - c(\bar{f})$ elbows.¹⁰

Proposition 9.5. Let $f \in \mathbf{B}_{k,n}^{c=1}$. Then C_f equals the number of maximal f -Deograms:

$$(9.1) \quad C_f = \#\text{Deo}_f^{\max}.$$

Proof. This is a simple consequence of the results of Deodhar [Deo85]. Let v, w be such that $f = f_{v,w}$. By Proposition 4.3, we have $\#\Pi_f^{\circ}(\mathbb{F}_q) = \#R_{v,w}^{\circ}(\mathbb{F}_q)$. Deodhar expressed $\#R_{v,w}^{\circ}(\mathbb{F}_q)$ as a certain sum over *distinguished subexpressions* for v inside a reduced word $w = s_{i_1}s_{i_2}\dots s_{i_l}$, where $l = \ell(w)$. Here, a *subexpression* for v is a way to write v as a product $v_1v_2\dots v_l$, where $v_j \in \{s_{i_j}, \text{id}\}$ for all $j = 1, 2, \dots, l$. A subexpression is *distinguished* if for all j such that $\ell(v_1\dots v_{j-1}s_{i_j}) < \ell(v_1\dots v_{j-1})$, we have $v_j = s_{i_j}$. Since w is k -Grassmannian, the terms in the product $w = s_{i_1}s_{i_2}\dots s_{i_l}$ correspond to the boxes of λ_w . Each Deogram $D \in \text{Deo}_f$ gives rise to a distinguished subexpression for v , so that the indices j such that $v_j = s_{i_j}$ correspond to the

⁹The terminology *Deodhar diagram* is borrowed from [KW13].

¹⁰The same statement does not hold for Richardson varieties: for $w = s_1s_2s_3s_2s_1$ and $v = s_2$ in S_4 , there are no subexpressions for v inside w skipping exactly $n - c(wv^{-1}) = 2$ indices.




FIGURE 6. For $n = 2k + 1$, maximal $f_{k,n}$ -Deograms are in bijection with non-crossing alternating trees; see Remark 9.7.

crossings in D . It is easy to see that this correspondence is bijective. Thus, the results of [Deo85] imply that

$$(9.2) \quad \#\Pi_f^\circ(\mathbb{F}_q) = \sum_{D \in \text{Deo}_f} (q-1)^{\text{elb}(D)} q^{(\text{xing}(D) - \ell(w))/2},$$

where $\text{elb}(D)$ and $\text{xing}(D)$ denote the number of elbows and crossings in D . By Remark 9.4, each maximal f -Deogram contributes $(q-1)^{n-1} q^{(\ell_{v,w} - n + 1)/2}$ to the right-hand side of (9.2). (Note that $\text{xing}(D) + \text{elb}(D) = \ell(w)$ is constant.) It remains to note that $\#\mathcal{X}_f^\circ(\mathbb{F}_q)$ is obtained by dividing $\#\Pi_f^\circ(\mathbb{F}_q)$ by $\#T(\mathbb{F}_q) = (q-1)^{n-1}$, and that C_f is by definition the $q = 1$ specialization of $\#\mathcal{X}_f^\circ(\mathbb{F}_q)$. \square

Let us focus on the case $f = f_{k,n}$ with $\gcd(k, n) = 1$. Explicitly, a maximal $f_{k,n}$ -Deogram is a way of placing $n - 1$ elbows in a $k \times (n - k)$ rectangle and filling the rest with crossings so that (i) the resulting permutation obtained by following the paths is the identity, and (ii) the distinguished condition in Definition 9.2 is satisfied. By Proposition 9.5, the number $\#\text{Deo}_{f_{k,n}}^{\max}$ of such objects equals the number $\#\text{Dyck}_{k,(n-k)}$ of Dyck paths inside a $k \times (n - k)$ -rectangle.

Problem 9.6. Give a bijective proof of Proposition 9.5. That is, find a bijection between $\text{Deo}_{f_{k,n}}^{\max}$ and $\text{Dyck}_{k,(n-k)}$ for the case $\gcd(k, n) = 1$.

For instance, Figures 1 and 5 both have 7 objects in them, but it is unclear which objects correspond to which. It would be also interesting to understand the area and div statistics in the language of $f_{k,n}$ -Deograms.

Remark 9.7. For the case $n = 2k + 1$ of the standard Catalan numbers, the maximal $f_{k,n}$ -Deograms are easily seen to be in bijection with *non-crossing alternating trees on $k + 1$ vertices* (item 62 in [Sta15]). Explicitly, given $D \in \text{Deo}_{f_{k,n}}^{\max}$, we assign a vertex to each of the $k + 1$ columns of D . One can show that every row of D must contain exactly two elbows, and connecting the two corresponding vertices by an edge for each of the k rows, one obtains a non-crossing alternating tree. The case $k = 3$, $n = 7$ is illustrated in Figure 6.

Remark 9.8. A recursive proof of Proposition 9.5 for the case $n = dk \pm 1$ ($d \geq 2$) was found by David Speyer (private communication). It appears that when $n = dk \pm 1$, the distinguished condition is automatically satisfied for any maximal $f_{k,n}$ -Deogram. However, this is not the case for instance when $k = 5$ and $n = 12$; see Figure 4(right). We were able to find a recursive proof of (9.1) for arbitrary k, n . This and some other enumerative consequences of our results will appear in a separate paper [GL21].

Remark 9.9. A probabilistic interpretation of f -Deograms and their weights in (9.2) in terms of *the stochastic colored six-vertex model* [KMMO16] was recently discovered in [Gal20]. In particular, a result closely related to Theorem 2.3 appears in [Gal20, Lemma 7.1 and Proposition 7.3].

REFERENCES

- [AHBC⁺16] Nima Arkani-Hamed, Jacob Bourjaily, Freddy Cachazo, Alexander Goncharov, Alexander Postnikov, and Jaroslav Trnka. *Grassmannian Geometry of Scattering Amplitudes*. Cambridge University Press, Cambridge, 2016.
- [AHLS20] Nima Arkani-Hamed, Thomas Lam, and Marcus Spradlin. Positive configuration space. [arXiv:2003.03904v2](#), 2020.
- [AMRW19] Pramod N. Achar, Shotaro Makisumi, Simon Riche, and Geordie Williamson. Koszul duality for Kac-Moody groups and characters of tilting modules. *J. Amer. Math. Soc.*, 32(1):261–310, 2019.
- [AR13] Pramod N. Achar and Simon Riche. Koszul duality and semisimplicity of Frobenius. *Ann. Inst. Fourier (Grenoble)*, 63(4):1511–1612, 2013.
- [AR16a] Pramod N. Achar and Simon Riche. Modular perverse sheaves on flag varieties I: tilting and parity sheaves. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(2):325–370, 2016. With a joint appendix with Geordie Williamson.
- [AR16b] Pramod N. Achar and Simon Riche. Modular perverse sheaves on flag varieties, II: Koszul duality and formality. *Duke Math. J.*, 165(1):161–215, 2016.
- [BB81] Alexandre Beilinson and Joseph Bernstein. Localisation de g -modules. *C. R. Acad. Sci. Paris Sér. I Math.*, 292(1):15–18, 1981.
- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne. Faisceaux pervers. In *Analysis and topology on singular spaces, I (Luminy, 1981)*, volume 100 of *Astérisque*, pages 5–171. Soc. Math. France, Paris, 1982.
- [Bea99] Arnaud Beauville. Counting rational curves on $K3$ surfaces. *Duke Math. J.*, 97(1):99–108, 1999.
- [Bei87] A. A. Beilinson. On the derived category of perverse sheaves. In *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 27–41. Springer, Berlin, 1987.
- [BGS96] Alexander Beilinson, Victor Ginzburg, and Wolfgang Soergel. Koszul duality patterns in representation theory. *J. Amer. Math. Soc.*, 9(2):473–527, 1996.
- [BGY06] K. A. Brown, K. R. Goodearl, and M. Yakimov. Poisson structures on affine spaces and flag varieties. I. Matrix affine Poisson space. *Adv. Math.*, 206(2):567–629, 2006.
- [BK81] J.-L. Brylinski and M. Kashiwara. Kazhdan-Lusztig conjecture and holonomic systems. *Invent. Math.*, 64(3):387–410, 1981.
- [BL94] Joseph Bernstein and Valery Lunts. *Equivariant sheaves and functors*, volume 1578 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [BT20] Roman Bezrukavnikov and Kostiantyn Tolmachov. Monodromic model for Khovanov-Rozansky homology. [arXiv:2008.11379v1](#), 2020.
- [BY13] Roman Bezrukavnikov and Zhiwei Yun. On Koszul duality for Kac-Moody groups. *Represent. Theory*, 17:1–98, 2013.
- [BZ97] Arkady Berenstein and Andrei Zelevinsky. Total positivity in Schubert varieties. *Comment. Math. Helv.*, 72(1):128–166, 1997.
- [CD16] Ivan Cherednik and Ivan Danilenko. DAHA and iterated torus knots. *Algebr. Geom. Topol.*, 16(2):843–898, 2016.
- [CGGS20] Roger Casals, Eugene Gorsky, Mikhail Gorsky, and José Simental. Algebraic Weaves and Braid Varieties. [arXiv:2012.06931v1](#), 2020.
- [CGGS21] Roger Casals, Eugene Gorsky, Mikhail Gorsky, and José Simental. Positroid Links and Braid varieties. [arXiv:2105.13948v1](#), 2021.
- [Che13] Ivan Cherednik. Jones polynomials of torus knots via DAHA. *Int. Math. Res. Not. IMRN*, (23):5366–5425, 2013.
- [Che20] Ivan Cherednik. Gröbner cells of punctual Hilbert schemes in dimension two. [arXiv:2005.05759v1](#), 2020.
- [CM18] Erik Carlsson and Anton Mellit. A proof of the shuffle conjecture. *J. Amer. Math. Soc.*, 31(3):661–697, 2018.
- [CP18] Ivan Cherednik and Ian Philipp. DAHA and plane curve singularities. *Algebr. Geom. Topol.*, 18(1):333–385, 2018.
- [CP19] Ivan Cherednik and Ian Philipp. Modules over plane curve singularities in any ranks and DAHA. *J. Algebra*, 520:186–236, 2019.
- [CR64] L. Carlitz and J. Riordan. Two element lattice permutation numbers and their q -generalization. *Duke Math. J.*, 31:371–388, 1964.
- [dCHM12] Mark Andrea A. de Cataldo, Tamás Hausel, and Luca Migliorini. Topology of Hitchin systems and Hodge theory of character varieties: the case A_1 . *Ann. of Math. (2)*, 175(3):1329–1407, 2012.
- [dCM04] Mark Andrea A. de Cataldo and Luca Migliorini. The Gysin map is compatible with mixed Hodge structures. In *Algebraic structures and moduli spaces*, volume 38 of *CRM Proc. Lecture Notes*, pages 133–138. Amer. Math. Soc., Providence, RI, 2004.

- [Del71] Pierre Deligne. Théorie de Hodge. II. *Inst. Hautes Études Sci. Publ. Math.*, (40):5–57, 1971.
- [Del80] Pierre Deligne. La conjecture de Weil. II. *Inst. Hautes Études Sci. Publ. Math.*, (52):137–252, 1980.
- [Deo85] Vinay V. Deodhar. On some geometric aspects of Bruhat orderings. I. A finer decomposition of Bruhat cells. *Invent. Math.*, 79(3):499–511, 1985.
- [DGR06] Nathan M. Dunfield, Sergei Gukov, and Jacob Rasmussen. The superpolynomial for knot homologies. *Experiment. Math.*, 15(2):129–159, 2006.
- [DM17] Gurbir Dhillon and Visu Makam. Extensions between Verma modules for dihedral groups. [arXiv:1712.05299v1](https://arxiv.org/abs/1712.05299), 2017.
- [EH19] Ben Elias and Matthew Hogancamp. On the computation of torus link homology. *Compos. Math.*, 155(1):164–205, 2019.
- [EMTW20] Ben Elias, Shotaro Makisumi, Ulrich Thiel, and Geordie Williamson. *Introduction to Soergel Bimodules*, volume 5 of *RSME Springer Series*. Springer International Publishing, 2020.
- [EW16] Ben Elias and Geordie Williamson. Soergel calculus. *Represent. Theory*, 20:295–374, 2016.
- [FPST17] Sergey Fomin, Pavlo Pylyavskyy, Eugenii Shustin, and Dylan Thurston. Morsifications and mutations. [arXiv:1711.10598v3](https://arxiv.org/abs/1711.10598), 2017.
- [FS00] Sergey Fomin and Michael Shapiro. Stratified spaces formed by totally positive varieties. *Michigan Math. J.*, 48:253–270, 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [FW87] John Franks and R. F. Williams. Braids and the Jones polynomial. *Trans. Amer. Math. Soc.*, 303(1):97–108, 1987.
- [FYH⁺85] P. Freyd, D. Yetter, J. Hoste, W. B. R. Lickorish, K. Millett, and A. Ocneanu. A new polynomial invariant of knots and links. *Bull. Amer. Math. Soc. (N.S.)*, 12(2):239–246, 1985.
- [FZ99] Sergey Fomin and Andrei Zelevinsky. Double Bruhat cells and total positivity. *J. Amer. Math. Soc.*, 12(2):335–380, 1999.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [Gal20] Pavel Galashin. Symmetries of stochastic colored vertex models. *Ann. Probab.*, to appear. [arXiv:2003.06330v1](https://arxiv.org/abs/2003.06330), 2020.
- [GH96] A. M. Garsia and M. Haiman. A remarkable q, t -Catalan sequence and q -Lagrange inversion. *J. Algebraic Combin.*, 5(3):191–244, 1996.
- [GH02] A. M. Garsia and J. Haglund. A proof of the q, t -Catalan positivity conjecture. volume 256, pages 677–717. 2002. LaCIM 2000 Conference on Combinatorics, Computer Science and Applications (Montreal, QC).
- [GH22] Eugene Gorsky and Matthew Hogancamp. Hilbert schemes and y -ification of Khovanov-Rozansky homology. *Geom. Topol.*, 26(2):587–678, 2022.
- [GHM21] Eugene Gorsky, Matthew Hogancamp, and Anton Mellit. Tautological classes and symmetry in Khovanov-Rozansky homology. [arXiv:2103.01212v1](https://arxiv.org/abs/2103.01212), 2021.
- [GKL19] Pavel Galashin, Steven N. Karp, and Thomas Lam. Regularity theorem for totally nonnegative flag varieties. *J. Amer. Math. Soc.*, to appear. [arXiv:1904.00527v2](https://arxiv.org/abs/1904.00527), 2019.
- [GKS21] Eugene Gorsky, Oscar Kivinen, and José Simental. Algebra and geometry of link homology. [arXiv:2108.10356v1](https://arxiv.org/abs/2108.10356), 2021.
- [GL19] Pavel Galashin and Thomas Lam. Positroid varieties and cluster algebras. *Ann. Sci. Éc. Norm. Supér.*, to appear. [arXiv:1906.03501v1](https://arxiv.org/abs/1906.03501), 2019.
- [GL20] Pavel Galashin and Thomas Lam. Parity duality for the amplituhedron. *Compos. Math.*, 156(11):2207–2262, 2020.
- [GL21] Pavel Galashin and Thomas Lam. Positroid Catalan numbers. [arXiv:2104.05701v1](https://arxiv.org/abs/2104.05701), 2021.
- [GL22] Pavel Galashin and Thomas Lam. Plabic links, quivers, and skein relations. [arXiv:2208.01175v1](https://arxiv.org/abs/2208.01175), 2022.
- [GM13] Evgeny Gorsky and Mikhail Mazin. Compactified Jacobians and q, t -Catalan numbers, I. *J. Combin. Theory Ser. A*, 120(1):49–63, 2013.
- [GM14] Evgeny Gorsky and Mikhail Mazin. Compactified Jacobians and q, t -Catalan numbers, II. *J. Algebraic Combin.*, 39(1):153–186, 2014.
- [GMM14] J. González-Meneses and P. M. G. Manchón. Closures of positive braids and the Morton-Franks-Williams inequality. *Topology Appl.*, 174:14–24, 2014.
- [GN15] Eugene Gorsky and Andrei Neguț. Refined knot invariants and Hilbert schemes. *J. Math. Pures Appl. (9)*, 104(3):403–435, 2015.
- [Gom06] Yasushi Gomi. The markov traces and the fourier transforms. *Journal of Algebra*, 303(2):566–591, 2006. Computational Algebra.
- [Gor12] E. Gorsky. q, t -Catalan numbers and knot homology. In *Zeta functions in algebra and geometry*, volume 566 of *Contemp. Math.*, pages 213–232. Amer. Math. Soc., Providence, RI, 2012.

- [GORS14] Eugene Gorsky, Alexei Oblomkov, Jacob Rasmussen, and Vivek Shende. Torus knots and the rational DAHA. *Duke Math. J.*, 163(14):2709–2794, 2014.
- [GSV10] Michael Gekhtman, Michael Shapiro, and Alek Vainshtein. *Cluster algebras and Poisson geometry*, volume 167 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [Hag03] J. Haglund. Conjectured statistics for the q, t -Catalan numbers. *Adv. Math.*, 175(2):319–334, 2003.
- [Hag08] James Haglund. *The q, t -Catalan numbers and the space of diagonal harmonics*, volume 41 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2008. With an appendix on the combinatorics of Macdonald polynomials.
- [Hag16] Jim Haglund. The combinatorics of knot invariants arising from the study of Macdonald polynomials. In *Recent trends in combinatorics*, volume 159 of *IMA Vol. Math. Appl.*, pages 579–600. Springer, [Cham], 2016.
- [Hai94] Mark D. Haiman. Conjectures on the quotient ring by diagonal invariants. *J. Algebraic Combin.*, 3(1):17–76, 1994.
- [Hai02] Mark Haiman. Vanishing theorems and character formulas for the Hilbert scheme of points in the plane. *Invent. Math.*, 149(2):371–407, 2002.
- [HM19] Matthew Hogancamp and Anton Mellit. Torus link homology. [arXiv:1909.00418v1](https://arxiv.org/abs/1909.00418), 2019.
- [Hog17] Matthew Hogancamp. Khovanov-Rozansky homology and higher Catalan sequences. [arXiv:1704.01562v1](https://arxiv.org/abs/1704.01562), 2017.
- [HRV08] Tamás Hausel and Fernando Rodriguez-Villegas. Mixed Hodge polynomials of character varieties. *Invent. Math.*, 174(3):555–624, 2008. With an appendix by Nicholas M. Katz.
- [Hum08] James E. Humphreys. *Representations of semisimple Lie algebras in the BGG category \mathcal{O}* , volume 94 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [Jon87] V. F. R. Jones. Hecke algebra representations of braid groups and link polynomials. *Ann. of Math. (2)*, 126(2):335–388, 1987.
- [Kho07] Mikhail Khovanov. Triply-graded link homology and Hochschild homology of Soergel bimodules. *Internat. J. Math.*, 18(8):869–885, 2007.
- [KL79] David Kazhdan and George Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53(2):165–184, 1979.
- [KL80] David Kazhdan and George Lusztig. Schubert varieties and Poincaré duality. In *Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, Proc. Sympos. Pure Math., XXXVI, pages 185–203. Amer. Math. Soc., Providence, R.I., 1980.
- [KLS13] Allen Knutson, Thomas Lam, and David E. Speyer. Positroid varieties: juggling and geometry. *Compos. Math.*, 149(10):1710–1752, 2013.
- [KMMO16] A. Kuniba, V. V. Mangazeev, S. Maruyama, and M. Okado. Stochastic R matrix for $U_q(A_n^{(1)})$. *Nuclear Phys. B*, 913:248–277, 2016.
- [KR08a] Mikhail Khovanov and Lev Rozansky. Matrix factorizations and link homology. *Fund. Math.*, 199(1):1–91, 2008.
- [KR08b] Mikhail Khovanov and Lev Rozansky. Matrix factorizations and link homology. II. *Geom. Topol.*, 12(3):1387–1425, 2008.
- [KW13] Yuji Kodama and Lauren Williams. The Deodhar decomposition of the Grassmannian and the regularity of KP solitons. *Adv. Math.*, 244:979–1032, 2013.
- [KWY13] Allen Knutson, Alexander Woo, and Alexander Yong. Singularities of Richardson varieties. *Math. Res. Lett.*, 20(2):391–400, 2013.
- [Lam16] Thomas Lam. Totally nonnegative Grassmannian and Grassmann polytopes. In *Current developments in mathematics 2014*, pages 51–152. Int. Press, Somerville, MA, 2016.
- [Lec16] B. Leclerc. Cluster structures on strata of flag varieties. *Adv. Math.*, 300:190–228, 2016.
- [LO08] Yves Laszlo and Martin Olsson. The six operations for sheaves on Artin stacks. II. Adic coefficients. *Publ. Math. Inst. Hautes Études Sci.*, (107):169–210, 2008.
- [LP20] Thomas Lam and Alex Postnikov. Polypositroids. [arXiv:2010.07120v1](https://arxiv.org/abs/2010.07120), 2020.
- [LS91] G. Lusztig and J. M. Smelt. Fixed point varieties on the space of lattices. *Bull. London Math. Soc.*, 23(3):213–218, 1991.
- [LS16] Thomas Lam and David E. Speyer. Cohomology of cluster varieties. I. Locally acyclic case. [arXiv:1604.06843v1](https://arxiv.org/abs/1604.06843), 2016.
- [LW09] Nicholas A. Loehr and Gregory S. Warrington. A continuous family of partition statistics equidistributed with length. *J. Combin. Theory Ser. A*, 116(2):379–403, 2009.
- [Mac60] Percy A. MacMahon. *Combinatory analysis*. Two volumes (bound as one). Chelsea Publishing Co., New York, 1960.

- [Maz07] Volodymyr Mazorchuk. Some homological properties of the category \mathcal{O} . *Pacific J. Math.*, 232(2):313–341, 2007.
- [Mel16] Anton Mellit. Toric braids and (m, n) -parking functions. [arXiv:1604.07456v1](https://arxiv.org/abs/1604.07456), 2016.
- [Mel17] Anton Mellit. Homology of torus knots. [arXiv:1704.07630v1](https://arxiv.org/abs/1704.07630), 2017.
- [Mel19] Anton Mellit. Cell decompositions of character varieties. [arXiv:1905.10685v1](https://arxiv.org/abs/1905.10685), 2019.
- [Mil13] James S. Milne. Lectures on etale cohomology (v2.21), 2013. Available at <http://www.jmilne.org/math/>.
- [Mor86] H. R. Morton. Seifert circles and knot polynomials. *Math. Proc. Cambridge Philos. Soc.*, 99(1):107–109, 1986.
- [MS13] Luca Migliorini and Vivek Shende. A support theorem for Hilbert schemes of planar curves. *J. Eur. Math. Soc. (JEMS)*, 15(6):2353–2367, 2013.
- [MS17] Greg Muller and David E. Speyer. The twist for positroid varieties. *Proc. Lond. Math. Soc. (3)*, 115(5):1014–1071, 2017.
- [MY14] Davesh Maulik and Zhiwei Yun. Macdonald formula for curves with planar singularities. *J. Reine Angew. Math.*, 694:27–48, 2014.
- [O’H90] Kathleen M. O’Hara. Unimodality of Gaussian coefficients: a constructive proof. *J. Combin. Theory Ser. A*, 53(1):29–52, 1990.
- [OR20] Alexei Oblomkov and Lev Rozansky. Soergel bimodules and matrix factorizations. [arXiv:2010.14546v1](https://arxiv.org/abs/2010.14546), 2020.
- [ORS18] Alexei Oblomkov, Jacob Rasmussen, and Vivek Shende. The Hilbert scheme of a plane curve singularity and the HOMFLY homology of its link. *Geom. Topol.*, 22(2):645–691, 2018. With an appendix by Eugene Gorsky.
- [Pio07] Jens Piontowski. Topology of the compactified Jacobians of singular curves. *Math. Z.*, 255(1):195–226, 2007.
- [Pos06] Alexander Postnikov. Total positivity, Grassmannians, and networks. Preprint, <http://math.mit.edu/~apost/papers/tpgrass.pdf>, 2006.
- [PS08] Chris A. M. Peters and Joseph H. M. Steenbrink. *Mixed Hodge structures*, volume 52 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2008.
- [PT87] Józef H. Przytycki and Paweł Traczyk. Conway algebras and skein equivalence of links. *Proc. Amer. Math. Soc.*, 100(4):744–748, 1987.
- [Ras07] Jacob Rasmussen. Khovanov-Rozansky homology of two-bridge knots and links. *Duke Math. J.*, 136(3):551–583, 2007.
- [Rid13] Laura Rider. Formality for the nilpotent cone and a derived Springer correspondence. *Adv. Math.*, 235:208–236, 2013.
- [Rol90] Dale Rolfsen. *Knots and links*, volume 7 of *Mathematics Lecture Series*. Publish or Perish, Inc., Houston, TX, 1990. Corrected reprint of the 1976 original.
- [Rou04] Raphaël Rouquier. Categorification of the braid groups. [arXiv:math/0409593v1](https://arxiv.org/abs/math/0409593), 2004.
- [Rou17] Raphaël Rouquier. Khovanov-Rozansky homology and 2-braid groups. In *Categorification in geometry, topology, and physics*, volume 684 of *Contemp. Math.*, pages 147–157. Amer. Math. Soc., Providence, RI, 2017.
- [RSW14] Simon Riche, Wolfgang Soergel, and Geordie Williamson. Modular Koszul duality. *Compos. Math.*, 150(2):273–332, 2014.
- [Sco06] J. S. Scott. Grassmannians and cluster algebras. *Proc. Lond. Math. Soc. (3)*, 92(2):345–380, 2006.
- [Sha93] B. Z. Shapiro. The mixed Hodge structure of the complement to an arbitrary arrangement of affine complex hyperplanes is pure. *Proc. Amer. Math. Soc.*, 117(4):931–933, 1993.
- [Soe89] W. Soergel. n -cohomology of simple highest weight modules on walls and purity. *Invent. Math.*, 98(3):565–580, 1989.
- [Soe90] Wolfgang Soergel. Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. *J. Amer. Math. Soc.*, 3(2):421–445, 1990.
- [Soe92] Wolfgang Soergel. The combinatorics of Harish-Chandra bimodules. *J. Reine Angew. Math.*, 429:49–74, 1992.
- [Soe07] Wolfgang Soergel. Kazhdan-Lusztig-Polynome und unzerlegbare Bimoduln über Polynomringen. *J. Inst. Math. Jussieu*, 6(3):501–525, 2007.
- [SSBW19] K. Serhiyenko, M. Sherman-Bennett, and L. Williams. Cluster structures in Schubert varieties in the Grassmannian. *Proc. Lond. Math. Soc. (3)*, 119(6):1694–1744, 2019.

- [Sta84] Richard P. Stanley. Combinatorial applications of the hard Lefschetz theorem. In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983)*, pages 447–453. PWN, Warsaw, 1984.
- [Sta12] Richard P. Stanley. *Enumerative combinatorics. Volume 1*, volume 49 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2012.
- [Sta15] Richard P. Stanley. *Catalan numbers*. Cambridge University Press, New York, 2015.
- [Str03] Catharina Stroppel. Category \mathcal{O} : gradings and translation functors. *J. Algebra*, 268(1):301–326, 2003.
- [Stu18] Eric Nathan Stucky. Parity-Unimodality and a Cyclic Sieving Phenomenon for Necklaces. [arXiv:1812.04578v3](https://arxiv.org/abs/1812.04578v3), 2018.
- [STWZ19] Vivek Shende, David Treumann, Harold Williams, and Eric Zaslow. Cluster varieties from Legendrian knots. *Duke Math. J.*, 168(15):2801–2871, 2019.
- [STZ17] Vivek Shende, David Treumann, and Eric Zaslow. Legendrian knots and constructible sheaves. *Invent. Math.*, 207(3):1031–1133, 2017.
- [SVW18] Wolfgang Soergel, Rahbar Virk, and Matthias Wendt. Equivariant motives and geometric representation theory. (with an appendix by F. Hörmann and M. Wendt). [arXiv:1809.05480v1](https://arxiv.org/abs/1809.05480v1), 2018.
- [Syl78] J. J. Sylvester. Proof of the hitherto undemonstrated fundamental theorem of invariants. *Philos. Mag.*, 5:178–188, 1878.
- [WW] Ben Webster and Geordie Williamson. The bounded below equivariant derived category. <http://mathserver.neu.edu/~bwebster/equivariant.pdf>.
- [WW08] Ben Webster and Geordie Williamson. A geometric model for Hochschild homology of Soergel bimodules. *Geom. Topol.*, 12(2):1243–1263, 2008.
- [WW11] Ben Webster and Geordie Williamson. The geometry of Markov traces. *Duke Math. J.*, 160(2):401–419, 2011.
- [WW17] Ben Webster and Geordie Williamson. A geometric construction of colored HOMFLYPT homology. *Geom. Topol.*, 21(5):2557–2600, 2017.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, 520 PORTOLA PLAZA, LOS ANGELES, CA 90025, USA

Email address: galashin@math.ucla.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 2074 EAST HALL, 530 CHURCH STREET, ANN ARBOR, MI 48109-1043, USA

Email address: tfylam@umich.edu