

Massey Products and Ideal Class Groups

Romyar T. Sharifi

July 26, 2006

Abstract

We consider certain Massey products in the cohomology of a Galois extension of fields with coefficients in p -power roots of unity. We prove formulas for these products both in general and in the special case that the Galois extension in question is the maximal extension of a number field unramified outside a set of primes S including those above p and any archimedean places. We then consider those \mathbf{Z}_p -Kummer extensions L_∞ of the maximal p -cyclotomic extension K_∞ of a number field K that are unramified outside S . We show that Massey products describe the structure of a certain “decomposition-free” quotient of a graded piece of the maximal unramified abelian pro- p extension of L_∞ in which all primes above those in S split completely, with the grading arising from the augmentation filtration on the group ring of the Galois group of L_∞/K_∞ . We explicitly describe examples of the maximal unramified abelian pro- p extensions of unramified outside p Kummer extensions of the cyclotomic field of all p -power roots of unity, for irregular primes p .

1 Introduction

1.1 Iwasawa modules over Kummer extensions

Classical Iwasawa theory takes place over the cyclotomic \mathbf{Z}_p -extension K_∞ of a number field K , for a prime p . One considers finitely generated modules for the Iwasawa algebra $\Lambda = \mathbf{Z}_p[[\text{Gal}(K_\infty/K)]]$, known as Iwasawa modules. This paper is concerned with Iwasawa theory over a \mathbf{Z}_p -extension L_∞ of K_∞ that is Galois over K . The pro- p group $\mathcal{G} = \text{Gal}(L_\infty/K)$ is in general nonabelian, being isomorphic to a semi-direct product of two copies of \mathbf{Z}_p : its normal subgroup $G = \text{Gal}(L_\infty/K_\infty)$ and any lift of its quotient $\Gamma = \text{Gal}(K_\infty/K)$. Whereas Λ is isomorphic to a power series ring in one variable over \mathbf{Z}_p , the Iwasawa algebra $\mathbf{Z}_p[[\mathcal{G}]]$ is isomorphic to a ring of \mathbf{Z}_p -power series

in two variables that do not commute if \mathcal{G} is nonabelian. The structure theory of finitely generated $\mathbf{Z}_p[[\mathcal{G}]]$ -modules is therefore considerably more complicated than in the classical case (see [V2] for a study).

Our idea is a simple one. Given a finitely generated $\mathbf{Z}_p[[\mathcal{G}]]$ -module M , it is possible to create an infinite sequence of Λ -modules, each a quotient of the last, as follows. Let I_G denote the augmentation ideal in $\mathbf{Z}_p[[G]]$, and consider the quotients

$$\mathrm{gr}_G^k M = I_G^k M / I_G^{k+1} M$$

for $k \geq 0$. The graded object

$$\mathrm{gr}_G M = \bigoplus_{k=0}^{\infty} \mathrm{gr}_G^k M$$

is a module over the graded ring $\mathrm{gr}_G \mathbf{Z}_p[[\mathcal{G}]]$. In particular, each $\mathrm{gr}_G^k M$ is a finitely generated $\Lambda \cong \mathrm{gr}_G^0 \mathbf{Z}_p[[\mathcal{G}]]$ -module. Since

$$\mathrm{gr}_G^k \mathbf{Z}_p[[G]] = I_G^k / I_G^{k+1} \cong G^{\otimes k},$$

the action of its first graded piece provides a surjection

$$\mathrm{gr}_G^k M \otimes_{\mathbf{Z}_p} G \twoheadrightarrow \mathrm{gr}_G^{k+1} M$$

of Λ -modules. (Note that G is isomorphic as a Λ -module to a twist of \mathbf{Z}_p by a possibly nonintegral power of the cyclotomic character.) Of course, a good deal of information on M can be lost in the passage to the graded module, but much is retained. For instance, we can detect the torsionness of M as a $\mathbf{Z}_p[[G]]$ -module from the finiteness of the $\mathrm{gr}_G^k M$ for large k .

Now, one of the most typical Iwasawa modules to consider is the Galois group \mathcal{Z}_K of the maximal unramified abelian pro- p extension of K_∞ . The action of Γ on \mathcal{Z}_K is given by lifting and conjugating. By class field theory, \mathcal{Z}_K can also be thought of as the inverse limit of p -parts of class groups of intermediate subfields of K , and the action is induced by the usual action of Γ on ideals. By a standard result of Iwasawa, \mathcal{Z}_K is a finitely generated, torsion Λ -module.

We wish to consider \mathcal{Z}_L , the Galois group of the maximal unramified abelian pro- p extension of L_∞ , and its relation to \mathcal{Z}_K . Let us assume that L_∞/K is unramified outside a finite set of primes of K . We remark that $\mathcal{Z}_L/I_G \mathcal{Z}_L$ is closely related to \mathcal{Z}_K , mapping to it with kernel and cokernel finitely generated over \mathbf{Z}_p . As a consequence,

\mathcal{Z}_L is finitely generated over $\mathbf{Z}_p[[\mathcal{G}]]$. Assume now that K_∞ contains all p -power roots of 1. We show in this paper that the higher graded quotients $\mathrm{gr}_G^k \mathcal{Z}_L$ for positive k relate to quotients of \mathcal{Z}_K by the image of certain operations in Galois cohomology called Massey products. To be precise in our description, we must replace \mathcal{Z}_K and \mathcal{Z}_L with closely related Iwasawa modules.

We may consider the quotient of \mathcal{Z}_K that is the Galois group \mathcal{A}_K of the maximal unramified abelian pro- p extension of K_∞ in which all primes above those in p split completely. We also have the analogously defined Galois group \mathcal{A}_L over L_∞ . Again by class field theory, the modules \mathcal{A}_K and \mathcal{A}_L are isomorphic to the inverse limits of p -parts of the quotients of class groups by the classes of primes above p in the intermediate subfields of K_∞ and L_∞ , respectively. It is the higher graded quotients of \mathcal{A}_L that relate to Massey products, as we shall describe below. We now state our main result (Theorem 6.3), with a strong extra restriction for simplicity of presentation.

Theorem A. *Assume that there is a unique prime above p in L_∞ and that L_∞/K_∞ is unramified outside of it. For each $k \geq 1$, there is a canonical isomorphism of Λ -modules,*

$$\mathrm{gr}_G^k \mathcal{A}_L \cong (\mathcal{A}_K / \mathcal{P}_{L/K}^{(k)}) \otimes_{\mathbf{Z}_p} G^{\otimes k},$$

where $\mathcal{P}_{L/K}^{(k)}$ is a group of inverse limits of $(k+1)$ -fold Massey products with k -copies of a Kummer generator of L_∞ .

Some applications of this result are given in Section 7 in the case that $K = \mathbf{Q}(\mu_p)$ with p odd. When L_∞ is defined by a sequence of cyclotomic p -units in $K_\infty = \mathbf{Q}(\mu_{p^\infty})$, we show that $\mathrm{gr}_G^1 \mathcal{A}_L$ may be either zero or nonzero by providing examples of both instances when $p = 37$ (Propositions 7.3 and 7.4). For such L_∞ , we conjecture that \mathcal{A}_L is finitely generated as a $\mathbf{Z}_p[[G]]$ -module (Conjecture 7.6). When L_∞ is a CM-field, however, we show that \mathcal{A}_L is not always finitely generated over $\mathbf{Z}_p[[G]]$ (Proposition 7.5). We have good reason to believe that the Massey products of cyclotomic p -units will provide highly interesting arithmetic objects with an interpretation in terms of p -adic L -functions of modular forms, but we leave that for future endeavors.

1.2 Construction of Massey products

In order to describe the Massey products of interest, it is easiest if we begin by working in a general context. For further technical details on the constructions we give in this

subsection, see Sections 2-4. Let H be a group, and let $N \geq 1$. Massey products are products of two or more elements of $H^1(H, \mathbf{Z}/N\mathbf{Z})$ that take values in a quotient of $H^2(H, \mathbf{Z}/N\mathbf{Z})$.

The simplest form of Massey product is the cup product. Let χ_1, χ_2 be two characters in $H^1(H, \mathbf{Z}/N\mathbf{Z})$. Recall that the cup product $\chi_1 \cup \chi_2 \in H^2(H, \mathbf{Z}/N\mathbf{Z})$ is represented by the two-cocycle $\nu: (h_1, h_2) \mapsto \chi_1(h_1)\chi_2(h_2)$ for $h_1, h_2 \in H$. The cup product provides the obstruction to the existence of a homomorphism $\rho: H \rightarrow GL_3(\mathbf{Z}/N\mathbf{Z})$ with

$$\rho(h) = \begin{pmatrix} 1 & \chi_1(h) & \kappa(h) \\ 0 & 1 & \chi_2(h) \\ 0 & 0 & 1 \end{pmatrix}$$

for some function $\kappa: H \rightarrow \mathbf{Z}/N\mathbf{Z}$. Put precisely, the two-coboundary of κ is $-\nu$, so the existence of ρ is equivalent to ν being a coboundary. Note that there is some ambiguity in the choice of κ , when it exists, being well-defined only up to a character in $H^1(H, \mathbf{Z}/N\mathbf{Z})$.

In general, we can define Massey products recursively. Let $\chi_1, \chi_2, \dots, \chi_q$ be characters in $H^1(H, \mathbf{Z}/N\mathbf{Z})$. Suppose we have a map $\rho: H \rightarrow GL_{q+1}(\mathbf{Z}/N\mathbf{Z})$ into the group of unipotent upper-triangular matrices T_{q+1} , with

$$\rho(h) = \begin{pmatrix} 1 & \chi_1(h) & \kappa_{1,3}(h) & \dots & \kappa_{1,q+1}(h) \\ & 1 & \chi_2(h) & \dots & \kappa_{2,q+1}(h) \\ & & \ddots & \ddots & \vdots \\ & & & 1 & \chi_q(h) \\ & & & & 1 \end{pmatrix} \quad (1.1)$$

and such that “modulo the upper right-hand corner,” ρ is a homomorphism. That is to say, letting Z_{q+1} denote the subgroup of T_{q+1} consisting of matrices with zeros everywhere off of the diagonal except the upper right-hand corner, we require that the composition $\bar{\rho}$ of ρ with the surjection $T_{q+1} \rightarrow T_{q+1}/Z_{q+1}$ is a homomorphism. Then

$$\nu: (h, h') \mapsto \chi_1(h)\kappa_{2,q+1}(h') + \kappa_{1,3}(h)\kappa_{3,q+1}(h') + \dots + \kappa_{1,q}(h)\chi_q(h'), \quad (1.2)$$

is a 2-cocycle. If ρ is a homomorphism, then ν is the negative of the coboundary of $\kappa_{1,q+1}$. Conversely, if ν has trivial class, then there exists a lift of $\bar{\rho}$ to a homomorphism to T_{q+1} .

Ideally, the cohomology class of the 2-cocycle ν of (1.2) would represent the cohomology class of the q -fold Massey product of the χ_1, \dots, χ_q . However, the ambiguity in

the choice of the $\kappa_{i,j}$ defining $\bar{\rho}$ means that this class is not well-defined. For instance, in the case that $q = 3$, we could modify $\kappa_{1,3}$ by a cocycle ψ , and this will modify ν by the 2-cocycle representing $\psi \cup \chi_3$. Similarly, changing $\kappa_{2,4}$ by a cocycle modifies ν by the cup product of χ_1 with a cocycle. The Massey triple product is therefore taken to be the class of ν in the quotient

$$H^2(H, \mathbf{Z}/N\mathbf{Z}) / (H^1(H, \mathbf{Z}/N\mathbf{Z}) \cup \chi_3 + \chi_1 \cup H^1(H, \mathbf{Z}/N\mathbf{Z})).$$

A similar story holds for higher-fold Massey products. In general, the choice of the maps defining $\bar{\rho}$ (the $\kappa_{i,j}$ with $1 \leq i < j - 1 \leq q$ and $(i, j) \neq (1, q + 1)$) is called a *defining system* of the Massey product of χ_1, \dots, χ_q . The indeterminacy in the Massey product arises from the choice of defining system.

In this paper, we specialize to the case of q -fold Massey products in which all the χ_i are equal, except possibly χ_q . Let $N = p^m$ for some $m \geq 1$. Assume that we are given a fixed surjective homomorphism $\chi: H \rightarrow \mathbf{Z}/p^n\mathbf{Z}$ for some n with $q \leq p^{n-m+1}$, and let χ_i be the reduction of χ modulo p^m for $i \leq q - 1$. Consider the subgroup C_{q+1} of T_{q+1} consisting of unipotent matrices with zeros outside the rightmost column. Our assumptions imply that there exist canonical choices of the maps $\kappa_{i,j}$ for $j \leq q$ such that ρ composed with $T_{q+1} \rightarrow T_{q+1}/C_{q+1}$ is a homomorphism with cyclic image. In fact, we take

$$\kappa_{i,j}(h) \equiv \begin{pmatrix} \tilde{\chi}(h) \\ j - i \end{pmatrix} \pmod{p^m}$$

for $h \in H$, where $\tilde{\chi}: H \rightarrow \mathbf{Z}$ is the lift of χ with image in $[0, p^n)$. Allowing only these canonical maps in our defining systems, the ambiguity in the definition of the Massey product of χ_1, \dots, χ_q is reduced to the choice of the $\kappa_{i,q+1}$ with $2 \leq i \leq q - 1$.

We now pass to the setting of Galois cohomology so that we may interpret our cohomology groups as groups of field elements (modulo powers) via Kummer theory. Suppose that H is the Galois group of some Galois extension of fields Ω/F , with F of prime-to- p characteristic containing the p^n th roots of unity μ_{p^n} . Then χ factors through the Galois group Q of a Kummer extension E/F of degree p^n . The group C_{q+1} is isomorphic to $(\mathbf{Z}/p^m\mathbf{Z})[Q]/I_Q^q$ as a $(\mathbf{Z}/p^m\mathbf{Z})[Q]$ -module via conjugation by a lift to T_{q+1} of the image of Q in T_{q+1}/C_{q+1} . We identify μ_{p^n} with $\mathbf{Z}/p^n\mathbf{Z}$ by a choice of primitive p^n th root of unity ζ_n . The Kummer dual $\text{Hom}(C_{q+1}, \mu_{p^n})$ of C_{q+1} is then isomorphic to a submodule of $(\mathbf{Z}/p^m\mathbf{Z})[Q]$. Its generators as a group, labelled $D^{(i)}$ for nonnegative $i \leq q - 1$, satisfy

$$(\sigma - 1)D^{(i)} = D^{(i-1)},$$

with $D^{(0)}$ the norm element and σ the generator of Q which χ takes to 1. These “derivative operators” are studied in Section 3.

Note that χ (resp., χ_q) corresponds to some $a \in F^\times - F^{\times p}$ (resp., $b \in F^\times$) that becomes a p^n th power (resp., p^m th power) in Ω by Kummer theory. By the same principle, if we can realize b as a norm of an element $y \in E^\times$, then the $D^{(i)}y$ with $i \leq q-1$ can be used to define the maps $\kappa_{q-i, q+1}$, so long as $D^{(i)}y \in \Omega^{\times p^m}$ for $i \leq q-2$. We refer to a defining system so obtained as a *proper defining system* and the resulting 2-cocycle as the “ q -fold Massey product of a with b .” It is not hard to see that if $D^{(q-1)}y \in \Omega^{\times p^m}$, then ρ is a homomorphism, so the class of the 2-cocycle ν of (1.2) that defines the Massey product is trivial. In general, $D^{(q-1)}y$ determines ν through a particular transgression map (Theorem 4.3). Section 2 provides the tools needed to derive this, giving a cohomological discussion of embedding problems in a rather general context, comparing nonabelian cohomology and the sequence of base terms of the Hochschild-Serre spectral sequence. The indeterminacy in the q -fold Massey product of a with b is given by the group of $(q-1)$ -fold Massey products of a with $c \in F^\times \cap \Omega^{\times p^m}$.

1.3 Massey products in number theory

We return to our earlier setting, beginning with a number field K that we now assume to contain the p^n th roots of unity. Let S be a set of primes of K containing all primes above p and any real places. We consider Massey products for the maximal unramified outside S extension Ω of K , using the construction of the previous subsection. Theorem A arises by taking inverse limits (increasing n) from a statement at the finite level, which in turn arises from a formula for the relevant Massey products. This is the subject of Sections 5 and 6.

We quickly introduce the relevant notation. Similarly to the previous subsection, let $a \in K^\times \cap \Omega^{\times p^n}$ with $a \notin K^{\times p}$, let $b \in K^\times \cap \Omega^{\times p^m}$, let L be the field given by adjoining a p^n th root α of a to K , and let σ be a generator of L such that $\alpha^{\sigma-1} = \zeta_n$. We use $P_{m, L/K}^{(k-1)}$ to denote the group of cohomology classes of all k -fold Massey products of a with some $c \in K^\times \cap \Omega^{\times p^m}$ and set $P_{m, L/K}^{(0)} = 0$. The class of the $(k+1)$ -fold Massey product of a with b is determined up to an element of $P_{m, L/K}^{(k-1)}$ when it admits a proper defining system. We denote the class of the resulting Massey product modulo $P_{m, L/K}^{(k-1)}$ by $(a, b)_{p^m, K, S}^{(k)}$.

Kummer theory provides a natural injection

$$A_{K,S}/p^m A_{K,S} \hookrightarrow H^2(G_{K,S}, \mu_{p^m}),$$

where $A_{K,S}$ denotes the p -part of the S -class group of K . The classes of proper defining systems actually lie in the quotient of $A_{K,S}/p^m$ by the image of $P_{m,L/K}^{(k-1)}$ in this group (recalling our rigidification of μ_{p^n}). We prove a formula (Theorem 5.4) for the Massey products which essentially states the following. If $(a, b)_{p^m, K, S}^{(k)}$ is defined, any proper defining system is given by the image modulo p^m of the ideal class of the norm of a fractional ideal \mathfrak{a} in the ring of S -integers of K such that $\mathfrak{a}^{(\sigma-1)^k}$ is (nearly) principal and has a generator of norm b^{-1} . This formula is a generalization of [McS, Theorem 2.4]. We use it to prove an analogue of our main result at the level of K (Proposition 5.5).

Passing up the cyclotomic tower, we can define Massey products at the level of K_∞ . Fix a generator of the Tate module of μ_{p^∞} so that we may identify it with \mathbf{Z}_p . Write $K_n = K(\mu_{p^n})$ for each $n \geq 1$ (we no longer fix n or assume $\mu_{p^n} \subset K$). Assume, for simplicity of presentation, that L_∞ is defined by a sequence $a = (a_n)$ with

$$a_n \in K_n^\times \cap \Omega^{\times p^n}, \quad a_n \notin K_n^{\times p}, \quad \text{and} \quad a_{n+1} a_n^{-1} \in K_{n+1}^{\times p^n}.$$

Let r be such that $k < p^r(p-1)$. Given a sequence b of p -units $b_n \in K_n^\times$ for which the images in $K_n/K_n^{\times p^{n-r}}$ are norm compatible and the $(k+1)$ -fold Massey products $(a_n, b_n)_{p^{n-r}, K_n, S}^{(k)}$ admit proper defining systems, we may consider the inverse limit of these products. This defines an element $(a, b)_{K, S}^{(k)}$ of \mathcal{A}_K , the $(k+1)$ -fold Massey product of a with b . We let $\mathcal{P}_{L/K}^{(k)}$ denote the submodule generated by all such inverse limits, varying b . Theorem A (and more generally, Theorem 6.3) then follows from its analogue at the finite level.

Acknowledgments. I thank Bill McCallum for our joint work on cup products [McS] which served as the inspiration for this paper, Dick Hain for introducing Massey products to me, Barry Mazur for his encouragement and advice, and Mike Hopkins for an important observation. Section 2 is a condensed version of the first chapter of my 1999 Ph.D. thesis [Sh1], and I thank Spencer Bloch for advising me on this. This research was undertaken, in part, thanks to funding from an NSF Postdoctoral Research Fellowship, Harvard University, the Max Planck Institute for Mathematics, and the Canada Research Chairs Program.

2 Galois embedding problems

In this section, we study homomorphisms of the absolute Galois group of a field to a finite group \mathcal{H} and their lifts to a larger finite group \mathcal{G} with \mathcal{H} as a quotient. That is, we study Galois embedding problems. Our point of view differs from those which we have been able to find in the literature (see, for instance, [ILF, Ko, NSW]), so we hope that this section will serve as a useful reference. The proofs are for the most part straightforward and left to the reader.

We use the following notational conventions. For a field K , we let G_K denote the Galois group of the separable closure of K over itself. For a Galois extension L/K , we let $G_{L/K} = \text{Gal}(L/K)$.

Now fix a field K . We consider an exact sequence of finite groups of the form

$$0 \rightarrow A \xrightarrow{\iota} \mathcal{G} \xrightarrow{\phi} \mathcal{H} \rightarrow 1,$$

where A is abelian and view these groups as trivial G_K -modules. We obtain a sequence of sets

$$0 \rightarrow \text{Hom}(G_K, A) \xrightarrow{\iota^*} \text{Hom}(G_K, \mathcal{G}) \xrightarrow{\phi^*} \text{Hom}(G_K, \mathcal{H}), \quad (2.1)$$

which is exact in the sense that those elements in the image of one map are exactly those taken to the trivial homomorphism by the next. Passing to nonabelian cohomology (e.g., [Se]), we have

$$0 \rightarrow H^1(G_K, A) \xrightarrow{\iota^*} H^1(G_K, \mathcal{G}) \xrightarrow{\phi^*} H^1(G_K, \mathcal{H}). \quad (2.2)$$

As A is not necessarily central in \mathcal{G} , we cannot quite extend the sequence (2.2) to $H^2(G_K, A)$. Instead, given a homomorphism $\bar{\rho} \in \text{Hom}(G_K, \mathcal{H})$ we can twist A by $\bar{\rho}$ so that the action of $\sigma \in G_K$ on $a \in A$ is now given by

$$a^\sigma = f(\sigma)af(\sigma)^{-1},$$

where f is any continuous function lifting $\bar{\rho}$ to \mathcal{G} . We denote this new module structure by $A_{\bar{\rho}}$. We then obtain an element $\Delta(\bar{\rho}) \in H^2(G_K, A_{\bar{\rho}})$ by lifting $\bar{\rho}$ to some f as above and defining the desired 2-cocycle by

$$a(\sigma_1, \sigma_2) = f(\sigma_1)f(\sigma_2)f(\sigma_1\sigma_2)^{-1}.$$

The image of a in cohomology is $\Delta(\bar{\rho})$, and this class does not change for homomorphisms cohomologous to $\bar{\rho}$. In particular, $\bar{\rho}$ (resp., its class $[\bar{\rho}]$) will be in the image of ϕ (resp., ϕ^*) if and only if $\Delta(\bar{\rho})$ is trivial.

Remark. The class $\Delta(\bar{\rho})$ is the obstruction to lifting $\bar{\rho}$ in a very real sense. If $\Delta(\bar{\rho}) = 0$, then a is a coboundary, so we can choose $\kappa: G_K \rightarrow A$ with $d\kappa = -a$. Then, as one can easily check, $\kappa \cdot f$ is a homomorphism lifting $\bar{\rho}$.

We can give a description of $\Delta(\bar{\rho})$ in terms of group extensions. Consider the fiber product

$$\mathcal{G} \times_{\mathcal{H}} G_K = \langle (g, \sigma) \mid \phi(g) = \bar{\rho}(\sigma), g \in \mathcal{G}, \sigma \in G_K \rangle,$$

which is the pullback in the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \mathcal{G} \times_{\mathcal{H}} G_K & \longrightarrow & G_K \longrightarrow 1 & (\dagger) \\ & & \parallel & & \downarrow & & \downarrow \bar{\rho} & \\ 0 & \longrightarrow & A & \longrightarrow & \mathcal{G} & \xrightarrow{\phi} & \mathcal{H} \longrightarrow 1. \end{array}$$

Lemma 2.1. *The class of (\dagger) in $H^2(G_K, A_{\bar{\rho}})$ is $\Delta(\bar{\rho})$.*

Let L denote the fixed field of the kernel of $\bar{\rho}$. Clearly, we can also take the pullback

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \phi^{-1}(G_{L/K}) & \longrightarrow & G_{L/K} \longrightarrow 1 & (\dagger\dagger) \\ & & \parallel & & \downarrow & & \downarrow \bar{\rho} & \\ 0 & \longrightarrow & A & \longrightarrow & \mathcal{G} & \xrightarrow{\phi} & \mathcal{H} \longrightarrow 1. \end{array}$$

The group extension $(\dagger\dagger)$ yields a class $\delta(\bar{\rho}) \in H^2(G_{L/K}, A_{\bar{\rho}})$. However, this class is not necessarily invariant under conjugation of $\bar{\rho}$. Hence we do not define δ on the class $[\bar{\rho}]$. In particular, we have by Lemma 2.1 that

$$\Delta(\bar{\rho}) = \text{Inf}(\delta(\bar{\rho})),$$

where Inf denotes the inflation from $G_{L/K}$ to G_K .

Next, we would like a suitable definition of the transgression map in the Hochschild-Serre spectral sequence in terms of group extensions. This description can obviously be made in the general setting of a profinite group \mathcal{P} , an open normal subgroup \mathcal{N} , and a discrete \mathcal{P} -module M upon which \mathcal{N} acts trivially, but we restrict to the case of interest ($\mathcal{P} = G_K$, and so forth) in this discussion. So, consider the transgression map

$$\text{Tra}: H^1(G_L, A_{\bar{\rho}})^{G_{L/K}} \rightarrow H^2(G_{L/K}, A_{\bar{\rho}}).$$

This map can be described as follows (e.g., [NSW, Proposition 1.6.5]). For a homomorphism $f \in H^1(G_L, A_{\bar{\rho}})^{G_{L/K}}$, extend f to a continuous map $h: G_K \rightarrow A_{\bar{\rho}}$ satisfying

$$h(\sigma\tau) = \sigma h(\tau) + h(\sigma) \quad \text{and} \quad h(\tau\sigma) = h(\tau) + h(\sigma) \quad (2.3)$$

for all $\tau \in G_L$ and $\sigma \in G_K$. Then dh induces a well-defined 2-cocycle on $G_{L/K}$, the class of which is $\text{Tra } f$.

In terms of group extensions, we can express the transgression map as follows. For $f \in H^1(G_L, A_{\bar{\rho}})^{G_{L/K}}$, define $g : G_L \rightarrow A_{\bar{\rho}} \rtimes G_K$ by $g(\tau) = (f(\tau), \tau)$. Then $g(G_L)$ is normal in $A_{\bar{\rho}} \rtimes G_K$, so define $\mathcal{E} = (A_{\bar{\rho}} \rtimes G_K)/g(G_L)$. This yields the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & A_{\bar{\rho}} \rtimes G_K & \longrightarrow & G_K \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \mathcal{E} & \longrightarrow & G_{L/K} \longrightarrow 1 \end{array} \quad (*)$$

Alternatively, we can describe \mathcal{E} as the pushout in the following commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & G_L & \longrightarrow & G_K & \longrightarrow & G_{L/K} \longrightarrow 1 \\ & & \downarrow -f & & \downarrow & & \parallel \\ 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \mathcal{E} & \longrightarrow & G_{L/K} \longrightarrow 1 \end{array} \quad (*)$$

Lemma 2.2. *The equivalence class of $(*)$ as a group extension in $H^2(G_{L/K}, A_{\bar{\rho}})$ is $\text{Tra } f$.*

We now link the two points of view discussed above: the study of lifts of Galois representations and nonabelian cohomology, and the study of group extensions and the transgression map. Given $\rho \in \text{Hom}(G_K, \mathcal{G})$ with image $\bar{\rho} \in \text{Hom}(G_K, \mathcal{H})$ we can define $\Lambda(\rho) \in H^1(G_L, A_{\bar{\rho}})^{G_{L/K}}$ by

$$\Lambda(\rho)(\tau) = -\rho(\tau)$$

for $\tau \in G_L$. Again, we do not define Λ on the class of ρ unless A is central in \mathcal{G} .

Lemma 2.3. *We have $\text{Tra } \Lambda(\rho) = \delta(\phi(\rho))$.*

Proof. Observe the following diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & G_L & \longrightarrow & G_K & \longrightarrow & G_{L/K} \longrightarrow 1 \\ & & \downarrow -\Lambda(\rho) & & \downarrow & & \parallel \\ 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \mathcal{E} & \longrightarrow & G_{L/K} \longrightarrow 1 \\ & & \parallel & & \vdots & & \parallel \\ 0 & \longrightarrow & A_{\bar{\rho}} & \longrightarrow & \phi^{-1}(G_{L/K}) & \longrightarrow & G_{L/K} \longrightarrow 1 \\ & & \parallel & & \downarrow & & \downarrow \bar{\rho} \\ 0 & \longrightarrow & A & \longrightarrow & \mathcal{G} & \xrightarrow{\phi} & \mathcal{H} \longrightarrow 1. \end{array} \quad (2.4)$$

Note that $\phi^{-1}(G_{L/K}) = \mathcal{G} \times_{\mathcal{H}} G_{L/K}$. We define $A_{\bar{\rho}} \rightarrow \mathcal{G} \times_{\mathcal{H}} G_{L/K}$ by $a \mapsto (a, 0)$ and $G_K \rightarrow \mathcal{G} \times_{\mathcal{H}} G_{L/K}$ by $\sigma \mapsto (\rho(\sigma), \bar{\sigma})$. These maps coincide on the images of $\tau \in G_L$ because $-\Lambda(\rho)(\tau) = \rho(\tau)$. By the universal property of the pushout \mathcal{E} , we obtain a map $\mathcal{E} \rightarrow \phi^{-1}(G_{L/K})$ for which (2.4) commutes. By the Five Lemma, it is an isomorphism. \square

Lemma 2.4. *Right multiplication by ρ induces a well-defined map from the group of 1-cocycles $Z^1(G_K, A_{\bar{\rho}})$ to $\text{Hom}(G_K, \mathcal{G})$ (and, in fact, from $H^1(G_K, A_{\bar{\rho}})$ to $H^1(G_K, \mathcal{G})$). Furthermore, for $k \in Z^1(G_K, A_{\bar{\rho}})$ we have*

$$\text{Res}[k] \cdot \Lambda(\rho) = \Lambda(k \cdot \rho),$$

where Res denotes restriction from G_K to G_L . Any lift of $\bar{\rho}$ has the form $k \cdot \rho$ for some k .

Lemma 2.5. *Two lifts ρ and ρ' of $\bar{\rho}$ satisfy $\Lambda(\rho) = \Lambda(\rho')$ if and only if $\rho' = t\rho$ for some $t \in Z^1(G_{L/K}, A_{\bar{\rho}})$.*

Putting everything together, we have the following ‘‘commutative diagram’’ in which certain of the maps, which we denote with dashed arrows, must be taken only on certain elements (which map to or come from $\bar{\rho}$ or a lift ρ of it) and in which exactness holds only in the appropriate sense and where appropriate. We include it merely to summarize what we have discussed above:

$$\begin{array}{ccccccc} Z^1(G_{L/K}, A_{\bar{\rho}}) & \hookrightarrow & Z^1(G_K, A_{\bar{\rho}}) & \xrightarrow{\iota} & \text{Hom}(G_K, \mathcal{G}) & \xrightarrow{\phi} & \text{Hom}(G_K, \mathcal{H}) & \xrightarrow{-\Delta} & H^2(G_K, A_{\bar{\rho}}) \\ \downarrow & & \downarrow & & \downarrow \Lambda & & \downarrow \delta & & \parallel \\ H^1(G_{L/K}, A_{\bar{\rho}}) & \xrightarrow{\text{Inf}} & H^1(G_K, A_{\bar{\rho}}) & \xrightarrow{\text{Res}} & H^1(G_L, A_{\bar{\rho}})^{G_{L/K}} & \xrightarrow{\text{Tra}} & H^2(G_{L/K}, A_{\bar{\rho}}) & \xrightarrow{\text{Inf}} & H^2(G_K, A_{\bar{\rho}}). \end{array} \quad (2.5)$$

In particular, we obtain the following proposition which relates liftings of Galois representations to group extensions.

Proposition 2.6. *Let $\bar{\rho}: G_K \rightarrow \mathcal{H}$, and let L denote the fixed field of its kernel. Then $\bar{\rho}$ lifts to some $\rho: G_K \rightarrow \mathcal{G}$ if and only if $\delta(\bar{\rho})$ is in the image of the transgression map. If $\delta(\bar{\rho}) = \text{Tra}\xi$, then ρ may be chosen so that $\Lambda(\rho) = \xi$. The choice of ρ is unique up to left multiplication by a cocycle in $Z^1(G_{L/K}, A_{\bar{\rho}})$. Furthermore, ρ will be surjective if and only if both $\bar{\rho}$ and ξ are surjective.*

Remark. In the case that A is central in \mathcal{G} , the module $A_{\bar{\rho}}$ is just A with trivial action, and all the maps we defined above pass to cohomology classes. Hence we obtain another “commutative diagram,” similar to (2.5), in which all the maps in the top row are defined on all elements:

$$\begin{array}{ccccccc}
0 & \longrightarrow & H^1(G_K, A) & \xrightarrow{\iota^*} & H^1(G_K, \mathcal{G}) & \xrightarrow{\phi^*} & H^1(G_K, \mathcal{H}) & \xrightarrow{\Delta} & H^2(G_K, A) \\
& & \parallel & & \downarrow \Lambda & & \downarrow \delta & & \parallel \\
0 & \longrightarrow & H^1(G_{L/K}, A) & \xrightarrow{\text{Inf}} & H^1(G_K, A) & \xrightarrow{\text{Res}} & H^1(G_L, A)^{G_{L/K}} & \xrightarrow{\text{Tra}} & H^2(G_{L/K}, A) & \xrightarrow{\text{Inf}} & H^2(G_K, A).
\end{array} \tag{2.6}$$

Remark. The entire discussion already given in this section goes through with G_K replaced by $G_{\Omega/K}$ for an algebraic Galois extension Ω/K .

In the case of Ω/K , in view of the latter remark, we denote the maps corresponding to Δ and Inf in (2.5) and (2.6) with a subscript Ω . On the other hand, we use Tra_{Ω} to denote

$$\text{Tra}_{\Omega}: H^1(G_{\Omega}, A_{\bar{\rho}})^{G_{\Omega/K}} \rightarrow H^2(G_{\Omega/K}, A_{\bar{\rho}})$$

and $\Lambda_{\Omega}(\rho)$ to denote the restriction of $\Lambda(\rho)$ to G_{Ω} .

The following will be useful in Section 4.

Lemma 2.7. *Let Ω be a Galois extension of K containing L , and assume that $\bar{\rho}$ lifts to a homomorphism $\rho: G_K \rightarrow \mathcal{G}$. The obstruction to lifting the map $\bar{\rho}_{\Omega}$ induced by $\bar{\rho}$ on $G_{\Omega/K}$ to an element of $\text{Hom}(G_{\Omega/K}, \mathcal{G})$ is given by the two equal quantities*

$$\Delta_{\Omega}(\bar{\rho}_{\Omega}) = \text{Tra}_{\Omega} \Lambda_{\Omega}(\rho).$$

Proof. That $\Delta_{\Omega}(\bar{\rho}_{\Omega})$ is the obstruction is immediate. By equation (2.5) for $G_{\Omega/K}$ and G_K , we have

$$\Delta_{\Omega}(\bar{\rho}_{\Omega}) = \text{Inf}_{\Omega} \delta(\rho) = \text{Inf}_{\Omega} \circ \text{Tra} \Lambda(\rho) = \text{Tra}_{\Omega} \Lambda_{\Omega}(\rho).$$

□

3 Kummer theory

We begin this section with an algebraic lemma, which we consider in only slightly more generality than we shall need. Let G be a finite group, and let R be $\mathbf{Z}/N\mathbf{Z}$ for some N .

Let $\iota: R[G] \rightarrow R[G]$ denote the linear map determined by the inversion map $g \mapsto g^{-1}$ on G . Given a left ideal J of $R[G]$, we may consider the right ideal J^ι of $R[G]$. Let J^\perp denote the left annihilator of J^ι , which is a left ideal of $R[G]$.

Lemma 3.1. *Let J be a left ideal in $R[G]$. Then*

$$\phi: \text{Hom}(R[G]/J, \mathbf{Q}/\mathbf{Z}) \rightarrow J^\perp,$$

with $\phi(f) = \sum_{g \in G} f(g)g$, is an isomorphism of left $R[G]$ -modules.

Proof. First, we remark that

$$\phi(hf) = \sum_{g \in G} f(h^{-1}g)g = h \sum_{g \in G} f(g)g.$$

Hence, ϕ is a homomorphism of left $R[G]$ -modules as a map into $R[G]$. Since $Q = \text{Hom}(R[G]/J, \mathbf{Q}/\mathbf{Z})$ is annihilated by J^ι , the image of ϕ is contained in J^\perp . Furthermore, we claim that there exists a homomorphism of left $R[G]$ -modules, $\psi: J^\perp \rightarrow Q$, given by

$$a = \sum_{g \in G} a_g g \mapsto (g \mapsto a_g).$$

To see this, we remark that it exists as a map to $\text{Hom}(R[G], \mathbf{Q}/\mathbf{Z})$, and for $r = \sum_{g \in G} r_g g \in J$, we have

$$\psi(a)(r) = \sum_{g \in G} r_g \psi(a)(g) = \sum_{g \in G} r_g a_g,$$

which is the coefficient of 1 in $\iota(r)a = 0$. Then ψ is clearly the inverse map to ϕ . \square

Remark. If J' is a left ideal containing J , then we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(R[G]/J', \mathbf{Q}/\mathbf{Z}) & \longrightarrow & \text{Hom}(R[G]/J, \mathbf{Q}/\mathbf{Z}) & \longrightarrow & \text{Hom}(J'/J, \mathbf{Q}/\mathbf{Z}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (J')^\perp & \longrightarrow & J^\perp & \longrightarrow & J^\perp / (J')^\perp \longrightarrow 0 \end{array}$$

in which the vertical arrows are isomorphisms. We also have $(J^\perp)^\perp = J$.

We will be applying Lemma 3.1 in the setting of Kummer theory. Let K be a field of characteristic not equal p containing the group μ_{p^n} of p^n th roots of unity. Let L/K

be a Galois extension, denote its Galois group by G , and consider a subgroup Φ of L^\times which is a left $\mathbf{Z}[G]$ -module. Then the field M obtained by adjoining all p^n th roots of elements of Φ to L is Galois over K . Let

$$\bar{\Phi} = \Phi / (\Phi \cap L^{\times p^n}).$$

Kummer theory provides a (noncanonical) isomorphism

$$G_{M/L} \cong \text{Hom}(\bar{\Phi}, \mathbf{Q}/\mathbf{Z}).$$

If $\bar{\Phi}$ is generated by one element as a module, we may apply Lemma 3.1 with $R = \mathbf{Z}/p^n\mathbf{Z}$ to obtain a second description of this Galois group.

Let us consider the case that G is cyclic of order p^n with generator σ . Let

$$D_\sigma^{(k)} = (-1)^k \sum_{i=0}^{p^n-1} \binom{i}{k} \sigma^{i-k} \in \mathbf{Z}[G]$$

for $0 \leq k \leq p^n - 1$. When no confusion about the choice of generator can result, we denote $D_\sigma^{(k)}$ simply by $D^{(k)}$. Letting $\mathbf{Z}[G]$ act on L^\times in the natural way, $D^{(0)}$ acts as $N_{L/K}$, where $N_{L/K}$ denotes the norm for the extension L/K . We leave the following lemma to the reader to verify.

Lemma 3.2. *Let k be a positive integer with $k \leq p^n - 1$. Then $D^{(k)}$ satisfies*

$$(\sigma - 1)D^{(k)} = D^{(k-1)} + (-1)^k \binom{p^n}{k} \sigma^{-k} \equiv D^{(k-1)} \pmod{p^{n-v_p(k)}\mathbf{Z}[G]}.$$

Therefore, letting $r_k = \max\{r \mid k \geq p^r\}$, we have

$$(\sigma - 1)^j D^{(k)} = D^{(k-j)} + (-\sigma)^{j-1-k} \sum_{i=0}^{j-1} \binom{p^n}{k-i} (\sigma^{-1} - 1)^{j-1-i} \equiv D^{(k-j)} \pmod{p^{n-r_k}\mathbf{Z}[G]}$$

for $1 \leq j \leq k$.

Let I_G denote the augmentation ideal of $R[G]$ for a given group G and coefficient ring R .

Lemma 3.3. *Let k and m be nonnegative integers with $m + r_k \leq n$ (that is, $k \leq p^{n-m+1} - 1$), and set $R = \mathbf{Z}/p^m\mathbf{Z}$. Then the ideal $(I_G^{k+1})^\perp$ of $R[G]$ is generated by $D^{(k)}$.*

Proof. Note that $(I_G^t)^t = I_G^t$ for all t . For $k = 0$, the statement of the lemma is obvious. We have the inclusion $(I_G^k)^\perp \subset (I_G^{k+1})^\perp$. Since I_G^k/I_G^{k+1} is cyclic of order dividing p^m , so is $(I_G^{k+1})^\perp/(I_G^k)^\perp$. It is then clearly generated by the element $D^{(k)}$ of order p^m , as $(\sigma - 1)D^{(k)} = D^{(k-1)}$ in $R[G]$. \square

It is not hard to pass from Lemma 3.3 to the general case of any k , m , and n , but we shall not require this. On the other hand, in Section 5, we shall require the following lemma that describes which $D^{(k)}$ become trivial in the group rings of quotients of G .

Lemma 3.4. *Let k be a nonnegative integer and m and n be positive integers with $k \leq p^{n-m+1} - 1$. Let G be a cyclic group of order p^n with generator σ , and let H be its quotient of order p^{n-m} . Let $\phi: R[G] \rightarrow R[H]$ denote the natural map on group rings, for $R = \mathbf{Z}/p^m\mathbf{Z}$. Then we have*

$$\phi(D_\sigma^{(k)}) = \begin{cases} 0 & \text{if } k < p^{n-m}(p-1), \\ p^{m-1}D_{\phi(\sigma)}^{(k-p^{n-m}(p-1))} & \text{if } k \geq p^{n-m}(p-1). \end{cases}$$

Proof. The proof we give is combinatorial. We have

$$\phi(D_\sigma^{(k)}) = (-1)^k \sum_{i=0}^{p^{n-m}-1} \sum_{j=0}^{p^m-1} \binom{i + jp^{n-m}}{k} \phi(\sigma)^{i-k}. \quad (3.1)$$

Note that

$$\binom{i + jp^{n-m}}{k} = \sum_{a=0}^k c_{i,k}(a) (jp^{n-m})^a, \quad (3.2)$$

where $c_{i,k}(0) = \binom{i}{k}$ and, for $a > 0$,

$$c_{i,k}(a) = \sum_{0 \leq b_1 < \dots < b_a < k} \frac{1}{k!} \prod_{\substack{d=0 \\ d \notin \{b_1, \dots, b_a\}}}^{k-1} (i - d). \quad (3.3)$$

Using the facts that $i < p^{n-m}$ and $k < p^{n-m+1}$, one can obtain a lower bound on the p -adic valuations of the terms in the sum in (3.3) which yields

$$p^{a(n-m)} c_{i,k}(a) \in \begin{cases} \mathbf{Z}_p & \text{if } a \leq p-1, \\ p\mathbf{Z}_p & \text{if } a \geq p. \end{cases} \quad (3.4)$$

Furthermore, one may check that

$$\sum_{j=0}^{p^m-1} j^a \equiv \begin{cases} -p^{m-1} \pmod{p^m} & \text{if } a \geq 1 \text{ and } a \equiv 0 \pmod{p-1}, \\ 0 \pmod{p^m} & \text{if } a = 0 \text{ or } a \not\equiv 0 \pmod{p-1} \end{cases} \quad (3.5)$$

unless $p = 2$, $m \geq 2$, a is odd and $a \geq 3$, in which case the sum is $0 \pmod{p^m}$. Note that equations (3.2), (3.4), and (3.5) reduce the computation to that of $c_{i,k}(p-1)$.

Now let us consider the specific case that $k = p^{n-m+1} - 1$. One checks in a manner similar to that of (3.4) that

$$p^{(p-1)(n-m)} c_{i,p^{n-m+1}-1}(p-1) \equiv \begin{cases} 0 \pmod{p} & \text{if } i \leq p^{n-m} - 2, \\ -1 \pmod{p} & \text{if } i = p^{n-m} - 1. \end{cases} \quad (3.6)$$

Putting equations (3.2), (3.5), and (3.6) together, we obtain

$$\sum_{j=0}^{p^m-1} \binom{i + jp^{n-m}}{p^{n-m+1} - 1} \equiv \begin{cases} 0 \pmod{p^m} & \text{if } i \leq p^{n-m} - 2, \\ p^{m-1} \pmod{p^m} & \text{if } i = p^{n-m} - 1. \end{cases}$$

The result for $k = p^{n-m+1} - 1$ now follows by recalling equation (3.1), since

$$p^{m-1} D_{\phi(\sigma)}^{(p^{n-m}-1)} \equiv p^{m-1} \pmod{p^m} \mathbf{Z}[H]$$

The result for all k then follows by applying Lemma 3.2. \square

It would be nice to have a more intuitive proof of Lemma 3.4. In any case, we can derive the following from it.

Lemma 3.5. *Let k , s , m , and n be nonnegative integers satisfying $1 \leq m \leq n$, $s \leq n$, and $k \leq p^{n-m}(p-1)$. Let G be a cyclic group of order p^n with generator σ , and let H be its quotient of order p^{n-s} . Let $R = \mathbf{Z}/p^m\mathbf{Z}$, and let $\phi: R[G] \rightarrow R[H]$ be the natural map on group rings. Let J be the ideal of $R[H]$ generated by $\phi(\sigma)^{p^{n-m}} - 1$. Then*

$$\phi((I_G^k)^\perp)^\perp \subseteq I_H^k + J^\perp.$$

Proof. The cases in which $s = 0$ and $s \geq m$ are trivial, so we assume that $1 \leq s < m \leq n$. If $k = 0$, then again the statement is trivial. Assume that we know the result for $k-1$ with $k \leq p^{n-m}(p-1)$. Then we have that

$$\phi((I_G^k)^\perp)^\perp \subseteq \phi((I_G^{k-1})^\perp)^\perp \subseteq I_H^{k-1} + J^\perp.$$

Take $x \in \phi((I_G^k)^\perp)^\perp$, so that

$$x \equiv a(\phi(\sigma) - 1)^{k-1} \pmod{(I_H^k + J^\perp)}. \quad (3.7)$$

for some $a \in R$. Now $x\phi(D_\sigma^{(k-1)}) = 0$ by Lemma 3.3, so noting Lemma 3.2, we have

$$a\phi(D_\sigma^{(0)}) = ap^s D_{\phi(\sigma)}^{(0)} \in J^\perp \phi((I_G^k)^\perp). \quad (3.8)$$

We define a generator \bar{N} of J^\perp by

$$\bar{N} = \sum_{i=0}^{p^{m-s}-1} \phi(\sigma)^{p^{n-m}i}.$$

Consider the quotient H' of G of order p^{n-m} , and let ϕ' denote the map on group rings. Since

$$D_{\phi(\sigma)}^{(0)} = \bar{N} \sum_{i=0}^{p^{n-m}-1} \phi(\sigma)^i,$$

the condition of (3.8) may be rewritten as

$$ap^s D_{\phi(\sigma)}^{(0)} \in \phi'((I_G^k)^\perp).$$

Now, by Lemma 3.4, we have $\phi'((I_G^k)^\perp) = 0$, since $k \leq p^{n-m}(p-1)$. Hence, we must have $a = p^{m-s}b$ for some $b \in R$. Going back to equation (3.7), we see that

$$x \equiv bp^{m-s}(\phi(\sigma) - 1)^{k-1} \equiv b\bar{N}(\phi(\sigma) - 1)^{k-1} \equiv 0 \pmod{(I_H^k + J^\perp)},$$

finishing the proof. \square

4 Massey products

We now fix a field K of characteristic not equal to p containing μ_{p^n} for some $n \geq 1$. We also fix a Galois extension Ω/K . As before, for $k \geq 1$, let r_k be the largest integer such that $p^{r_k} \leq k$. Fix $k \leq p^n - 1$ and $m \leq n - r_k$, and consider an array of the form

$$C = (C_{i,j} = \mu_{p^m}^{\otimes j-i} \mid 1 \leq i < j \leq k+2).$$

This forms a multiplicative system as in [D, p. 183], in that there are multiplication maps

$$C_{i,j} \otimes C_{j,k} \rightarrow C_{i,k}.$$

Let $a \in K^\times \cap \Omega^{\times p^n}$ and $b \in K^\times \cap \Omega^{\times p^m}$. We will be interested in the $(k+1)$ -fold Massey product of k -copies of a with b , denoted $(a, b)_{p^m, \Omega/K}^{(k)}$ when defined, which is to say it has a defining system described as follows. A defining system is a set of continuous maps

$$\kappa_{i,j}: G_{\Omega/K} \rightarrow C_{i,j}$$

for $1 \leq i < j \leq k+2$ with $(i, j) \neq (1, k+2)$, that satisfy

- a. $\kappa_{i,i+1}$ is the Kummer character associated to a p^m th root of a for all positive $i \leq k$,
- b. $\kappa_{k+1,k+2}$ is the Kummer character associated to a p^m th root of b , and
- c. for any i and j as above and $\sigma_1, \sigma_2 \in G_{\Omega/K}$, we have

$$\kappa_{i,j}(\sigma_1 \sigma_2) = \kappa_{i,j}(\sigma_1) + \kappa_{i,j}(\sigma_2) + \sum_{l=1}^{j-i-1} \kappa_{i,l+i}(\sigma_1) \kappa_{l+i,j}(\sigma_2).$$

The $(k+1)$ -fold Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ with respect to such a defining system is the class in $H^2(G_{\Omega/K}, \mu_{p^m}^{\otimes k+1})$ of the 2-cocycle

$$(\sigma_1, \sigma_2) \mapsto \sum_{i=2}^{k+1} \kappa_{1,i}(\sigma_1) \kappa_{i,k+2}(\sigma_2). \quad (4.1)$$

Therefore, when the Massey product is defined, it is defined only up to a choice of defining system (in fact, modulo some subgroup of $H^2(G_{\Omega/K}, \mu_{p^m}^{\otimes k+1})$). Later, we shall restrict to a subset of all defining systems in order to reduce this ambiguity (and, consequently, the set of Massey products which are considered to be defined).

We may interpret defining systems for Massey products in terms of matrices. Throughout, we fix a choice of primitive p^n th root of unity ζ , which we use to make an identification $\mu_{p^n} \cong \mathbf{Z}/p^n\mathbf{Z}$. We shall later show the resulting Massey product to be independent of this choice. We envision the multiplicative system as defining the group of upper triangular unipotent matrices $\mathcal{T}_m^{(k)}$ in $GL_{k+2}(\mathbf{Z}/p^m\mathbf{Z})$. Let $\mathcal{Z}_m^{(k)}$ denote the center of this group (the cyclic subgroup generated by the matrix $Z = I + E_{1,k+2}$, where $E_{i,j}$ is the matrix which is zero in all the (i, j) th entry, which is 1). A defining system for $(a, b)_{p^m, \Omega/K}^{(k)}$ then provides a homomorphism

$$\bar{\rho}: G_{\Omega/K} \rightarrow \mathcal{T}_m^{(k)} / \mathcal{Z}_m^{(k)},$$

as in (1.1), taking $\chi_i = \kappa_{i,i+1}$. By definition of Δ_Ω as in Section 2 with $A = \mathcal{Z}^{(k)} \cong \mathbf{Z}/p^m\mathbf{Z}$, the cocycle defining $\Delta_\Omega(\bar{\rho})$ is, under the appropriate identification (taking Z to 1), the negative of the cocycle in (4.1) for the Massey product relative to the given defining system.

Let $\mathcal{G}_m^{(k)}$ be the subgroup of $\mathcal{T}_m^{(k)}$ generated by the two elements $X = I + \sum_{i=1}^k E_{i,i+1}$ and $Y = I + E_{k+1,k+2}$. The quotient H_m of $\mathcal{G}_m^{(k)}$ by the normal subgroup $\mathcal{N}_m^{(k)}$ generated by Y is cyclic of order p^{m+r_k} , generated by the image of X . We remark that $\mathcal{G}_m^{(k)} \cong \mathcal{N}_m^{(k)} \rtimes H_m$. Letting $G = H_{n-r_k}$, which has order p^n , we may view H_m canonically as a quotient of G for any fixed m with $m + r_k \leq n$.

For i with $0 \leq i \leq k$, we let $Y_i = I + E_{k+1-i,k+2}$, so that

$$Y_i = [X, [X, \dots [X, Y] \dots]],$$

where the commutator is of length $i + 1$ (using the convention $[X, Y] = XYX^{-1}Y^{-1}$). The group $\mathcal{N}^{(k)}$ is isomorphic to $(\mathbf{Z}/p^m\mathbf{Z})[G]/I_G^{k+1}$ as a $(\mathbf{Z}/p^m\mathbf{Z})[G]$ -module, under the isomorphism taking Y to 1.

Let $\alpha^{p^n} = a$, and set $L = K(\alpha)$. We assume for simplicity that $[L : K] = p^n$ throughout. Let σ generate $G_{L/K}$, and let ζ be such that $\sigma\alpha = \zeta\alpha$. For any $t \leq n$, let $\zeta_t = \zeta^{p^{n-t}}$. Assume that the norm residue symbol $(a, b)_{p^n, K}$ is trivial, which is to say that there exists $y \in L^\times$ with $N_{L/K}y = b$. We identify $G_{L/K}$ with G under the map taking σ to the image of $X \in \mathcal{G}_{n-r_k}^{(k)}$. Let Φ denote the $\mathbf{Z}[G]$ -submodule of L^\times generated by y . Let M denote the Kummer extension of L given by adjoining the p^m th roots of all elements of Φ . In the notation of Section 3, we have an identification $\bar{\Phi} \cong (\mathbf{Z}/p^m\mathbf{Z})[G]/J$ of $(\mathbf{Z}/p^m\mathbf{Z})[G]$ -modules taking the image of y to the image of 1, for some ideal J . Lemma 3.1 then implies that $G_{M/L} \cong J^\perp$ (with $R = \mathbf{Z}/p^m\mathbf{Z}$, under the isomorphism defined by ζ_m). For $j \leq k$, let M_j denote the subextension corresponding to the submodule $((I_G^{j+1})^\perp + J)/J$ of $\bar{\Phi}$. Also, let $c_j = D^{(j)}y$ and $\eta_j^{p^m} = c_j$. By Lemma 3.3, we have $M_j = L(\eta_0, \dots, \eta_j)$. We remark that $G_{M_j/L}$ has the form $J^\perp/(I_G^{j+1} \cap J^\perp)$ as a $(\mathbf{Z}/p^m\mathbf{Z})[G]$ -module.

Lemma 4.1. *For positive integers k and m with $m + r_k \leq n$, there is a homomorphism*

$$\rho^{(k)} : G_{M_k/K} \rightarrow \mathcal{G}_m^{(k)}$$

under which $G_{M_k/L}$ is considered as a subgroup of $\mathcal{N}_m^{(k)}$ via the injection

$$J^\perp/(I_G^{k+1} \cap J^\perp) \rightarrow (\mathbf{Z}/p^m\mathbf{Z})[G]/I_G^{k+1} \quad (4.2)$$

and for which, fixing a lift $\tilde{\sigma}$ of σ to $G_{M_k/K}$, we have

$$\rho^{(k)}(\tilde{\sigma}) = X + tE_{k+1,k+2},$$

where $\zeta_m^t = \lambda(\tilde{\sigma})$, where λ is the Kummer character associated to a p^m th root of b .

Proof. We must check that $\rho^{(k)}$, defined on $G_{M_k/L}$ and $\tilde{\sigma}$ as in the lemma, extends to a homomorphism on $G_{M_k/K}$ to the semidirect product $\mathcal{G}_m^{(k)}$. To see this, we begin by noting that

$$Y^{(\rho^{(k)}(\tilde{\sigma})-1)^j} = Y_j \tag{4.3}$$

for each j with $0 \leq j \leq k$.

We consider an element e of $G_{M_k/L}$ as an element of $(\mathbf{Z}/p^m\mathbf{Z})[G]/I_G^{k+1}$ via the injection in (4.2). Note that the norm element $N_G \in (\mathbf{Z}/p^m\mathbf{Z})[G]$ satisfies

$$N_G \equiv p^{m-1}(\sigma - 1)^{p^{n-m+1}-1} \pmod{I_G^{k+1}}.$$

Thus, unless $e = 0$, or $k = p^{n-m+1} - 1$ and e is a multiple of $p^{m-1}(\sigma - 1)^{p^{n-m+1}-1}$, we have that $\sigma\tilde{e} \neq \tilde{e}$ for any lift of e to

$$G_{M/L} \cong J^\perp \leq (\mathbf{Z}/p^m\mathbf{Z})[G].$$

Therefore, if $k \leq p^{n-m+1} - 2$, then $\tilde{\sigma}^{p^n} = 1$ and

$$G_{M_k/K} \cong G_{M_k/L} \rtimes G,$$

so (4.3) suffices to prove the lemma in this case.

Now take $k = p^{n-m+1} - 1$, and set $\rho = \rho^{(k)}$. We are reduced to checking that $\rho(\tilde{\sigma}^{p^n}) = \rho(\tilde{\sigma})^{p^n}$. As $\tilde{\sigma}^{p^n}$ is an element of $G_{M/L}$ fixed by the action of $G_{L/K}$, we have seen that $\rho(\tilde{\sigma}^{p^n})$ is a power of $Y_{p^{n-m+1}-1}$. Note also that

$$\rho(\tilde{\sigma})^{p^n} = Y_{p^{n-m+1}-1}^{p^{m-1}t},$$

so we need only check that

$$\lambda(\tilde{\sigma})^{p^{m-1}} = \lambda'(\tilde{\sigma}^{p^n}),$$

where λ' denotes the Kummer character associated to a p^m th root of $c_{p^{n-m+1}-1}$. This follows from the following two facts in the appropriate group rings:

$$\tilde{\sigma}^{p^n} - 1 = (\tilde{\sigma} - 1) \sum_{i=0}^{p^n-1} \tilde{\sigma}^i$$

and

$$N_G D^{(p^{n-m+1}-1)} = p^{m-1} N_G.$$

□

Lemma 4.1 has the following consequence for defining systems of Massey products using their definition in terms of matrices. For $x \in \Omega^\times$, let $[x]$ denote the cohomology class in $H^1(G_\Omega, \mu_{p^m})$ associated to x by Kummer theory.

Proposition 4.2. *Let $\rho^{(k)}$ be the homomorphism defined in Lemma 4.1. If $c_{k-1} \in \Omega^{\times p^m}$, then the induced homomorphism*

$$\bar{\rho}_\Omega^{(k)} : G_{\Omega/K} \rightarrow \mathcal{G}_m^{(k)} / \mathcal{Z}_m^{(k)}$$

provides a defining system for the $(k+1)$ -fold Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ with class

$$\text{Tra}_\Omega[c_k] \otimes \zeta_m^{\otimes k} \in H^2(G_{\Omega/K}, \mu_{p^m}^{\otimes k+1}).$$

Given y and ζ , this class does not depend upon the choice of lift $\tilde{\sigma}$ of σ .

Proof. The first statement follows by definition, using the identification given by the choice of ζ , since the assumption on c_{k-1} implies that $M_{k-1} \subseteq \Omega$. The class of the defining system is $-\Delta_\Omega(\bar{\rho}^{(k)}) \otimes \zeta_m^{\otimes k}$. By Lemma 2.7 (with $L = M_{k-1}$), we have

$$\Delta_\Omega(\bar{\rho}_\Omega^{(k)}) = \text{Tra}_\Omega \Lambda_\Omega(\rho),$$

where ρ denotes the inflation of $\rho^{(k)}$ to G_K . By definition, $\Lambda_\Omega(\rho)$ is the negative of the class of $\rho|_{G_\Omega}$. By the injection in (4.2), this class is equal to $[c_k]$. That this class does not depend upon the choice of lift $\tilde{\sigma}$ is clear. □

We make the following definitions for $m \leq n - r_k$. If there exists $y \in L^\times$ with $N_{L/K}y = b$ and $c_{k-1} \in \Omega^{\times p^m}$, then the defining system provided by y as in Proposition 4.2 is called a proper defining system. Let $U_{m,L/K}^{(k)}$ denote the group of $b \in K^\times \cap \Omega^{\times p^m}$ for which the Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ has a proper defining system. Let $P_{m,L/K}^{(0)}$ be the trivial group, and for $k \geq 1$, let $P_{m,L/K}^{(k)}$ denote the group of classes in $H^2(G_{\Omega/K}, \mu_{p^m})$ which are twists by $\zeta_m^{\otimes -k}$ of proper defining systems of $(a, b)_{p^m, \Omega/K}^{(k)}$ for any $b \in U_{m,L/K}^{(k)}$. By Proposition 4.2, we may write this more precisely as

$$P_{m,L/K}^{(k)} = \langle \text{Tra}_\Omega [D^{(k)}y] \mid y \in L^\times, D^{(k-1)}y \in \Omega^{\times p^m} \rangle.$$

When we wish to indicate the dependence of these definitions on Ω , we will write $U_{m,\Omega/L/K}^{(k)}$ and $P_{m,\Omega/L/K}^{(k)}$ in place of $U_{m,L/K}^{(k)}$ and $P_{m,L/K}^{(k)}$, respectively.

Restricting to the set of proper defining systems (for a fixed n) reduces the ambiguity in the class of the Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ to the choice of an $x \in L^\times$ with $D^{(k-2)}x \in \Omega^{\times p^m}$, as we may replace y with $N_{L/K}y = b$ by $yx^{\sigma^{-1}}$. This changes the resulting class by a twist of the class associated with the proper defining system of $(a, N_{L/K}x)_{p^m, \Omega/K}^{(k-1)}$. In particular, $P_{m,L/K}^{(k)}$ contains $P_{m,L/K}^{(k-1)}$.

From now on, we take the Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ for $b \in U_{m,L/K}^{(k)}$ to be the unique element of

$$H^2(G_{\Omega/K}, \mu_{p^m}^{\otimes k+1}) / (P_{m,L/K}^{(k-1)} \otimes \mu_{p^m}^{\otimes k})$$

associated to any proper defining system. (We will show this to be independent of the choice of ζ in Proposition 4.4. We will also examine its dependence upon n , which we exclude from the notation.) With these definitions, the following theorem, which is the main result of this section, is a direct corollary of Proposition 4.2.

Theorem 4.3. *Let $a \in K^\times \cap \Omega^{\times p^n}$, let $L = K(\alpha)$ for $\alpha \in \Omega^\times$ with $\alpha^{p^n} = a$, and assume that $[L : K] = p^n$. Let σ denote a generator of $G_{L/K}$ with $\sigma\alpha = \zeta\alpha$. Let k and m be positive integers with $m + r_k \leq n$. Let $b \in U_{m,L/K}^{(k)}$, and let $y \in L^\times$ be such that $N_{L/K}y = b$ and $D_\sigma^{(k-1)}y \in \Omega^{\times p^m}$. Then the $(k+1)$ -fold Massey product satisfies*

$$(a, b)_{p^m, \Omega/K}^{(k)} \otimes \zeta_m^{\otimes -k} = \text{Tra}_\Omega[D_\sigma^{(k)}y] \pmod{P_{m,L/K}^{(k-1)}},$$

where $\zeta_m = \zeta^{p^{n-m}}$.

We now verify that our definition of the Massey product is independent of the original choice of ζ .

Proposition 4.4. *The Massey product $(a, b)_{p^m, \Omega/K}^{(k)}$ is independent of the choice of ζ . Furthermore, $P_{m,L/K}^{(k)}$ is independent of the choice of the element $a \in K$ defining L .*

Proof. If we replace ζ by ζ^j for some j prime to p , then by its definition, σ is replaced by σ^j as well. By Lemma 3.3, both $D^{(k)}$ and $D_j^{(k)} = D_{\sigma^j}^{(k)}$ have image generating $(I_G^{k+1})^\perp / (I_G^k)^\perp$ for $R = \mathbf{Z}/p^m\mathbf{Z}$, and so they agree modulo $(I_G^k)^\perp$ up to a constant. To compute this constant, note $(\sigma - 1)^k D^{(k)} = N_G$ and

$$N_G = (\sigma^j - 1)^k D_j^{(k)} = \left(\sum_{i=0}^{j-1} \sigma^i \right)^k \cdot (\sigma - 1)^k D_j^{(k)} = j^k (\sigma - 1)^k D_j^{(k)}.$$

Since $D^{(k-1)}$ generates $(I_G^k)^\perp$ as a $(\mathbf{Z}/p^m\mathbf{Z})[G]$ -module by Lemma 3.3, we then have that

$$\mathrm{Tra}_\Omega[c_k] \equiv j^k \mathrm{Tra}_\Omega[D_j^{(k)} y] \pmod{P_{m,L/K}^{(k-1)}}$$

so

$$\mathrm{Tra}_\Omega[c_k] \otimes \zeta_m^{\otimes k} \equiv \mathrm{Tra}_\Omega[D_j^{(k)} y] \otimes (\zeta_m^j)^{\otimes k} \pmod{P_{m,L/K}^{(k-1)} \otimes \mu_{p^m}^{\otimes k}}.$$

The first statement now follows by Theorem 4.3.

Finally, we remark that replacing a by a^j but fixing σ amounts to replacing ζ by ζ^j , hence merely multiplies the Massey product by j^k , not affecting the group $P_{m,L/K}^{(k)}$. \square

We now list the relationships between Massey products that result when one of m , n , K , and Ω is allowed to vary. We leave the proofs to the reader.

Lemma 4.5. *Let*

$$Q_{m,\Omega/L/K}^{(k)} = H^2(G_{\Omega/K}, \mu_{p^m}^{\otimes k+1}) / P_{m,\Omega/L/K}^{(k-1)}$$

and take $b \in U_{m,\Omega/L/K}^{(k)}$.

- a. Fix $n' \leq n$ with $m + r_k \leq n'$, and let L' be the unique subextension of L/K of degree $p^{n'}$. Then the quotient map $Q_{m,\Omega/L/K}^{(k)} \rightarrow Q_{m,\Omega/L'/K}^{(k)}$ takes $(a, b)_{p^m,\Omega/L/K}^{(k)}$ to $(a, b)_{p^m,\Omega/L'/K}^{(k)}$.
- b. For $m' \leq m$, the canonical map $Q_{m,\Omega/L/K}^{(k)} \rightarrow Q_{m',\Omega/L/K}^{(k)}$ takes $(a, b)_{p^m,\Omega/K}^{(k)}$ to $(a, b)_{p^{m'},\Omega/K}^{(k)}$.
- c. Assume that K' is an extension of K contained in Ω with $L \cap K' = K$ and that b' is an element of $U_{m,\Omega/LK'/K'}^{(k)}$ satisfying $N_{K'/K} b' = b$. Then corestriction induces a map $Q_{m,\Omega/LK'/K'}^{(k)} \rightarrow Q_{m,n,\Omega/L/K}^{(k)}$ taking $(a, b')_{p^m,\Omega/K'}^{(k)}$ to $(a, b)_{p^m,\Omega/K}^{(k)}$.
- d. If Ω' is a Galois extension of K containing Ω , then inflation induces a map $Q_{m,\Omega/L/K}^{(k)} \rightarrow Q_{m,n,\Omega'/L/K}^{(k)}$ taking $(a, b)_{p^m,\Omega/K}^{(k)}$ to $(a, b)_{p^m,\Omega'/K}^{(k)}$.

Remark. We have assumed that $[L : K] = p^n$. It would be nice to have the case of general degree written up for completeness. We have not done this, as it would have complicated our discussion of Massey products, and it is not needed later in this paper. Note that if $(a, b)_{p^m,\Omega/K}^{(k)}$ has a proper defining system, then one should have

$$(a^i, b)_{p^m,\Omega/K}^{(k)} = i^k \cdot (a, b)_{p^m,\Omega/K}^{(k)} \pmod{P_{m,L_i/K}^{(k-1)}},$$

where $L_i = K(\alpha^i)$, for any $i \in \mathbf{Z}$ (not just prime to p).

5 Class groups

We now take K to be a number field, and we let S be a set of primes of K including those above p and all real places. Recall that we assume $\mu_{p^n} \subset K^\times$. We let $G_{K,S}$ denote the Galois group of the maximal extension Ω of K unramified outside S , and we denote the Massey products for Ω/K by $(a, b)_{p^m, K, S}^{(k)}$.

Let $\text{Cl}_{K,S}$ denote the S -class group of K , i.e., the quotient of the class group of K modulo the classes of all finite primes in S . Equivalently, it is the class group of the ring of S -integers $\mathcal{O}_{K,S}$ of K , obtained from the integer ring of K by inverting all finite primes in S . The cohomology group $H^1(G_{K,S}, \mathcal{O}_{\Omega,S}^\times)$ is generated by the classes of the cocycles f_x for $x \in \Omega^\times$ with $x\mathcal{O}_{\Omega,S} = \mathfrak{A}\mathcal{O}_{\Omega,S}$, where $f_x(\tau) = x^{\tau-1}$ for $\tau \in G_{K,S}$ and \mathfrak{A} is a fractional ideal of $\mathcal{O}_{K,S}$. The map taking the class of f_x to the ideal class of \mathfrak{A} provides a canonical isomorphism $H^1(G_{K,S}, \mathcal{O}_{\Omega,S}^\times) \cong \text{Cl}_{K,S}$ (see, for example, [NSW, Proposition 8.3.10]). Via the coboundary map

$$\partial: H^1(G_{K,S}, \mathcal{O}_{\Omega,S}^\times) \rightarrow H^2(G_{K,S}, \mu_{p^n}),$$

we have a canonical injection

$$\text{Cl}_{K,S}/p^n \text{Cl}_{K,S} \rightarrow H^2(G_{K,S}, \mu_{p^n}),$$

and we denote the image of \mathfrak{A} in this group by $[\mathfrak{A}]$.

Lemma 5.1. *Let $x \in \Omega^\times$ be such that $x\mathcal{O}_{\Omega,S}$ is the lift of a fractional ideal \mathfrak{A} of $\mathcal{O}_{K,S}$. Then $\text{Tra}_\Omega[x] = -[\mathfrak{A}]$.*

Proof. Let $\xi^{p^n} = x$. Let g denote a cochain with p^n th power equal to f_x , i.e., a cochain $g(\delta) = \xi^{\tilde{\delta}-1}$, choosing a lift $\tilde{\delta} \in G_K$ for each $\delta \in G_{K,S}$ (with $\tilde{1} = 1$). By definition, we have that

$$dg(\delta, \delta') = \frac{\xi^{\tilde{\delta}-1}(\xi^{\tilde{\delta}'-1})^{\tilde{\delta}}}{\xi^{\tilde{\delta}\tilde{\delta}'-1}} = \frac{\xi^{\tilde{\delta}-1}\xi^{\tilde{\delta}\tilde{\delta}'-\tilde{\delta}}}{\xi^{\tilde{\delta}\tilde{\delta}'-1}} = \xi^{\tilde{\delta}\tilde{\delta}'-\tilde{\delta}\tilde{\delta}'}$$

Define

$$\theta(\delta, \delta') = \tilde{\delta}\tilde{\delta}'^{-1}\tilde{\delta}\tilde{\delta}'$$

Letting f_ξ denote the p^n th Kummer cocycle on G_Ω associated to x , we see that the cocycle defining \mathfrak{A} is given by

$$dg(\delta, \delta') = (\xi^{\theta(\delta, \delta')-1})^{\tilde{\delta}\tilde{\delta}'} = f_\xi(\theta(\delta, \delta')),$$

since $\xi^{\theta(\delta, \delta')^{-1}} \in \mu_{p^n}$.

Now fix an extension $h: G_K \rightarrow \mu_{p^n}$ of f_ξ with $h(\tilde{\delta}) = 1$ for all $\delta \in G_{K,S}$ and

$$h(\tilde{\delta}\tau) = h(\tau\tilde{\delta}) = h(\tau)$$

for $\tau \in G_K$. The coboundary dh defines a cocycle on the quotient $G_{K,S}$, and the transgression $\text{Tra}_\Omega[x]$ is given, since h satisfies (2.3), by its class. Since

$$dh(\delta, \delta') = -h(\tilde{\delta}\tilde{\delta}') = -h(\tilde{\delta}\tilde{\delta}'\theta(\delta, \delta')) = -h(\theta(\delta, \delta')) = -f_\xi(\theta(\delta, \delta')),$$

we have our result. □

For a finite extension F of K , we let $H_{F,S}$ denote the p -Hilbert S -class field of F , which is the maximal unramified abelian p -extension of F in which all primes of F above those in S split completely. Let $A_{F,S}$ denote the p -part of $\text{Cl}_{F,S}$. We have an identification $A_{F,S} \cong \text{Gal}(H_{F,S}/F)$ via class field theory. We begin by recalling the following basic lemma [McS, Lemma 6.1].

Lemma 5.2. *Let L/K be a cyclic p -extension unramified outside S . The norm map $N_{L/K}: A_{L,S} \rightarrow A_{K,S}$ has image $\text{Gal}(H_{K,S}/L \cap H_{K,S})$ and kernel equal to the subgroup of $A_{L,S}$ generated by $(\sigma - 1)A_{L,S}$ and the intersection of $A_{L,S}$ with the subgroup of $\text{Gal}(H_{L,S}/K)$ generated by all decomposition groups of primes in S .*

Lemma 5.2 has the following easy corollary.

Corollary 5.3. *The map*

$$A_{L,S}/(\sigma - 1)A_{L,S} \rightarrow A_{K,S},$$

induced by $N_{L/K}$ is surjective if $L \cap H_{K,S} = K$ and injective if there is at most one prime in S that does not split completely in L/K .

As in Section 4, let $a \in K^\times \cap \Omega^{\times p^n}$, $\alpha^{p^n} = a$, and $L = K(\alpha)$, and assume $[L : K] = p^n$. Let $\sigma \in G_{L/K}$ be such that $\sigma\alpha = \zeta\alpha$ for our fixed p^n th root of unity ζ . For i with $0 \leq i \leq n$, we let L_i/K denote the unique subextension of L/K of degree p^i . Let $I_{F,S}$ denote the S -ideal group of a finite extension F of K .

Theorem 5.4. *Let k and m be positive integers with $k \leq p^{n-m}(p-1)$. Let $b \in U_{m,L/K}^{(k)}$, and let $y \in L^\times$ with $N_{L/K}y = b$ and $D^{(k-1)}y \in \Omega^{\times p^m}$. Then we may write*

$$y\mathcal{O}_{L,S} \equiv \mathfrak{Y}^{(\sigma-1)^k} \mathfrak{B} \pmod{p^m I_{L,S}},$$

with $\mathfrak{Y} \in I_{L,S}$ and $\mathfrak{B} \in I_{L_{n-m},S}$, and we have

$$(a, b)_{p^m, K, S}^{(k)} \otimes \zeta_m^{\otimes -k} = -[N_{L/K}\mathfrak{Y}] \pmod{P_{m,L/K}^{(k-1)}}$$

unless $k = p^{n-m}(p-1)$, in which case

$$(a, b)_{p^m, K, S}^{(k)} \otimes \zeta_m^{\otimes -k} = -[N_{L/K}\mathfrak{Y}] - p^{m-1}[N_{L_{n-m}/K}\mathfrak{B}] \pmod{P_{m,L/K}^{(k-1)}}.$$

Proof. The first statement is a consequence of Lemma 3.5, which can be seen as follows. For each prime ideal \mathfrak{q} of $\mathcal{O}_{K,S}$, the $\mathbf{Z}[G_{L/K}]$ -module summand $M_{\mathfrak{q}}$ of $I_{L,S}$ that is generated by the primes of $\mathcal{O}_{L,S}$ lying over \mathfrak{q} is cyclic, isomorphic to $\mathbf{Z}[H]$ for some quotient H of $G_{L/K}$. Consider the projection $y_{\mathfrak{q}}$ of $y\mathcal{O}_{L,S}$ to $M_{\mathfrak{q}}$. Since $D^{(k-1)}y \in \Omega^{\times p^m}$, we have that

$$D^{(k-1)}y_{\mathfrak{q}} \equiv 0 \pmod{p^m M_{\mathfrak{q}}}.$$

We are now in the setting of Lemma 3.5, with $R = \mathbf{Z}/p^m\mathbf{Z}$, $|H| = p^{n-s}$, and $G = G_{L/K}$. Identifying $R[H]$ with $M_{\mathfrak{q}}/p^m M_{\mathfrak{q}}$, we have that the image of $y_{\mathfrak{q}}$ is contained in $\phi((I_G^k)^\perp)^\perp$ and therefore, we may write

$$y_{\mathfrak{q}} \equiv \mathfrak{Y}_{\mathfrak{q}}^{(\sigma-1)^k} \mathfrak{B}_{\mathfrak{q}} \pmod{p^m M_{\mathfrak{q}}}$$

for some $\mathfrak{Y}_{\mathfrak{q}} \in M_{\mathfrak{q}}$ and $\mathfrak{B}_{\mathfrak{q}} \in M_{\mathfrak{q}}^{G_{L/L_{n-m}}}$. Since \mathfrak{q} does not ramify in L/K , we have that

$$M_{\mathfrak{q}}^{G_{L/L_{n-m}}} = M_{\mathfrak{q}} \cap I_{L_{n-m},S}.$$

The claim then results from taking the product over all \mathfrak{q} .

As for the second statement, Lemma 3.2 yields that

$$D^{(k)}\mathfrak{Y}^{(\sigma-1)^k} \equiv N_{L/K}\mathfrak{Y} \pmod{p^m I_{L,S}}$$

Furthermore, Lemma 3.4 implies that $D^{(k)}\mathfrak{B} \in p^m I_{L,S}$ unless $k = p^{n-m}(p-1)$, in which case

$$D^{(k)}\mathfrak{B} \equiv p^{m-1}N_{L_{n-m}/K}\mathfrak{B} \pmod{p^m I_{L,S}}.$$

Recalling the first statement of the theorem and the fact that Ω contains the p -Hilbert class field of L (and without loss of generality, replacing b by a prime-to- p power of itself), we conclude that there exists $w \in \Omega^\times$ with

$$(w^{p^m} c_k) \mathcal{O}_{\Omega, S} = \begin{cases} N_{L/K} \mathfrak{Y} & \text{if } k < p^{n-m}(p-1) \\ N_{L/K} \mathfrak{Y} \cdot N_{L_{n-m}/K} \mathfrak{B}^{p^{m-1}} & \text{if } k = p^{n-m}(p-1), \end{cases} \quad (5.1)$$

for $c_k = D^{(k)} y$ as before. By Theorem 4.3, we need only make the following computation in cohomology with μ_{p^m} -coefficients:

$$\text{Tra}_\Omega[c_k] = \text{Tra}_\Omega[w^{p^m} c_k] = \begin{cases} -[N_{L/K} \mathfrak{Y}] & \text{if } k < p^{n-m}(p-1) \\ -[N_{L/K} \mathfrak{Y}] - p^{m-1} [N_{L_{n-m}/K} \mathfrak{B}] & \text{if } k = p^{n-m}(p-1), \end{cases} \quad (5.2)$$

where the last step follows from (5.1) and Lemma 5.1. \square

Remark. The case $k = 1$ and $m = n$ of the above result is Theorem 2.4 of [McS] on cup products, giving an entirely different proof of that result which, when boiled down to what is needed for that specific case, seems simpler. (Actually, we have restricted to the case that the degree is p^n here, but for cup products, the general case follows almost immediately.)

For $m \leq n$ and $k \geq 1$, let

$$J_{m, L/K}^{(k)} = \langle \mathfrak{a} \in A_{L, S} \mid \mathfrak{a}^{(\sigma-1)^k} \in \phi(A_{L_{n-m}, S}) + p^m A_{L, S} \rangle, \quad (5.3)$$

where $\phi: A_{L_{n-m}, S} \rightarrow A_{L, S}$ is the natural map.

Proposition 5.5. *Let $k \geq 1$ and $m \geq 1$ be such that $k \leq p^{n-m}(p-1) - 1$. The map $N_{L/K}: A_{L, S} \rightarrow A_{K, S}$ induces a surjective map*

$$J_{m, L/K}^{(k)} / (J_{m, L/K}^{(k)} \cap (p^m, \sigma - 1) A_{L, S}) \rightarrow P_{m, L/K}^{(k)}$$

which is an isomorphism if there is a prime v of K in S that does not split at all in L and v is the only prime in S which does not split completely in L .

Proof. Given $\mathfrak{a} \in I_{L, S}$ with ideal class in $J_{m, L/K}^{(k)}$, we can find $b \in K^\times$ associated with \mathfrak{a} as follows. Apply $(\sigma - 1)^k$ to \mathfrak{a} , multiply it by an element of $I_{L_{n-m}, S} \cdot I_{L, S}^m$ to make

it principal, and apply $N_{L/K}$ to a generator of the resulting principal ideal to obtain b . One sees that $b \in U_{m,L/K}^{(k)}$, and then Theorem 5.4 implies that $[N_{L/K}\mathfrak{a}] \in P_{m,L/K}^{(k)}$. Hence, $N_{L/K}J_{m,L/K}^{(k)}$ modulo p^m is contained in $P_{m,L/K}^{(k)}$.

Conversely, take an element $t \in P_{m,L/K}^{(k)}$ associated to a proper defining system for some $b \in U_{m,L/K}^{(k)}$ and y as in Theorem 5.4. Then the first statement of Theorem 5.4 supplies an ideal \mathfrak{a} with class in $J_{m,L/K}^{(k)}$ such that $t = [N_{L/K}\mathfrak{a}]$, noting the equality of cohomology classes demonstrated in (5.2). Therefore, we have surjectivity.

Finally, if v is as stated, then the map in Corollary 5.3 is an isomorphism, so the kernel of the map $A_{L,S} \rightarrow A_{K,S}/p^m$ induced by norm is given by $(p^m, \sigma - 1)A_{L,S}$. Thus we have injectivity. \square

6 Iwasawa theory

Now let F be a number field, and let $K = F(\mu_p)$ if p is odd and $K = F(\mu_4)$ if $p = 2$. In this section, we pass to the cyclotomic \mathbf{Z}_p -extension $K_\infty = K(\mu_{p^\infty})$ of K , and we consider a \mathbf{Z}_p -extension L_∞ of K_∞ unramified outside a finite set of primes of K_∞ . We will consider Massey products in the inverse limits of class groups up the cyclotomic tower.

Fix any set of primes S of K containing those above p and those which ramify in L_∞/K . Let $\zeta = (\zeta_n)$ be a generator of the Tate module, and let $K_n = K(\mu_{p^n})$ for $n \geq 1$. There exists a sequence of elements $a = (a_n)_{n \geq N}$ for N sufficiently large and a nondecreasing sequence $(l_n)_{n \geq N}$ with infinite limit and $l_n \leq n$ for all n , such that $a_n \in K_n^\times \cap \Omega^{\times p^{l_n}}$, $a_n \notin K_n^{\times p}$, with

$$a_{n+1}a_n^{-1} \in K_{n+1}^{\times p^{l_n}}$$

and such that, setting $\alpha_n^{p^{l_n}} = a_n$ and $L_n = K_n(\alpha_n)$, the field L_∞ is the union of the L_n .

Remark. Any prime w in K_∞ lying over a prime other than p which is unramified in a pro- p extension F/E with E containing K_∞ must split completely in F/E , since the completion of K_∞ at w contains the unramified \mathbf{Z}_p -extension of the completion K_w .

Given this remark, we let \mathcal{A}_K denote the Galois group of the maximal pro- p unramified abelian extension of K_∞ in which all primes above p split completely, and let \mathcal{A}_L be similarly defined for L_∞ . Class field theory provides canonical isomorphisms

$$\mathcal{A}_K \cong \varprojlim A_{K_n,S} \cong \varprojlim A_{K_n,S}/p^{m_n},$$

with inverse limits taken with respect to norm maps, for any nondecreasing sequence m_n with infinite limit. Similarly, \mathcal{A}_L may be viewed as an inverse limit of the $A_{L_n, S}$ (modulo p^{m_n}) via norm maps, even though the extensions may not be Galois.

Let $G = G_{L_\infty/K_\infty}$, and let I_G denote the augmentation ideal of $\mathbf{Z}_p[[G]]$. Let $\tilde{\Lambda} = \mathbf{Z}_p[[G_{K_\infty/F}]]$. We remark that restriction induces a homomorphism

$$\mathcal{A}_L/I_G\mathcal{A}_L \rightarrow \mathcal{A}_K$$

of \mathbf{Z}_p -modules which is a homomorphism of $\tilde{\Lambda}$ -modules if L_∞/K is Galois.

Let m_n be any (fixed) nondecreasing sequence with infinite limit and satisfying $k < p^{l_n - m_n}(p - 1)$ for all (sufficiently large) n . We define

$$\mathcal{P}_{L/K}^{(k)} = \varprojlim P_{m_n, L_n/K_n}^{(k)}.$$

That $\mathcal{P}_{L/K}^{(k)}$ is well-defined and its definition is independent of the choices of l_n and m_n follows from the first three parts of Lemma 4.5. It is also independent of the choice of S , as the following lemma will make clear. Define $\mathcal{J}_{L/K}^{(k)}$ to be the kernel of $(\sigma - 1)^k$ on \mathcal{A}_L , where σ is a topological generator of G .

Lemma 6.1. *The restriction map $\mathcal{A}_L \rightarrow \mathcal{A}_K$ induces a surjective homomorphism of \mathbf{Z}_p -modules (of $\tilde{\Lambda}$ -modules if L_∞/K is Galois):*

$$\mathcal{J}_{L/K}^{(k)}/(\mathcal{J}_{L/K}^{(k)} \cap I_G\mathcal{A}_L) \rightarrow \mathcal{P}_{L/K}^{(k)}. \quad (6.1)$$

Proof. We begin by verifying that

$$\mathcal{J}_{L/K}^{(k)}/(\mathcal{J}_{L/K}^{(k)} \cap I_G\mathcal{A}_L) \cong \varprojlim J_n/(J_n \cap (p^{m_n} + I_G)A_{L_n, S}), \quad (6.2)$$

where we define J_n to be $J_{m_n, L_n/K_n}^{(k)}$ as defined in (5.3). The inverse limit in (6.2) must be interpreted appropriately as sequences of norm compatible elements of the subgroups J_n of $A_{L_n, S}$, since $N_{L_t/L_n}J_t$ need not be contained in J_n for $t > n$.

By definition, the image of $\mathcal{J}_{L/K}^{(k)}$ under the natural map to $A_{L_n, S}$ is contained in J_n . Conversely, take a sequence $\mathbf{a} = (\mathbf{a}_n)$ of norm compatible elements

$$\mathbf{a}_n \in J_n + (p^{m_n} + I_G)A_{L_n, S}.$$

Let $t > n$, and let ϕ_t denote the natural map from the S -class group of an extension of K in L_t to that of L_n . We have by definition of J_t that

$$\mathbf{a}_n^{(\sigma-1)^k} = N_{L_t/L_n}(\mathbf{a}_t^{(\sigma-1)^k}) \in N_{L_t/L_n}\phi_t(A_{L_t-m_t, S}) + p^{m_t}A_{L_n, S} + I_G^{k+1}A_{L_n, S}.$$

Note that if t is such that $t \geq m_t + n$, then we have

$$N_{L_t/L_n} \phi_t(A_{L_{t-m_t}, S}) = p^{m_t} N_{L_{t-m_t}/L_n} A_{L_{t-m_t}, S} \subseteq p^{m_t} A_{L_n, S}.$$

Therefore, noting that m_t has infinite limit, we have

$$\mathfrak{a}_n^{(\sigma-1)^k} \in \bigcap_t (p^{m_t} A_{L_n, S} + I_G^{k+1} A_{L_n, S}) = I_G^{k+1} A_{L_n, S}.$$

This yields that $\mathfrak{a}^{(\sigma-1)^k} \in I_G^{k+1} \mathcal{A}_L$, and therefore, that $\mathfrak{a} \in \mathcal{J}_{L/K}^{(k)} + I_G \mathcal{A}_L$, finishing the verification of (6.2).

Now consider the inverse limit of the surjective maps

$$\psi_n: J_n / (J_n \cap (p^{m_n} + I_G) A_{L_n, S}) \rightarrow P_{m_n, L_n / K_n}^{(k)}$$

of Proposition 5.5 (for n sufficiently large). The inverse limit of the ψ_n is also surjective, as the kernels of the ψ_n are finite abelian groups. The map in (6.1) is a surjection as the composition of the isomorphism in (6.2) with $\lim_{\leftarrow} \psi_n$. \square

Let \mathcal{H}_K and \mathcal{H}_L denote the fields such that

$$\mathcal{A}_K = G_{\mathcal{H}_K/K_\infty} \quad \text{and} \quad \mathcal{A}_L = G_{\mathcal{H}_L/L_\infty}.$$

Let $\mathcal{D}_{L/K}$ be the intersection of \mathcal{A}_L with the subgroup of $G_{\mathcal{H}_L/K_\infty}$ generated by all decomposition groups of primes of K_∞ . We also define

$$\mathcal{A}_{L/K} = \text{Gal}(\mathcal{H}_K/\mathcal{H}_K \cap L_\infty) \leq \mathcal{A}_K.$$

Lemma 6.2. *The restriction map*

$$\mathcal{A}_L / I_G \mathcal{A}_L \rightarrow \mathcal{A}_K$$

has kernel $\mathcal{D}_{L/K} / (\mathcal{D}_{L/K} \cap I_G \mathcal{A}_L)$ and image $\mathcal{A}_{L/K}$.

Proof. The result follows immediately from Lemma 5.2 by taking inverse limits. \square

Note that Lemma 6.1 implies that $\mathcal{A}_{L/K}$ contains $\mathcal{P}_{L/K}^{(k)}$ for all $k \geq 0$. We define the *decomposition-free quotient* of the k th graded piece

$$\text{gr}_G^k \mathcal{A}_L = I_G^k \mathcal{A}_L / I_G^{k+1} \mathcal{A}_L$$

in the augmentation filtration of \mathcal{A}_L by

$$\mathcal{Q}_{L/K}^{(k)} = I_G^k \mathcal{A}_L / (I_G^k \mathcal{D}_{L/K} + I_G^{k+1} \mathcal{A}_L)$$

for each $k \geq 0$. Regarding these quotients, we have the following obvious remark (which follows from Corollary 5.3).

Remark. If at most one prime above p in K_∞ does not split completely in L_∞ , then in fact $\mathcal{Q}_{L/K}^{(k)} = \text{gr}_G^k \mathcal{A}_L$ for all k .

Our main result now describes the decomposition-free quotients in terms of Massey products.

Theorem 6.3. *For any $k \geq 0$, we have a canonical isomorphism of \mathbf{Z}_p -modules*

$$\mathcal{Q}_{L/K}^{(k)} \xrightarrow{\sim} \mathcal{A}_{L/K} / \mathcal{P}_{L/K}^{(k)} \otimes_{\mathbf{Z}_p} G^{\otimes k}$$

which is an isomorphism of $\tilde{\Lambda}$ -modules if L_∞/K is Galois.

Proof. Taking the quotient of the surjection

$$\mathcal{A}_L / I_G \mathcal{A}_L \rightarrow \mathcal{A}_{L/K}$$

by the surjection in Lemma 6.1 yields a surjection

$$\mathcal{A}_L / (\mathcal{J}_{L/K}^{(k)} + I_G \mathcal{A}_L) \rightarrow \mathcal{A}_{L/K} / \mathcal{P}_{L/K}^{(k)}$$

with kernel given by the image of $\mathcal{D}_{L/K}$. By definition of $\mathcal{Q}_{L/K}^{(k)}$, we need only observe that there is a canonical isomorphism

$$\mathcal{A}_L / (\mathcal{J}_{L/K}^{(k)} + I_G \mathcal{A}_L) \otimes_{\mathbf{Z}_p} G^{\otimes k} \xrightarrow{\sim} \text{gr}_G^k \mathcal{A}_L$$

given by $\mathfrak{a} \otimes \sigma^{\otimes k} \mapsto \mathfrak{a}^{(\sigma-1)^k}$. □

This has the following immediate corollary, describing a setting in which we have an exact comparison between the augmentation filtration of \mathcal{A}_L and the group \mathcal{A}_K modulo the image of Massey products.

Corollary 6.4. *Assume that there exists a unique prime v of K_∞ that does not split completely in L_∞ and that v does not split at all in L_∞ . Then there are canonical isomorphisms*

$$\text{gr}_G^k \mathcal{A}_L \xrightarrow{\sim} \mathcal{A}_K / \mathcal{P}_{L/K}^{(k)} \otimes_{\mathbf{Z}_p} G^{\otimes k}$$

for all $k \geq 0$.

Let

$$\mathcal{U}_K = \varprojlim (\mathcal{O}_{K_n, S}^\times \otimes \mathbf{Z}_p),$$

the inverse limit being taken with respect to norm maps. This group is easily seen to satisfy

$$\mathcal{U}_K \cong \varprojlim \mathcal{O}_{K_n, S}^\times / \mathcal{O}_{K_n, S}^{\times p^n} \cong \varprojlim K_n^\times / K_n^{\times p^n},$$

the latter isomorphism following from the fact that no prime splits completely in the cyclotomic \mathbf{Z}_p -extension a number field. We write an element of \mathcal{U}_K as a sequence $b = (b_n)$ of elements $b_n \in \mathcal{O}_{K_n, S}^\times$ with reduction modulo p^n th powers forming a norm compatible sequence.

For $k \geq 1$, let

$$\mathcal{U}_{L/K}^{(k)} = \langle b \in \mathcal{U}_K \mid b_n \in U_{m_n, L_n/K_n}^{(k)} \text{ for all } n \gg 0 \rangle.$$

For $b \in \mathcal{U}_{L/K}^{(k)}$, we define Massey products

$$(a, b)_{p^\infty, K_\infty}^{(k)} \otimes \zeta^{\otimes -k} = \varprojlim (a_n, b_n)_{p^{m_n}, K_n, S}^{(k)} \otimes \zeta_{m_n}^{\otimes -k} \in \mathcal{A}_K / \mathcal{P}_{L/K}^{(k-1)}.$$

One would like to say that $\mathcal{P}_{L/K}^{(k)}$ is generated (modulo $\mathcal{P}_{L/K}^{(k-1)}$) by such Massey products. This is indeed the case.

Proposition 6.5. *For $k \geq 1$, we have an equality of groups*

$$\mathcal{P}_{L/K}^{(k)} / \mathcal{P}_{L/K}^{(k-1)} = \langle (a, b)_{p^\infty, K_\infty}^{(k)} \otimes \zeta^{\otimes -k} \mid b \in \mathcal{U}_{L/K}^{(k)} \rangle.$$

Proof. Let $\mathfrak{A} \in \mathcal{P}_{L/K}^{(k)}$. Then \mathfrak{A} is the image of some $\mathfrak{a} = (\mathfrak{a}_n) \in \mathcal{J}_{L/K}^{(k)}$ by Lemma 6.1. Pick a representative of \mathfrak{a}_n , apply $(\sigma - 1)^k$ to it, and let y_n be a generator. The element $b_n = N_{L_n/K_n} y_n$ is determined by \mathfrak{a} up to the norm of an element of $\mathcal{O}_{L_n, S}^\times$. The sequence (b_n) therefore yields a norm compatible sequence of elements in

$$\varprojlim U_{m_n, L_n/K_n}^{(k)} / N_{L_n/K_n} \mathcal{O}_{L_n, S}^\times.$$

By Theorem 5.4, we have that $\mathfrak{A} \pmod{\mathcal{P}_{L/K}^{(k-1)}}$ is the inverse limit of the Massey products

$$(a_n, b_n)_{p^{m_n}, K_n, S}^{(k)} \otimes \zeta_{m_n}^{\otimes -k}.$$

We have only to check that the map

$$\mathcal{U}_{L/K}^{(k)} \cong \varprojlim U_{m_n, L_n/K_n}^{(k)} / \mathcal{O}_{K_n, S}^{\times p^{m_n}} \rightarrow \varprojlim U_{m_n, L_n/K_n}^{(k)} / (\mathcal{O}_{K_n, S}^{\times p^{m_n}} \cdot N_{L_n/K_n} \mathcal{O}_{L_n, S}^\times)$$

is surjective, but this follows as $U_{m_n, L_n/K_n}^{(k)}$ may be replaced by $U_{m_n, L_n/K_n}^{(k)} \cap \mathcal{O}_{K_n, S}^\times$ in the inverse limits and $\mathcal{O}_{K_n, S}^\times / \mathcal{O}_{K_n, S}^{\times p^{m_n}}$ is finite. \square

We conclude this section with the following statement on which elements of \mathcal{U} are trivial under all Massey products with a .

Proposition 6.6. *Let $\mathcal{U}_{L/K} = \bigcap_{k \geq 1} \mathcal{U}_{L/K}^{(k)} \subseteq \mathcal{U}_K$. Then*

$$\mathcal{U}_{L/K} = \langle b \in \mathcal{U} \mid b_n \in N_{L_n/K_n}(\mathcal{O}_{L_n, S}^\times \otimes \mathbf{Z}_p) \text{ for all } n \geq N \rangle. \quad (6.3)$$

Proof. The group on the right-hand side of (6.3) is contained in $\mathcal{U}_{L/K}$ by Theorem 5.4. Assume, on the other hand, that $b \in \mathcal{U}_{L/K}$. Fix $n \geq N$ and choose k large enough so that $I_G^{k-1} A_{L_n, S} = 0$. For any $t \geq n$ satisfying $t - m_t \geq r_k$, we have that $b_t \in U_{m_t, L_t/K_t}^{(k)}$. Hence, Theorem 5.4 yields that $b_t = N_{L/K} y_t$ for some $y_t \in L_t^\times$ with $y_t \mathcal{O}_{L_t, S} = \mathfrak{Y}_t^{(\sigma-1)^k} \mathfrak{B}_t$, where $\mathfrak{Y}_t \in I_{L_t, S}$ and $\mathfrak{B}_t \in I_{L_t - m_t, S}$. Let $y_n = N_{L_t/L_n} y_t$. Then we have

$$y_n \mathcal{O}_{L_n, S} = (N_{L_t/L_n} \mathfrak{Y}_t)^{(\sigma-1)^k} N_{L_t/L_n} \mathfrak{B}_t$$

and $N_{L_n/K_n} y_n = b_n v^{p^n}$ for some $v \in \mathcal{O}_{K_n, S}^\times$. If $t - m_t \geq n$ as well, then $N_{L_t/L_n} \mathfrak{B}_t \in p^{m_t} I_{L_n, S}$. As in the proof of Lemma 6.1, since m_t can be made arbitrarily large and n was arbitrary, we see that $y_n \mathcal{O}_{L_n, S} = \mathfrak{C}^{(\sigma-1)^k}$ for some $\mathfrak{C} \in I_{L_n, S}$. Since $I_G^{k-1} A_{L_n, S}$ is trivial by assumption, some prime to p -power of y_n (which we assume without loss of generality to be 1) has the form $u \cdot x^{\sigma-1}$ for some $u \in \mathcal{O}_{L_n, S}^\times$ and $x \in L_n^\times$, which tells us that $b_n = N_{L_n/K_n}(uv^{-1})$. \square

Remark. Let \mathcal{U}_L be the group $\varprojlim (\mathcal{O}_{L_n, S}^\times \otimes \mathbf{Z}_p)$ of universal norm sequences from L_∞ . There is an obvious norm map

$$N_{L_\infty/K_\infty} : \mathcal{U}_L \rightarrow \mathcal{U}_K.$$

As a consequence of Proposition 6.6, we have

$$\mathcal{U}_{L/K} = N_{L_\infty/K_\infty} \mathcal{U}_L,$$

the latter group being easily seen to be equal to the group on the right-hand side of (6.3).

7 Examples

In this section, we apply the results of Section 6 to the case in which $F = \mathbf{Q}$ and $K = \mathbf{Q}(\mu_p)$ for an odd prime p . Let S be the set consisting of the unique prime $(1 - \zeta_1)$ above p in K . We take L_∞ to be a \mathbf{Z}_p -extension of $K_\infty = \mathbf{Q}(\mu_{p^\infty})$ which is Galois over \mathbf{Q} and unramified outside p . Let $G = G_{L_\infty/K_\infty}$ and $\mathcal{G} = G_{L_\infty/\mathbf{Q}}$. We remark that \mathcal{A}_K is equal to the Galois group of the maximal pro- p unramified abelian extension of K_∞ .

Let $T = \mathbf{Z}_p \times \mathbf{Z}/(p-1)\mathbf{Z}$. We view \mathbf{Z} as a subgroup of T via the diagonal map. For $n \geq 1$, we have a natural surjection $T \rightarrow \mathbf{Z}/p^{n-1}(p-1)\mathbf{Z}$ which allows us to speak of congruences modulo $p^{n-1}(p-1)$ between elements of T , and we let t_n denote a lift to \mathbf{Z} of the image of $t \in T$ under this map. For $t \in T$, define

$$\mathbf{Z}_p(t) = \lim_{\leftarrow} (\mathbf{Z}/p^n\mathbf{Z})(t_n).$$

Let $\Delta = \text{Gal}(K/\mathbf{Q})$, let ω denote the Teichmüller character, and set

$$\epsilon_i = \sum_{\delta \in \Delta} \omega(\delta)^{-i} \delta \in \mathbf{Z}_p[\Delta]$$

for $i \in T$. We have the following corollary of Theorem 6.3.

Lemma 7.1. *For $i \in \mathbf{Z}$ and $k \geq 0$, there is a canonical isomorphism of $\tilde{\Lambda}$ -modules:*

$$\epsilon_{i+kt}(\text{gr}_G^k \mathcal{A}_L) \otimes_{\mathbf{Z}_p} G^{\otimes -k} \cong \epsilon_i(\mathcal{A}_{L/K}/\mathcal{P}_{L/K}^{(k)})$$

For $t \in T$ with t odd (modulo $p-1$), define

$$\lambda_{t,n} = \prod_{\substack{i=1 \\ (i,p)=1}}^{p^n-1} (1 - \zeta_n^i)^{it^{n-1}}.$$

Then

$$\lambda_{t,n+1} \lambda_{t,n}^{-1} \in K_{n+1}^{\times p^n}.$$

We fix an odd $t \in T$ and assume for now that L_∞/K_∞ is defined by the sequence $\lambda_t = (\lambda_{t,n})_{n \geq 1}$. The field L_∞ is then a \mathbf{Z}_p -extension of K_∞ unramified outside p which satisfies $G \cong \mathbf{Z}_p(t)$ as a $\tilde{\Lambda}$ -module. We remark that L_∞/K_∞ is totally ramified if $\epsilon_t \mathcal{A}_K = 0$.

Let A_K be the p -part of the class group of K , so $A_K = A_{K,S}$. Recall that Vandiver's conjecture states that the real part A_K^+ of A_K is zero and implies the weaker statement

that $\epsilon_s A_K$ is cyclic for all odd s . Vandiver's conjecture is known to hold for all $p < 12,000,000$ [BCE]. We assume the weaker condition of cyclicity of an odd eigenspace of A_K in the following proposition (see [Ku] for a discussion of this condition). For $r \in \mathbf{Z}$, $n \geq 1$ (or ∞), and $a, b \in \mathcal{O}_{K_n, S}^\times$, we let

$$\langle a, b \rangle_{p^n, r} = \epsilon_{2-r}(a, b)_{p^n, K_n, S}^{(1)}$$

(see [McS] for a study of this cup product pairing in the case $n = 1$).

Proposition 7.2. *Let r be even such that $\epsilon_{1-r} A_K$ has p -rank 1. Let M be a $\mathbf{Z}_p[[\mathcal{G}]]$ -submodule of \mathcal{A}_L generated by a choice of element with restriction generating $\epsilon_{1-r} \mathcal{A}_{L/K}$. If $\langle \lambda_{t,1}, \lambda_{r-t,1} \rangle_{p,r}$ is nonzero, then $M \cong \epsilon_{1-r} \mathcal{A}_K$.*

Proof. The condition of $\epsilon_{1-r} A_K$ having p -rank 1 is equivalent to that of $\epsilon_{1-r} \mathcal{A}_K$ being principally generated as a $\Lambda = \mathbf{Z}_p[[G_{K_\infty/K}]]$ -module [Ku, Remark 1.5]. Set

$$X = \epsilon_{1-r}(\mathcal{A}_K / \mathcal{P}_{L/K}^{(1)}).$$

Let \mathfrak{m} denote the maximal ideal of Λ . The assumption on the pairing implies that

$$\epsilon_{2-r}(A_K \otimes \mu_p) / \langle \langle \lambda_{t,1}, \lambda_{r-t,1} \rangle_{p,r} \rangle$$

is trivial. Now, $X/\mathfrak{m}X$ is isomorphic to a quotient of this, hence is trivial, which in turn implies $X = 0$. On the other hand, Lemma 7.1 yields

$$\mathrm{gr}_G^1 M \cong \epsilon_{1-r+t}(\mathrm{gr}_G^1 \mathcal{A}_L) \cong \epsilon_{1-r}(\mathcal{A}_{L/K} / \mathcal{P}_{L/K}^{(1)}) \hookrightarrow X.$$

Hence, we have $I_G M = 0$ and $\epsilon_{1-r} \mathcal{A}_{L/K} = \epsilon_{1-r} \mathcal{A}_K$, and, therefore, we have

$$M = M / I_G M \cong \epsilon_{1-r} \mathcal{A}_K.$$

□

We now apply Proposition 7.2 to one case in which the pairing $\langle \cdot, \cdot \rangle_{p,r}$ has been computed (see [McS]). Let \mathcal{Z}_L denote the Galois group of the maximal unramified abelian pro- p extension of L_∞ .

Proposition 7.3. *For $p = 37$ and $t \in T$ with $t \not\equiv 5, 27 \pmod{36}$, we have isomorphisms of $\mathbf{Z}_p[[\mathcal{G}]]$ -modules,*

$$\mathcal{A}_L \cong \mathcal{A}_K \cong \mathbf{Z}_p(s),$$

for some $s \in T$ with $s \equiv 5 \pmod{36}$. If, furthermore, $t \not\equiv 31 \pmod{36}$, then $\mathcal{Z}_L \cong \mathbf{Z}_p(s)$ as well.

Proof. Note that $p \mid B_{32}$. By [McS, Theorem 7.5], the pairing value $\langle \lambda_{t,1}, \lambda_{32-t,1} \rangle_{37,32}$ is nonzero unless $t \not\equiv 5, 27 \pmod{36}$. (Note that if $t \equiv 5 \pmod{36}$, then L_∞/K_∞ is not totally ramified at 37.) It follows from Proposition 7.2 that $\mathcal{A}_L \cong \mathcal{A}_K$. Furthermore, the λ -invariant of $\mathcal{A}_K = \epsilon_5 \mathcal{A}_K$ is 1, so $\mathcal{A}_K \cong \mathbf{Z}_p(s)$ for some $s \equiv 5 \pmod{36}$.

Note that

$$\mathcal{A}_L \cong \mathcal{Z}_L / \langle \mathfrak{p} \rangle,$$

where \mathfrak{p} denotes the class of the unique prime of L_∞ above p . Since L_∞/K_∞ is totally ramified at p , we have

$$\mathcal{Z}_L / I_G \mathcal{Z}_L \cong \mathcal{A}_L / I_G \mathcal{A}_L \cong \mathcal{A}_K.$$

In particular, we have $\mathfrak{p} \in I_G \mathcal{Z}_L$. The fact that $I_G \mathcal{A}_L = 0$ implies that $I_G \mathcal{Z}_L$ is generated by \mathfrak{p} and that

$$I_G \mathcal{Z}_L \cong \epsilon_{5+t}(I_G \mathcal{Z}_L / I_G^2 \mathcal{Z}_L).$$

Since $\mathfrak{p} \in \mathcal{Z}_L^{\mathcal{G}}$, we conclude that $I_G \mathcal{Z}_L = 0$ if $t \not\equiv 31 \pmod{36}$. \square

We now illustrate that \mathcal{A}_L can have larger p -rank than \mathcal{A}_K as well.

Proposition 7.4. *Let $r \in T$, and assume that $\epsilon_{1-r} \mathcal{A}_K \cong \mathbf{Z}_p(1-r)$ and $\epsilon_r \mathcal{A}_K = \epsilon_{2r-1} \mathcal{A}_K = 0$. Let L_∞ be the \mathbf{Z}_p -extension of K_∞ defined by λ_{2r-1} . Then we have*

$$\epsilon_{1-r}(\mathrm{gr}_G^k \mathcal{A}_L \otimes_{\mathbf{Z}_p} G^{\otimes -k}) \cong \mathbf{Z}_p(1-r).$$

if $k = 0$ or 1 , and for $k = 2$ if $3r \neq 2$.

Proof. For $k = 0$, the result is Lemma 6.2.

For $k = 1$, let $1 - \zeta$ denote the universal norm for K_∞/K given by the sequence $(1 - \zeta_n)$. We remark that $\epsilon_{1-r} \mathcal{P}_{L/K}^{(1)}$ is generated as a Λ -module by $\langle \lambda_{2r-1}, 1 - \zeta \rangle_{p^\infty, r}$, since $\epsilon_r \mathcal{A}_K = 0$ implies that $\epsilon_r \mathcal{U}_K$ is generated by universal norms arising from cyclotomic units. At the level of K_n , we have

$$\langle \lambda_{2r-1, n}, 1 - \zeta_n \rangle_{p^n, r} = \langle 1 - \zeta_n, \lambda_{1-r, n} \rangle_{p^n, r}, \quad (7.1)$$

from which we conclude that

$$\epsilon_{1-r} \mathcal{P}_{L/K}^{(1)} = \epsilon_{1-r} \mathcal{P}_{H/K}^{(1)},$$

where H_∞/K_∞ is the \mathbf{Z}_p -extension defined by λ_{1-r} . The extension H_∞/K_∞ is unramified. Using Lemma 5.2, and defining \mathcal{A}_H to be the Galois group of the maximal pro- p abelian unramified extension of H_∞ in which all primes above p split completely, we see by Lemma 6.2 that

$$\epsilon_{1-r}(\mathcal{A}_H/I_G\mathcal{A}_H) = 0.$$

Since the right hand term of (7.1) generates the ϵ_{1-r} -eigenspace of $P_{n,H_n/K_n}^{(1)}$, where H_n/K_n is the extension given by a p^n th root of $\lambda_{1-r,n}$. Since

$$\mathcal{P}_{H/K}^{(1)} = \varprojlim P_{n,H_n/K_n}^{(1)},$$

Theorem 5.4 implies that $\epsilon_{1-r}\mathcal{P}_{H/K}^{(1)}$ is trivial.

For $k = 2$, we note that the p -completion of the p -units in K_n is topologically generated by ζ_n and those p -units fixed by complex conjugation. But this means only even eigenspaces and the eigenspace of $\mathcal{P}_{L/K}^{(2)}$ generated by

$$(\lambda_{2r-1}, \zeta)_{p^\infty, K_\infty}^{(2)} \otimes \zeta^{\otimes -2}$$

can be nontrivial. Galois acts on the latter element by the $(3 - 4r)$ th power of the cyclotomic character. On the other hand, this element generates a subgroup of $\epsilon_{1-r}\mathcal{A}_K \cong \mathbf{Z}_p(1 - r)$ by Lemma 7.1. Since $3 - 4r = 1 - r$ if and only if $3r = 2 \in T$, we are done. \square

Remark. The condition of λ -invariant 1 implicitly assumed in Proposition 7.4 is also known to hold for all nontrivial eigenspaces of \mathcal{A}_K for $p < 12,000,000$ [BCE]. It is not expected to hold in general.

Venjakob [V2, Section 8] and Coates [C, Section 3] asked if \mathcal{Z}_L is always pseudo-null as a $\mathbf{Z}_p[[\mathcal{G}]]$ -module (see [V1, Definition 3.1] for the definition). By [V2, Proposition 6.5], the $\mathbf{Z}_p[[\mathcal{G}]]$ -module \mathcal{Z}_L is pseudo-null if and only if it (or, equivalently, \mathcal{A}_L) is torsion as a $\mathbf{Z}_p[[G]]$ -module (since \mathcal{Z}_L is finitely generated as a $\mathbf{Z}_p[[G]]$ -module). This would be a natural extension of a conjecture of Greenberg's [G, Conjecture 3.5] to this noncommutative case. As is demonstrated in a forthcoming paper with Hachimori [HS], this is not so if we take L_∞ to be a CM-field and the (-1) -eigenspace $\mathcal{A}_{\bar{K}}$ of \mathcal{A}_K under complex conjugation has \mathbf{Z}_p -rank at least 2. In fact, a result of considerably greater generality is proven. We give another proof of the above-mentioned result (slightly weakened for simplicity) here.

Proposition 7.5. *Let L_∞ be a CM \mathbf{Z}_p -extension of K_∞ which is Galois over K , unramified outside p , and totally ramified at p . Then \mathcal{A}_L^- is not pseudo-null if the \mathbf{Z}_p -rank of \mathcal{A}_K^- is at least 2.*

Proof. Note that G is fixed under complex conjugation, since L is CM. It is easily checked that

$$\mathrm{rank}_{\mathbf{Z}_p[[G]]} \mathcal{A}_L^- = \lim_{k \rightarrow \infty} \mathrm{rank}_{\mathbf{Z}_p} \mathrm{gr}_G^k \mathcal{A}_L^-.$$

By Theorem 6.3, this yields

$$\mathrm{rank}_{\mathbf{Z}_p[[G]]} \mathcal{A}_L^- = \mathrm{rank}_{\mathbf{Z}_p} \mathcal{A}_K^- - \lim_{k \rightarrow \infty} \mathrm{rank}_{\mathbf{Z}_p} (\mathcal{P}_{L/K}^{(k)})^-, \quad (7.2)$$

since L/K is totally ramified at p . Now \mathcal{U}_K^- is simply the Tate module of p -power roots of unity, and the Kummer generator a of L_∞ lies in the (-1) -eigenspace of the pro- p completion of K_∞^\times . Hence $(\mathcal{P}_{L/K}^{(k)})^-$ is generated over \mathbf{Z}_p by $(a, \zeta)_{p^\infty, K_\infty, S}^{(k)} \otimes_{\mathbf{Z}_p} \zeta^{\otimes -k}$. By (7.2), we are done. \square

Note that \mathcal{A}_K^- has \mathbf{Z}_p -rank (at least) 2 for $p = 157, 353, 379, \dots$. The CM condition implies that L_∞ is *not* generated by a sequence of p -power roots of p -units over K_∞ . In the situation that L_∞/K_∞ is generated by cyclotomic p -units, it is still an open question whether \mathcal{Z}_L is always pseudo-null or not. We phrase it as a conjecture (see [HS] for a related question).

Conjecture 7.6. *Let L_∞ be a \mathbf{Z}_p -extension of $K_\infty = \mathbf{Q}(\mu_{p^\infty})$ unramified outside p and generated over K_∞ by p -power roots of p -units in K_∞ . Then \mathcal{Z}_L is pseudo-null.*

Proposition 7.3 provides examples in which \mathcal{Z}_L is indeed pseudo-null. In [Sh2], we establish a sufficient and computable condition for the nontriviality of the cup product pairing which holds at least for $p < 1000$, in particular furnishing a much larger class of examples of this form. Beyond pseudo-nullity, it would be interesting to know if it ever occurs that $I_G^k \mathcal{Z}_L$ is nontrivial for all k when L_∞ arises from roots of p -units.

We end by remarking on a possible counterexample to Conjecture 7.6.

Remark. In [W, Proposition 3.1] and [NQD, Exemple 3.2], it was claimed that the Galois group of the maximal pro- p abelian unramified extension of K_∞ (in which all primes above p split completely) is free pro- p if Vandiver's conjecture holds at p . Under Vandiver's conjecture, any unramified \mathbf{Z}_p -extension L_∞/K_∞ is generated by p -power

roots of cyclotomic p -units. Assuming the freeness result, it is easily seen that \mathcal{Z}_L is not pseudo-null whenever the \mathbf{Z}_p -rank of \mathcal{A}_K is at least 2 and Vandiver’s conjecture holds at p . (We thank Manabu Ozaki for pointing out this example.) Unfortunately, however, the proofs of the above-mentioned result both have subtle but vital mistakes. For example, the Galois group of the maximal unramified pro-157 extension of K_∞ for $p = 157$ is actually isomorphic to $\mathbf{Z}_{157} \oplus \mathbf{Z}_{157}$ (note that 157 divides B_{62} and B_{110}). This follows from Proposition 7.2 and the nontriviality of the cup product $\langle \lambda_{47,1}, \lambda_{15,1} \rangle_{157,62}$ (for this nontriviality, see [McS, Sh2]).

References

- [BCE] J. Buhler, R. Crandall, R. Ernvall, T. Metsänkylä, M. A. Shokrollahi, Irregular primes and cyclotomic invariants to 12 million, *J. Symbolic Comput.* **31** (2001), 89–96.
- [C] J. Coates, Iwasawa algebras and arithmetic, Séminaire Bourbaki, no. 896, November 2001.
- [CS] J. Coates and R. Sujatha, Fine Selmer groups of elliptic curves over p -adic Lie extensions, *Math. Ann.* **331** (2005), 809–839.
- [D] W. G. Dwyer, Homology, Massey products and maps between groups, *J. Pure Appl. Algebra* **6** (1975), 177–190.
- [G] R. Greenberg, Iwasawa theory—past and present, Class Field Theory—Its Centenary and Prospect, Adv. Stud. Pure Math., vol. 30, Math. Soc. Japan, Tokyo, 2001, 335–385.
- [HS] Y. Hachimori and R. Sharifi, On the failure of pseudo-nullity of Iwasawa modules, *J. Algebraic Geom.* **14** (2005), 567–591.
- [ILF] V. Ishkhanov, B. Lur’e, D. Faddeev, The Embedding Problem in Galois Theory, Translations of Mathematical Monographs, vol. 165, Amer. Math. Soc., Providence, 1997.
- [Ko] H. Koch, Algebraic Number Theory, Springer, Berlin, 1997.

- [Ku] M. Kurihara, Ideal class groups of cyclotomic fields and modular forms of level 1, *J. Number Theory* **45** (1993), 281–294.
- [McS] W. McCallum and R. Sharifi, A cup product in the Galois cohomology of number fields, *Duke Math. J.* **120** (2003), 269–310.
- [NQD] T. Nguyen Quang Do, K_3 et formules de Riemann-Hurwitz p -adiques, *K-theory* **7** (1993), 429–441.
- [NSW] J. Neukirch, A. Schmidt, K. Wingberg, Cohomology of Number Fields, Springer, Berlin, 2000.
- [Se] J.-P. Serre, Galois Cohomology, Springer, New York, 1997.
- [Sh1] R. Sharifi, Twisted Heisenberg representations and local conductors, Ph.D. Thesis, The University of Chicago, 1999.
- [Sh2] R. Sharifi, Iwasawa Theory and the Eisenstein Ideal, preprint, arXiv:math.NT/0501236, to appear in *Duke Math. J.*
- [V1] O. Venjakob, On the structure theory of the Iwasawa algebra of a p -adic Lie group, *J. Eur. Math. Soc.* **4** (2002), 271–311.
- [V2] O. Venjakob, A noncommutative Weierstrass preparation theorem and applications to Iwasawa theory, *J. Reine angew. Math.* **559** (2003), 153–191.
- [W] K. Wingberg, On the maximal unramified p -extension of an algebraic number field, *J. Reine angew. Math.* **440** (1993), 129–156.

Department of Mathematics and Statistics
 McMaster University
 1280 Main Street West
 Hamilton, Ontario L8S 4K1
 Canada
 email address: sharifi@math.mcmaster.ca