

Matricial R-circular Systems and Random Matrices

Romuald Lenczewski

Instytut Matematyki i Informatyki
Politechnika Wrocławska

UC Berkeley, March 2014

Motivations:

- describe asymptotic distributions of random blocks
- construct new random matrix models
- unify concepts of noncommutative independence

- 1 If $Y(u, n)$ is a standard complex Hermitian Gaussian random matrix, it converges under the trace to a **semicircular** operator

$$\lim_{n \rightarrow \infty} Y(u, n) \rightarrow \omega(u)$$

under $\tau(n) = \mathbb{E} \circ \text{Tr}(n)$

- 2 If $Y(u, n)$ is a standard complex non-Hermitian Gaussian random matrix, it converges to a **circular** operator

$$\lim_{n \rightarrow \infty} Y(u, n) \rightarrow \eta(u)$$

under $\tau(n) = \mathbb{E} \circ \text{Tr}(n)$.

- ① free probability and freeness
- ② operator-valued free probability and freeness with amalgamation
- ③ matricially free probability and matricial freeness

- 1 Complex independent Hermitian Gaussian random matrices converge to a **free semicircular family**

$$\{Y(u, n) : u \in \mathcal{U}\} \rightarrow \{\omega(u) : u \in \mathcal{U}\}$$

- 2 Complex independent Non-Hermitian Gaussian random matrices converge to a **free circular family**

$$\{Y(u, n) : u \in \mathcal{U}\} \rightarrow \{\eta(u) : u \in \mathcal{U}\}$$

- 3 Generalizations (Dykema, Schlakhtyenko, Hiai-Petz, Benaych-Georges and others)

By asymptotic freeness, large random matrix is a free random variable, so it is natural to

- ① decompose it into large blocks
- ② look for a concept of 'matricial' independence for blocks
- ③ find operatorial realizations for large blocks
- ④ reduce computations to properties of these operators

- 1 decompose the set $[n] := \{1, \dots, n\}$ into disjoint intervals

$$[n] = N_1 \cup \dots \cup N_r$$

and set $n_j = |N_j|$,

- 2 use **normalized partial traces**

$$\tau_j(n) = \mathbb{E} \circ \text{Tr}_j(n)$$

where

$$\text{Tr}_j(n)(A) = \frac{n}{n_j} \text{Tr}(n)(D_j A D_j)$$

and D_j is the $n_j \times n_j$ unit matrix embedded in $M_n(\mathbb{C})$ at the right place.

- 1 decompose random matrices $Y(u, n)$ into independent blocks

$$S_{i,j}(u, n) = D_i Y(u, n) D_j$$

- 2 decompose symmetric blocks $T_{i,j}(u, n)$, built from blocks of the same color:

$$Y(u, n) = \begin{pmatrix} S_{1,1}(u, n) & S_{1,2}(u, n) & \dots & S_{1,r}(u, n) \\ S_{2,1}(u, n) & S_{2,2}(u, n) & \dots & S_{2,r}(u, n) \\ \cdot & \cdot & \ddots & \cdot \\ S_{r,1}(u, n) & S_{r,2}(u, n) & \dots & S_{r,r}(u, n) \end{pmatrix}$$

In order to define a 'matricial' concept of independence,

- 1 replace families of variables and subalgebras by arrays

$$\{a_i, i \in I\} \rightarrow (a_{i,j})_{(i,j) \in J}$$

$$\{\mathcal{A}_i, i \in I\} \rightarrow (\mathcal{A}_{i,j})_{(i,j) \in J}$$

- 2 replace one distinguished state in a unital algebra by an array of states

$$\varphi \rightarrow (\varphi_{i,j})_{(i,j) \in J}$$

where we set $\varphi_{i,j} = \varphi_j$ (state 'under condition' j)

The definition of **matricial freeness** is based on two conditions

① 'freeness condition'

$$\varphi_{i,j}(a_1 a_2 \dots a_n) = 0$$

where $a_k \in \mathcal{A}_{i_k, j_k} \cap \text{Ker} \varphi_{i_k, j_k}$ and neighbors come from different algebras

② 'matriciality condition': subalgebras are not unital, but they have internal units $1_{i,j}$, such that the unit condition

$$1_{i,j} w = w$$

holds only if w is a 'reduced word' matricially adapted to (i,j) and otherwise it is zero.

The concept of matricial freeness allows to

- 1 unify the main notions of independence
- 2 give a unified approach to sums and products of a large class of independent random matrices
- 3 find a unified combinatorial description of limit distributions (non-crossing colored partitions)
- 4 derive explicit formulas for arbitrary multiplicative convolutions of Marchenko-Pastur laws
- 5 find random matrix models for boolean independence, monotone independence and s -freeness (noncommutative independence defined by subordination)
- 6 construct a natural random matrix model for free Meixner laws

Definition

By the **matricially free Fock space of tracial type** we understand

$$\mathcal{M} = \bigoplus_{j=1}^r \mathcal{M}_j,$$

where each summand is of the form

$$\mathcal{M}_j = \mathbb{C}\Omega_j \oplus \bigoplus_{m=1}^{\infty} \bigoplus_{\substack{j_1, \dots, j_m \\ u_1, \dots, u_m}} \mathcal{H}_{j_1, j_2}(u_1) \otimes \dots \otimes \mathcal{H}_{j_m, j}(u_m),$$

Definition

Define **matricially free creation operators** on \mathcal{M} as partial isometries with the action onto basis vectors

$$\begin{aligned}\wp_{i,j}(u)\Omega_j &= e_{i,j}(u) \\ \wp_{i,j}(u)(e_{j,k}(s)) &= e_{i,j}(u) \otimes e_{j,k}(s) \\ \wp_{i,j}(u)(e_{j,k}(s) \otimes w) &= e_{i,j}(u) \otimes e_{j,k}(s) \otimes w\end{aligned}$$

for any $i, j, k \in [r]$ and $u, s \in \mathcal{U}$, where $e_{j,k}(s) \otimes w$ is a basis vector. Their actions onto the remaining basis vectors give zero.

Relations

One square matrix of creation operators $(\varphi_{i,j})$ gives an array of partial isometries satisfying relations

$$\sum_{j=1}^r \varphi_{i,j} \varphi_{i,j}^* = \varphi_{k,i}^* \varphi_{k,i} - \varphi_i \quad \text{for any } k$$

$$\sum_{j=1}^r \varphi_{k,j}^* \varphi_{k,j} = 1 \quad \text{for any } k$$

where φ_i is the projection onto $\mathbb{C}\Omega_j$. The corresponding C^* -algebras are [Toeplitz-Cuntz-Krieger algebras](#).

Arrays of matricially free Gaussian operators

$$\omega_{i,j}(u) = \sqrt{d_j}(\varphi_{i,j}(u) + \varphi_{i,j}^*(u))$$

play the role of **matricial semicircular operators**

$$[\omega(u)] = \begin{pmatrix} \omega_{1,1}(u) & \omega_{1,2}(u) & \dots & \omega_{1,r}(u) \\ \omega_{2,1}(u) & \omega_{2,2}(u) & \dots & \omega_{2,r}(u) \\ \cdot & \cdot & \ddots & \cdot \\ \omega_{r,1}(u) & \omega_{r,2}(u) & \dots & \omega_{r,r}(u) \end{pmatrix}$$

and generalize semicircular operators. From now on we incorporate scalars like $\sqrt{d_j}$ or more general $\sqrt{b_{i,j}(u)}$ in the operators.

Theorem [Voiculescu]

If $Y(u, n)$ are Hermitian standard Gaussian independent random matrices with complex entries, then

$$\lim_{n \rightarrow \infty} Y(u, n) = \omega(u)$$

in the sense of mixed moments under the complete trace $\tau(n)$.

Theorem

If $Y(u, n)$ are Hermitian Gaussian independent random matrices with **i.b.i.d.** complex entries, then

$$\lim_{n \rightarrow \infty} T_{i,j}(u, n) = \hat{\omega}_{i,j}(u)$$

in the sense of mixed moments under partial traces $\tau_j(n)$, where

$$\hat{\omega}_{i,j}(u) = \begin{cases} \omega_{j,j}(u) & \text{if } i = j \\ \omega_{i,j}(u) + \omega_{i,j}(u) & \text{if } i \neq j \end{cases}$$

are **symmetrized Gaussian operators**.

The symmetric blocks are called

- 1 **balanced** if $d_i > 0$ and $d_j > 0$
- 2 **unbalanced** if $d_i = 0 \wedge d_j > 0$ or $d_i > 0 \wedge d_j = 0$
- 3 **evanescent** if $d_i = 0$ and $d_j = 0$

where the numbers

$$\lim_{n \rightarrow \infty} \frac{|N_j|}{n} = d_j \geq 0$$

are called **asymptotic dimensions** .

Special cases

In the general formula for mixed moments, we get

- 1 $T_{i,j}(u, n) \rightarrow \hat{\omega}_{i,j}(u)$ if block is balanced
- 2 $T_{i,j}(u, n) \rightarrow \omega_{i,j}(u)$ if block is unbalanced, $j = 0 \wedge i > 0$
- 3 $T_{i,j}(u, n) \rightarrow \omega_{j,i}(u)$ if block is unbalanced, $j > 0 \wedge i = 0$
- 4 $T_{i,j}(u, n) \rightarrow 0$ if block is evanescent

Theorem [Voiculescu]

If $Y(u, n)$ are standard complex non-Hermitian Gaussian independent random matrices, then

$$\lim_{n \rightarrow \infty} Y(u, n) = \eta(u)$$

in the sense of moments under $\tau(n)$, where each $\eta(u)$ is circular.

Theorem

If $Y(u, n)$ are non-Hermitian Gaussian independent random matrices with i.b.i.d. complex entries, then

$$\lim_{n \rightarrow \infty} T_{i,j}(u, n) = \eta_{i,j}(u)$$

where

$$\eta_{i,j}(u) = \hat{\wp}_{i,j}(2u - 1) + \hat{\wp}_{i,j}^*(2u)$$

are called **matricial circular operators** and $\hat{\wp}_{i,j}(s)$ are symmetrizations of $\wp_{i,j}(s)$ (two partial isometries for each $\eta_{i,j}(u)$).

Lemma

We can identify partial isometries $\wp_{i,j}(u)$ with

$$\wp_{i,j}(u) = \ell(i,j,u) \otimes e(i,j) \in M_r(\mathcal{A})$$

where

- 1 the family $\{\ell(i,j,u) : i,j \in [r], u \in \mathcal{U}\}$ is a system of *-free creation operators w.r.t. state φ on \mathcal{A}
- 2 $(e_{i,j})$ is a system of matrix units in $M_r(\mathbb{C})$
- 3 $\Psi_j = \varphi \otimes \psi_j$, where ψ_j is the state associated with $e(j) \in \mathbb{C}^r$.

Lemma

Consequently,

$$\omega_{i,j}(u) = \ell(i,j,u) \otimes e(i,j) + \ell(i,j,u)^* \otimes e(j,i)$$

$$\widehat{\omega}_{i,j}(u) = \begin{cases} g(i,j,u) \otimes e(i,j) + g(i,j,u)^* \otimes e(j,i) & \text{if } i < j \\ f(i,u) \otimes e(i,i) & \text{if } i = j \end{cases}$$

where

$$\{g(i,j,u) : i,j \in [r], u \in \mathcal{U}\}, \quad \{f(i,u) : i \in [r], u \in \mathcal{U}\}$$

are $*$ -free generalized circular and semicircular systems, respectively.

Lemma

We can also identify matricial circular operators with

$$\eta_{i,j}(u) = \begin{cases} g(i,j,u) \otimes e(i,j) + g(j,i,u) \otimes e(j,i) & \text{if } i < j \\ g(i,i,u) \otimes e(i,i) & \text{if } i = j \end{cases}$$

Theorem

If $Y(u, n)$ are complex Gaussian independent random matrices with i.b.i.d. entries, then

$$\lim_{n \rightarrow \infty} S_{i,j}(u, n) = \zeta_{i,j}(u)$$

where

$$\zeta_{i,j}(u) = g(i, j, u) \otimes e(i, j)$$

are called **matricial R-circular operators** .

Consider one off-diagonal $\zeta_{i,j}$ of the form

$$\zeta_{i,j} = (l_1 + l_2^*) \otimes e(i,j)$$

where $i \neq j$ and l_1, l_2 are free creation operators with covariances γ_1 and γ_2 , respectively. We have

$$\Psi_j(\zeta_{i,j}^* \zeta_{i,j} \zeta_{i,j}^* \zeta_{i,j}) = \varphi(l_1^* l_1 l_1^* l_1) + \varphi(l_1^* l_2^* l_2 l_1) = \gamma_1^2 + \gamma_1 \gamma_2$$

whereas all remaining *-moments of $\zeta_{i,j}$ in the state Ψ_j vanish.

Theorem

The array of *-subalgebras $(\mathcal{M}_{i,j})$ of $M_r(\mathcal{A})$, each generated by

$$\{\ell(i,j,u) \otimes e(i,j) : u \in \mathcal{U}\},$$

with the unit

$$1_{i,j} = t(i,j) \otimes e(i,i) + 1 \otimes e(j,j)$$

where

$$t(i,j) = \begin{cases} \sum_{u \in \mathcal{U}} \ell(i,j,u) \ell(i,j,u)^* & \text{if } i \neq j \\ 0 & \text{if } i = j \end{cases}$$

is matricially free with respect to $(\Psi_{i,j})$, where $\Psi_{j,j} = \varphi \otimes \psi_j$.

Theorem

The array $(\mathcal{M}_{i,j})$ of $*$ -subalgebras of $M_r(\mathcal{A})$, each generated by

$$\{\ell(i,j,u) \otimes e(i,j), \ell(j,i,u) \otimes e(j,i) : u \in \mathcal{U}\}$$

and the symmetrized unit

$$1_{i,j} = 1 \otimes e(i,i) + 1 \otimes e(j,j),$$

is symmetrically matricially free with respect to $(\Psi_{i,j})$.

Corollary

The array $(\mathcal{M}_{i,j})$ of $*$ -subalgebras of $M_r(\mathcal{A})$ such that one of the following cases holds:

- 1 each $\mathcal{M}_{i,j}$ is generated by $\{\widehat{\omega}_{i,j}(u) : u \in \mathcal{U}\}$,
- 2 each $\mathcal{M}_{i,j}$ is generated by $\{\eta_{i,j}(u) : u \in \mathcal{U}\}$,
- 3 each $\mathcal{M}_{i,j}$ is generated by $\{\zeta_{i,j}(u) : u \in \mathcal{U}\}$,

is symmetrically matricially free with respect to $(\Psi_{i,j})$.

If $a_k = c_k \otimes e(i_k, j_k) \in M_r(\mathcal{A})$ for $k = 1, \dots, m$, where $c_k \in \mathcal{A}$, we will denote by

$$\mathcal{NC}_m(a_1, \dots, a_m)$$

the set of non-crossing partitions of $[m]$ which are **adapted** to $(e(i_1, j_1), \dots, e(i_m, j_m))$ which means that this tuple is cyclic

$$j_1 = i_2, j_2 = i_3, \dots, j_n = i_1$$

together with all tuples associated with the blocks of π . Its subset consisting of pair partitions will be denoted $\mathcal{NC}_m^2(a_1, \dots, a_m)$.

Example

For $(e(i, j), e(j, i), e(i, j), e(j, i))$, where $i \neq j$, there are three non-crossing partitions adapted to it:

$$\{\{1, 2, 3, 4\}\}, \{\{1, 4\}, \{2, 3\}\}, \{\{1, 2\}, \{3, 4\}\}.$$

In turn, the partitions

$$\{\{1, 2, 5\}, \{3, 4\}\}, \{\{1, 2, 3, 4, 5\}\} \quad \{\{1, 2, 3\}, \{4, 5\}\}$$

are the only non-crossing partitions adapted to the tuple $(e(i, j), e(j, k), e(k, i), e(i, m), e(m, i))$, where $i \neq j \neq k \neq i \neq m$.

Lemma

With the above notations, let $a_k = \zeta_{i_k, j_k}^{\epsilon_k}(u_k)$, where $i_k, j_k \in [r]$, $u_k \in \mathcal{U}$, and $\epsilon_k \in \{1, *\}$ for $j \in [m]$ and $m \in \mathbb{N}$. Then

$$\Psi_j(a_1 \dots a_m) = \sum_{\pi \in \mathcal{NC}_m^2(a_1, \dots, a_m)} b_j(\pi, f)$$

where j is equal to the second index of last matrix unit (associated with a_m) and f is the unique coloring of π adapted to (a_1, \dots, a_m) ,

$$b_j(\pi, f) = \prod_k b_j(\pi_k, f)$$

with $b_j(\pi_k) = b_{c(k), c(o(k))}(u)$, where $c(k)$, $c(o(k))$, j are colors assigned to π_k , its nearest outer block $\pi_{o(k)}$ and to the imaginary block. In the remaining cases, the moment vanishes.

Definition

A family of multilinear functions $\kappa_\pi[\cdot; j]$, where $j \in [r]$, of matricial variables $a_k = c_k \otimes e(i_k, j_k) \in M_r(\mathcal{A})$ is **cyclically multiplicative** over the blocks π_1, \dots, π_s of $\pi \in \mathcal{NC}_m(a_1, \dots, a_m)$ if

$$\kappa_\pi[a_1, \dots, a_m; q] = \prod_{k=1}^s \kappa(\pi_k)[a_1, \dots, a_m; j(k)],$$

for any a_1, \dots, a_m , where

$$\kappa(\pi_k)[a_1, \dots, a_m; j(k)] = \kappa_s(a_{q_1}, \dots, a_{q_s}; j_{q_s})$$

for the block $\pi_k = (q_1 < \dots < q_n)$, where $\{\kappa_s(\cdot; j) : s \geq 1, j \in [r]\}$ is a family of multilinear functions. If $\pi \notin \mathcal{NC}_m(a_1, \dots, a_m)$, we set $\kappa_\pi[a_1, \dots, a_m; j] = 0$ for any j .

Definition

By the **cyclic cumulants** we shall understand the family of multilinear cyclically multiplicative functionals over the blocks of non-crossing partitions

$$\pi \rightarrow \kappa_{\pi}[\cdot; j],$$

defined by r moment-cumulant formulas

$$\Psi_j(a_1 \dots a_m) = \sum_{\pi \in \mathcal{NC}_m(a_1, \dots, a_m)} \kappa_{\pi}[a_1, \dots, a_m; j],$$

where $j \in [r]$ and $\Psi_j = \varphi \otimes \psi_j$.

Take

$$\zeta_{i,j} = (\ell_1 + \ell_2^*) \otimes e(i,j) \quad \text{and} \quad \zeta_{i,j}^* = (\ell_1^* + \ell_2) \otimes e(j,i),$$

where $i \neq j$. The non-trivial second order cumulants are

$$\begin{aligned} \kappa_2(\zeta_{i,j}^*, \zeta_{i,j}; j) &= \Psi_j(\zeta_{i,j}^* \zeta_{i,j}) = \varphi(\ell_1^* \ell_1) = \gamma_1 \\ \kappa_2(\zeta_{i,j}, \zeta_{i,j}^*; i) &= \Psi_i(\zeta_{i,j} \zeta_{i,j}^*) = \varphi(\ell_2^* \ell_2) = \gamma_2 \end{aligned}$$

whereas all higher order cumulants vanish. For instance,

$$\begin{aligned} \kappa_2(\zeta_{i,j}^*, \zeta_{i,j}, \zeta_{i,j}^*, \zeta_{i,j}; j) &= \Psi_j(\zeta_{i,j}^* \zeta_{i,j} \zeta_{i,j}^* \zeta_{i,j}) \\ &\quad - \kappa_2(\zeta_{i,j}^*, \zeta_{i,j}; j) \kappa_2(\zeta_{i,j}^*, \zeta_{i,j}; j) \\ &\quad - \kappa_2(\zeta_{i,j}^*, \zeta_{i,j}; j) \kappa_2(\zeta_{i,j}, \zeta_{i,j}^*; i) \\ &= \gamma_1^2 + \gamma_1 \gamma_2 - \gamma_1^2 - \gamma_1 \gamma_2 = 0, \end{aligned}$$

Cyclic R-transforms

Let $\zeta := (\zeta_{i,j})$ and $\zeta^* := (\zeta_{i,j}^*)$. **Cyclic R-transforms** of this pair of arrays are formal power series in $z := (z_{i,j})$ and $z^* := (z_{i,j}^*)$ of the form

$$R_{\zeta, \zeta^*}(z, z^*; j) \\ = \sum_{n=1}^{\infty} \sum_{\substack{i_1, \dots, i_n \\ j_1, \dots, j_n}} \sum_{\epsilon_1, \dots, \epsilon_n} \kappa_n(\zeta_{i_1, j_1}^{\epsilon_1}, \dots, \zeta_{i_n, j_n}^{\epsilon_n}; j) z_{i_1, j_1}^{\epsilon_1} \dots z_{i_n, j_n}^{\epsilon_n},$$

where $i_1, \dots, i_n, j, j_1, \dots, j_n \in [r]$ and $\epsilon_1, \dots, \epsilon_n \in \{1, *\}$.

Theorem

If $\zeta := (\zeta_{i,j})$ is the square array of matricial R-circular operators and $\zeta^* := (\zeta_{i,j}^*)$, then its cyclic R-transforms are of the form

$$R_{\zeta, \zeta^*}(z, z^*; j) = \sum_{i=1}^r (b_{i,j} z_{i,j}^* z_{i,j} + b_{i,j} z_{j,i} z_{j,i}^*)$$

where $j \in [r]$, where $b_{i,j} = \kappa_2(\xi_{i,j}^*, \xi_{i,j}; j) = \kappa_2(\xi_{j,i}, \xi_{j,i}^*; j)$.

Theorem

If $\zeta := \zeta_{i,j}$, where $i \neq j$, then non-trivial cyclic R-transforms of the pair $\{\zeta, \zeta^*\}$ are of the form

$$\begin{aligned}R_{\zeta, \zeta^*}(z, z^*; j) &= b_{i,j} z_{i,j}^* z_{i,j} \\ R_{\zeta, \zeta^*}(z, z^*; i) &= b_{j,i} z_{i,j} z_{i,j}^*\end{aligned}$$

and if $\zeta := \zeta_{j,j}$, then

$$R_{\zeta, \zeta^*}(z, z^*; j) = b_{j,j}(z_{j,j}^* z_{j,j} + z_{j,j} z_{j,j}^*)$$

Corollary

If $b_{i,j} = d_i$ for any i, j , then $c := \sum_{i,j} \zeta_{i,j}$ is circular with respect to $\Psi = \sum_{j=1}^r d_j \Psi_j$ and the corresponding R-transform takes the form

$$R_{c,c^*}(z_1, z_2) = \sum_{j=1}^r d_j R_{\zeta,\zeta^*}(z, z^*; j) = z_1 z_2 + z_2 z_1,$$

where $R_{\zeta,\zeta^*}(z, z^*; j)$ are the cyclic R-transforms considered above in which all entries of arrays z and z^* are identified with z_1 and z_2 , respectively.