



ELSEVIER

Contents lists available at ScienceDirect

Journal of Theoretical Biology

journal homepage: www.elsevier.com/locate/jtbi

A mathematical model of intercellular signaling during epithelial wound healing[☆]

Filippo Posta^{a,*}, Tom Chou^{a,b,**}^a Department of Biomathematics, UCLA, Los Angeles, CA 90095-1766, USA^b Department of Mathematics, UCLA, Los Angeles, CA 90095-1555, USA

ARTICLE INFO

Article history:

Received 16 October 2009

Received in revised form

18 April 2010

Accepted 21 May 2010

Available online 2 June 2010

Keywords:

MAPK

ROS

EGF

Ligand

Traveling wave

ABSTRACT

Recent experiments monitoring the healing process of wounded epithelial monolayers have demonstrated the necessity of MAPK activation for coordinated cell movement after damage. This MAPK activity is characterized by two wave-like phenomena. One MAPK “wave” that originates immediately after injury, propagates deep into the cell sheet, away from the edge, and then rebounds back to the wound interface. After this initial MAPK activity has largely disappeared, a second MAPK front propagates slowly from the wound interface and also continues into the cell sheet, maintaining a sustained level of MAPK activity throughout the cell sheet. It has been suggested that the first wave is initiated by Reactive Oxygen Species (ROS) generated at the time of injury. In this work, we develop a minimal mathematical model that reproduces the observed behavior. The main ingredients of our model are a competition between ligand (e.g., Epithelial Growth Factor) and ROS for the activation of Epithelial Growth Factor Receptor, and a feedback loop between receptor occupancy and MAPK activation. We explore the mathematical properties of the model and look for traveling wave solutions consistent with the experimentally observed MAPK activity patterns.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Coordinated cell movement is an essential feature of many biological processes, such as wound healing, embryonic morphogenesis, and tumor growth (Martin and Parkhurst, 2004). In wound healing, cell migration and cell contraction are the two main mechanisms responsible for wound closure. Cell contraction is the dominant mechanism in the closing of small wounds through the so called “purse-string” process (Kiehart, 1999). For larger wounds, cell contraction is not sufficient, and surrounding cells must migrate to close larger wounds. While the two mechanisms are not mutually exclusive, there are cases where cell migration is the only healing process (Sherratt and Murray, 1990), such as when a strip of cells from an epithelial layer is removed (Poujade et al., 2007). Despite the existence of experimental assays targeting cell migration during wound healing, there are still many open questions. For example, what mechanical and biochemical phenomena regulate the switch from resting cells before injury to motile cells after injury? Is it the availability of free space that leads cells to move toward wound closure? What determines the speed of cell migration? To be able to

answer these questions, we need to understand both the mechanical and biochemical aspects of cell migration, and how they might regulate each other. While some of the physical mechanisms of cell movement have been studied (DiMilla et al., 1991; Sherratt and Murray, 1990; Murray, 2003; Maini et al., 2004), the complex regulation of the wound healing process by biochemical signals and feedback pathways remains poorly understood. Matsubayashi et al. monitored cell migration of monolayers of epithelial cells after wounding and observed two “waves” of phosphorylation of Mitogen Activated Protein Kinase (MAPK) (Matsubayashi et al., 2004). Within this context, a MAPK activation wave consists of a sequential increase of MAPK activity from cells at the wound edge to cells farther away from it (see Fig. 1). In more recent experiments, Nikolić et al. observed the same MAPK activation patterns under various wounding protocols, and identified Reactive Oxygen Species (ROS) as regulatory factors of MAPK activity during healing (Nikolić et al., 2006). A schematic of these experiments is shown in Fig. 1. Our goal is to build a mathematical representation of the intercellular signaling mechanisms observed in these two sets of experiments (Matsubayashi et al., 2004; Nikolić et al., 2006).

In general, MAPK activation is achieved through a cascade characterized by the sequential activation of three protein kinases (Ferrell, 1996). Because of its relevance and ubiquity, MAPK and its activation pathway remain the subject of many computational and mathematical modeling studies (Orton et al., 2005). Huang and Ferrell (1996) proposed a system of 18 differential equations

[☆] Funded by: NIH.

* Corresponding author.

** Corresponding author at: Department of Biomathematics, UCLA, Los Angeles, CA 90095-1766, USA.

E-mail addresses: fposta@ucla.edu (F. Posta), tomchou@ucla.edu (T. Chou).




Fig. 1. Schematic of MAPK activation during epithelial wound healing (Matsubayashi et al., 2004; Nikolić et al., 2006). The system extends ~ 3 mm in the z -direction normal to the cell monolayer. (a) The main elements of the experiments: a monolayer of epithelial cells; the cells have either MAPK activity (yellow) or not (no color filling). There are two diffusible signals responsible for MAPK activation, ROS (red circles) and EGF (blue squares). (b) At time $t=0$ a strip of cells is removed from the epithelial layer creating a wound. As a result of the injury, ROS molecules are released at wound edge (Nikolić et al., 2006). (c) ROS diffuses and promotes MAPK activation. This is the first MAPK wave observed in Matsubayashi et al. (2004) and Nikolić et al. (2006), it reaches its farthest point from the wound edge after 3 min. (d) The first wave “rebounds” back to wound edge as ROS generated by the wound is either consumed or diffuses away. At the same time the feedback loop between MAPK/EGFR/EGF promotes EGF release by MAPK-active epithelial cells (blue dots near wound edge). The “rebound” wave returns to the cells around the wound edge after 10 min from injury. (e) About 30 min after injury, the cells begin to move into the empty space left by the wound. Cell movement promotes ROS release. The combination of ROS production by migrating cells and EGF release by MAPK-active cells ignites the second sustained MAPK wave.

representing the 10 reactions that compose the three-kinases MAPK cascade. A mathematical and computational analysis of the system showed that the cascade has the effect of amplifying an input signal (e.g., receptor phosphorylation) in such a way that its overall behavior can be compared to that of a cooperative enzyme (Keener and Sneyd, 1998). These first computational studies sparked additional modeling efforts that, coupled with ongoing discoveries by experimentalists, have led to many more advanced models that exhibit characteristics (e.g., bistability, ultrasensitivity, oscillations, etc.) of the MAPK signaling pathway

(Qiao et al., 2007; Hornberg et al., 2005; Schoeberl et al., 2002; Sasagawa et al., 2005).

Here, we are not interested in the intracellular dynamics of MAPK, but rather in its role within the wound healing signaling network. For this reason, we will treat the MAPK cascade as a black box, using the results in Ferrell (1996, 1997) to essentially represent the whole cascade as a switch for signal transmission. In this approach, MAPK is the output of the “signaling switch”. ROS are one input that can activate the switch, but are unlikely to be the only one because of the different properties of the two activation waves. As suggested in Nikolić et al. (2006) and in other wound healing experiments (Xu et al., 2004; Block et al., 2004), other candidate inputs are diffusible ligands and their cell receptors. Epidermal Growth Factor (EGF) and EGF Receptor (EGFR) also play essential roles in promoting cell migration, proliferation and wound closure (Wiley et al., 2003; Joslin et al., 2007). Moreover, positive and negative feedbacks between EGFR signaling and the MAPK cascade have been demonstrated experimentally and verified computationally (Santos et al., 2007; Kholodenko, 2006, 2007). Although Nikolić et al. suggest EGF, as well as other molecules, as possible signals, they did not pursue the topic in their work. However, they did identify ROS as direct regulators of MAPK activity in their wound healing experiments. Since ROS have been shown to induce EGFR activation in the absence of EGF (Reynolds et al., 2003), this finding is in agreement with other studies that showed the regulatory role of ROS in wound healing (Roy et al., 2006; Sen and Roy, 2008) and MAPK signaling (Torres, 2003; McCubrey et al., 2006).

Because of the documented connection between diffusible signals, wound healing, and MAPK activation, we propose a mechanistic model based on ligand-mediated intercellular signaling. We use a continuum approach that is justified by the facts that the observed MAPK patterning spans many epithelial cells, and that the concentrations of the species relevant to MAPK activation are high (Pribyl et al., 2003b; Muratov et al., 2009). Because we are focusing on the signaling dynamics of the system, we do not model directly cell migration. Accordingly, we represent the positive feedback loop between MAPK signaling and cell migration through ROS production due to the mechanical stresses induced by migrating cells (Sen and Roy, 2008). As a result, the model reproduces the observed MAPK activation pattern, and is consistent with the qualitative experimental features observed in Matsubayashi et al. (2004) and Nikolić et al. (2006).

2. MAPK dynamics during wound healing

Before introducing the mathematical model, we present the experimental observations on which our model is based. In Matsubayashi et al. (2004) studied wound healing in epithelial monolayers of Madin–Darby canine kidney (MDCK) cells. They wounded the monolayers by removing a strip of cells from them, and then monitored cellular responses to the wound itself. The authors observed that, after injury, both the cells at wound margin and the submarginal cells migrate to close the wound. The number of cells remained essentially constant, an indication that cell proliferation did not play a role in this system. To investigate the cellular response after wounding, the authors stained the cells with ERK1/2 antibodies. ERK1/2 is a MAPK that had been shown to contribute to wound healing. The images showed a peculiar pattern of MAPK activation consisting of two “waves”: a fast transient one and a slow sustained one. Within this context, a MAPK activation wave consists of a sequential increase of MAPK activity from cells at the wound edge to cells farther away from it

(see Fig. 1). The rapid transient wave is generated at wound onset and spreads from marginal to submarginal cells. This wave reaches its peak (i.e., the furthest distance from the wound margin) after 3 min and quickly recedes back to the wound edge after about 10 min. During the retreat of the wave, MAPK activity disappears sequentially from cells farther away from the wound to cells at the wound edge. After the first wave, MAPK activation is confined to the wound margin until the onset of a second “wave”. The second MAPK activation wave starts 30 min after wounding and propagates from the marginal cells to the submarginal ones at a constant speed that is much slower (about two orders of magnitude) than the first wave. The authors observed that the second wave kept propagating for the duration of the experiments (4–6 h), and also that MAPK was inactive once the opposing edges of the wound confronted one another.

Matsubayashi et al. concluded that the observed MAPK activity is necessary for wound healing, but could not identify a specific mechanism or signal that contributes to MAPK activation. They did propose the existence of a feedback loop between cell movement and MAPK activation that promotes coordinated cell migration.

Nikolić et al. extended Matsubayashi et al.’s experiments by probing the epithelial wound healing assay with different wounding protocols to further elucidate the properties of MAPK activation (Nikolić et al., 2006). They observed that the two wave motifs are always present after mechanical injury. The kinematic description of the MAPK waves matches the one given by Matsubayashi et al., with the first wave reaching a maximum distance from the wound edge of about 480 μm after 3 min, then receding back to the wound edge until the onset of the second wave, 30 min after wounding, as depicted in Fig. 1. The authors were also able to detect ROS production at the wound site immediately following injury and investigated the role of ROS with respect to MAPK activation. They discovered that ROS is necessary for MAPK activation since both MAPK waves were absent during experiments in which ROS were inhibited. Nikolić et al. concluded that both MAPK waves are necessary for wound healing, and proposed a sequence of events to explain MAPK dynamics. This sequence starts with ROS production at the wound edge after mechanical wounding, ROS then ignites the first, fast, MAPK activation wave that promotes coordinated cell migration. The second MAPK wave and cell migration seem to be linked through a feedback loop, since they have the same kinematic properties, as also suggested by Matsubayashi et al.

3. Mathematical model

The experimental results from Matsubayashi et al. (2004) and Nikolić et al. (2006) provide evidence of precise spatio-temporal MAPK signaling for the regulation of cell migration during wound healing, without determining the exact biochemical events that govern it. Nikolić et al. identified ROS as signaling molecules responsible for the initial wave of MAPK, but they also stated that more diffusible signals are needed to explain the MAPK dynamics (Nikolić et al., 2006). As stated in the Introduction, we propose that two diffusible signals, ROS and EGF, are sufficient to explain the observed spatio-temporal MAPK pattern. In fact, ROS and EGF molecules are both able to diffuse in the extracellular space (ROS can also move across the cell membrane) and to phosphorylate the EGF membrane receptor (EGFR), that activates the MAPK cascade. EGF induces phosphorylation of the cytoplasmic tail of EGFR by binding to it. Reynolds et al. (2003) showed that ROS can also induce EGFR phosphorylation even in the absence of EGF by binding to intracellular phosphatases in the domain of the EGFR molecules. Also well documented is the positive feedback




Fig. 2. Schematic of biophysical events during wound healing. (a) Diffusible ligands (L) phosphorylate membrane receptors by binding to them, activating the MAPK cascade that leads to the production of intracellular protease (P). Protease induces ligand release. (b) ROS (S) can also interact with membrane receptors by inducing phosphorylation of their cytoplasmic tail, which also activates the MAPK cascade and promotes release of ligands into the extracellular matrix. Diffusible ligand and ROS represent two independent triggers of the MAPK cascade. (c) Schematic of the three-kinase cascade that characterizes MAPK signaling. Note that in our model we do not represent the cascade in details, but we replace it with a Hill function. (d) The stresses caused by cell movement toward wound closure lead to ROS release (Rhee et al., 2000; Torres, 2003; Ali et al., 2006).

between EGF and the MAPK cascade, and its ability to produce long range signaling through autocrine relays (Pribyl et al., 2003b). Finally, ROS can be generated by mechanical stresses like the ones generated by the migrating cells in the wounded epithelium (Torres, 2003), thus providing a feedback loop between MAPK activation, cell motility, and further ROS production. These four signaling mechanisms are summarized in Fig. 2 and constitute the main ingredients of our mathematical model.

The wound healing system is a three-dimensional one, and the migration of individual cells toward wound closure is not always normal to the wound edge as shown by cell tracking experiments (Nikolić et al., 2006). Here, we simplify the analysis by considering the cell layer in cross section as a semi-infinite straight line, with the wound initially positioned at the origin (cf. Fig. 2(d)). The resulting two-dimensional system consists of a semi-infinite cell layer that is immersed in medium of infinite height (the medium in Nikolić et al., 2006 is 3 mm thick which is much larger than any other length scale in the problem). We model four species: ROS, one diffusible ligand (e.g., EGF), one ligand receptor (e.g., EGFR), and playing the role of the output of the MAPK cascade “black box”, a protease that is the intracellular precursor of the ligand (e.g., the piece completing the feedback loop between EGF and the MAPK cascade in Figs. 2(a) and (b)). We denote the local concentrations of ROS, EGF, EGFR, and protease by L , S , R , and P , respectively. As depicted in Fig. 2, the signal can be transmitted to the cell in two different ways: through a ligand-receptor complex ($C_L \equiv R \cdot L$) and a ROS-receptor complex¹ ($C_S \equiv R \cdot S$). We will assume that the number of available cell membrane receptors is in excess, implying that R is approximately constant and that ROS-ligand-receptor complexes are negligible (Pribyl et al., 2003b; Muratov et al., 2009).

¹ ROS are known to activate EGFR by associating with its cytoplasmic tail, and inactivating its phosphatase activity. Kinetically, modeling this process is equivalent to irreversible ROS-EGFR ($C_S \equiv R \cdot S$) complex formation.




Fig. 3. Schematic of simplified signaling pathway during wound healing: ligands (L) and ROS (S) are free to diffuse and bind to cell surface receptors. This binding forms complexes (C_S and C_L) that activate MAPK signaling. Complexes are lost due to endocytosis or, in the case of ligand-receptor complexes, through ligand unbinding and endocytosis. Intracellular proteases (P) release extracellular ligands (L). Note that receptors are assumed to be in excess. Conversely, ligand-ROS-receptor complexes are negligible.

A schematic representation of the system is given in Fig. 3. The governing equations and boundary conditions of our model are

$$\frac{\partial L(X,Z,T)}{\partial T} = D_L \left(\frac{\partial^2 L(X,Z,T)}{\partial X^2} + \frac{\partial^2 L(X,Z,T)}{\partial Z^2} \right), \quad L(X,Z=\infty,T) = 0, \quad (1)$$

$$D_L \frac{\partial L(X,0,T)}{\partial Z} = k_{on}^L R L(X,0,T) - k_{off}^L C_L(X,T) - g_L P(X,T), \quad (2)$$

$$\frac{\partial C_L(X,T)}{\partial T} = k_{on}^L R L(X,0,T) - (k_{off}^L + k_{ec}^L) C_L(X,T), \quad (3)$$

$$\frac{\partial S(X,Z,T)}{\partial T} = D_S \left(\frac{\partial^2 S(X,Z,T)}{\partial X^2} + \frac{\partial^2 S(X,Z,T)}{\partial Z^2} \right), \quad S(X,Z=\infty,T) = 0, \quad (4)$$

$$D_S \frac{\partial S(X,0,T)}{\partial Z} = k_{on}^S R S(X,0,T) - g_S \Pi_S(P,X,T), \quad (5)$$

$$\frac{\partial C_S(X,T)}{\partial T} = k_{on}^S R S(X,0,T) - k_{ec}^S C_S(X,T), \quad (6)$$

$$\frac{\partial P(X,T)}{\partial T} = -k_P P(X,T) + g_P \Pi_P(C_L, C_S). \quad (7)$$

Eqs. (1) and (4) describe the diffusion of ligands and ROS in the extracellular medium with diffusion constants D_L and D_S , respectively. Eq. (2) accounts for the flux of ligand across the surface of the cellular layer consisting of ligand-receptor complex formation with rate constant k_{on}^L , ligand-receptor complex dissociation with rate constant k_{off}^L , and extracellular ligand release by intracellular protease with rate g_L . Eq. (3) governs the kinetics of ligand-receptor complexes, with new complexes forming at rate k_{on}^L and dissociating with rate k_{off}^L . The rate of receptor-mediated endocytosis of the ligand-bound receptor complexes is represented by k_{ec}^L . Eq. (5) describes the diffusive flux of ROS due to formation of ROS-receptor complexes with rate k_{on}^S and to ROS production by the functional $\Pi_S(C_L, X, T)$. The kinetics of ROS-receptor complexes is represented by Eq. (6) and its terms are analogous to the ones in Eq. (3), except that there is no ROS release from ROS-receptor complexes in accordance with the experimental results by Reynolds et al. (2003). The last equation describes the cellular response to extracellular signaling through the activity of intracellular proteases. Within the wound healing framework, protease activity is directly related to ERK1/2 activity measured in Nikolić et al. (2006). In particular, the protease dynamics is characterized by a degradation term with rate constant k_P and a source term $\Pi_P(C_L, C_S)$ with maximum production rate g_P . To complete the description of the

mathematical model, we need to impose reasonable functional forms for Π_P and Π_S .

The role of the functional $\Pi_P(C_L, C_S)$ is to represent the intermediate biochemical steps that lead to protease production. These steps include the MAPK cascade and any other reaction in the feedback loop between ligand binding and ligand release (e.g., the solid box in Fig. 2(c)). In the literature, Π_P is usually represented as a sigmoidal function of cell surface complexes such as the Hill function (Ferrell, 1996, 1997; Shvartsman et al., 2002). If the level of receptor signaling is given by the total concentration of complexes ($C_L + C_S$), we propose the following functional form:

$$\Pi_P(C_L, C_S) = \frac{(C_L + C_S)^n}{C_A^n + (C_L + C_S)^n}, \quad (8)$$

where n is an effective Hill coefficient, and C_A represents the activation threshold of the signaling pathway.

Defining the ROS source Π_S is more problematic since the experimental evidence suggests an interplay between cellular signaling and cell migration, thus involving mechanical forces whose description goes beyond the scope of this study. Conversely, the production of ROS due to wound induction is embedded in the initial conditions of the system and it is not described in Π_S . Generally, ROS production increases with cell metabolism (Torres, 2003). In our case, metabolic increase may be related to cells becoming motile (Ali et al., 2006; Torres, 2003) and/or to ligand signaling (Rhee et al., 2000). We use a Hill function multiplied by a decaying exponential to represent the spatially-dependent Π_S :

$$\Pi_S(C_L, X, T) = \begin{cases} 0 & T < T_D, \\ \frac{(P(X, T - T_D))^m}{S_A^m + (P(X, T - T_D))^m} e^{-k_s X^2} & T \geq T_D. \end{cases} \quad (9)$$

The Hill functional represents ROS production from motile cells after MAPK activation. The delay T_D represents the delay between MAPK activation and cell migration. The exponential factor in Eq. (9) describes reduction in ROS production due to the decrease in cell motility further from the wound edge at $X=0$. This choice of Π_S represents the feedback loop between MAPK signalling and cell deformation through its signaling pathways. We assume that cell motility is initiated by MAPK signaling through ligand and ROS binding. Then, motile cells release ROS as a byproduct of the mechanical stresses arising from cell motion.

3.1. Dimensionless model

Upon introducing the following dimensionless quantities

$$t = k_p T, \quad x = X \sqrt{\frac{k_p}{D_L}}, \quad z = Z \sqrt{\frac{k_p}{D_L}},$$

$$l = L \frac{k_p k_{ec}^L k_{on}^L R}{g_L g_P (k_{off}^L + k_{ec}^L)}, \quad c_L = C_L \frac{k_p k_{ec}^L}{g_L g_P}, \quad c_S = C_S \frac{k_{ec}^S}{g_S},$$

$$s = S \frac{k_{on}^S R}{g_S} \quad \text{and} \quad p = P \frac{k_p}{g_P},$$

we express the system of equations in dimensionless form:

$$\frac{\partial l(x, z, t)}{\partial t} = \frac{\partial^2 l(x, z, t)}{\partial x^2} + \frac{\partial^2 l(x, z, t)}{\partial z^2}, \quad (10)$$

$$\alpha l_z(x, 0, t) = l(x, 0, t) - \beta [l(x, 0, t) - c_L(x, t)] - p(x, t), \quad (11)$$

$$\varepsilon \frac{\partial c_L(x, t)}{\partial t} = l(x, 0, t) - c_L(x, t), \quad (12)$$

$$\delta \frac{\partial c_S(x, t)}{\partial t} = s(x, 0, t) - c_S(x, t), \quad (13)$$

$$\frac{\partial s(x, z, t)}{\partial t} = \eta \left(\frac{\partial^2 s(x, z, t)}{\partial x^2} + \frac{\partial^2 s(x, z, t)}{\partial z^2} \right), \quad (14)$$

$$v s_z(x, 0, t) = s(x, 0, t) - \pi_s(p, x, t), \quad (15)$$

$$\frac{\partial p}{\partial t} = -p + \pi_p(c_L, c_S), \quad (16)$$

where the dimensionless parameters are

$$\alpha = \frac{\sqrt{D_L k_p} (k_{off}^L + k_{ec}^L)}{k_{on}^L R k_{ec}^L}, \quad \beta = \frac{k_{off}^L}{k_{ec}^L}, \quad \delta = \frac{k_p}{k_{ec}^S},$$

$$\varepsilon = \frac{k_p}{k_{off}^L + k_{ec}^L}, \quad \eta = \frac{D_S}{D_L}, \quad v = \sqrt{\frac{k_p}{D_L}} \frac{D_S}{k_{on}^S R}. \quad (17)$$

The parameters α and v characterize the relative rates of diffusion and binding, while β represents the strength of complex degradation relative to ligand dissociation. The parameters ε and δ describe the speed of binding and endocytosis relative to ligand release mediated by intracellular species, and η is the ratio of diffusivity between ROS and EGF ligand.

The dimensionless protease and ROS production functions become

$$\pi_p(c_L, c_S) = \frac{(c_L + \gamma c_S)^n}{c_A^n + (c_L + \gamma c_S)^n}, \quad (18)$$

$$\pi_s(c_L, x, t) = \begin{cases} 0 & t < \tau, \\ \frac{(p(x, t - \tau))^m}{s_A^m + (p(x, t - \tau))^m} e^{-\lambda x^2} & t \geq \tau, \end{cases} \quad (19)$$

where $c_A = C_A (k_p k_{ec}^L) / (g_L g_P)$, $\gamma = (g_S k_p k_{ec}^L) / (g_L g_P k_{ec}^S)$, $\tau = T_D k_p$, $\lambda = k_x L$, and $s_A = (S_A k_p) / g_P$.

3.2. Fast-binding approximation

Before we proceed to the analysis of the model, we make an assumption that significantly simplifies the model and that is also justifiable biophysically (Pribyl et al., 2003b; Muratov et al., 2009). The range of values of k_p , k_{ec}^S , k_{off}^L , and k_{ec}^L given in Table 1 indicates that k_p is approximately one order of magnitude smaller than the other parameters (there is no value for k_{ec}^S in Table 1, but from the literature about ROS it is safe to assume that $k_{ec}^S > k_{ec}^L$ (Torres, 2003)). In our analysis we assume that ligand dissociation

Table 1

Typical values of model parameters.

Parameter	Typical value	Ref.
D_L	10^{-8} – 10^{-6} cm ² s ⁻¹	Pribyl et al. (2003b)
k_{on}^L	10^{-15} – 10^{-12} cm ³ s ⁻¹	Pribyl et al. (2003b)
k_{off}^L	10^{-3} – 10^{-2} s ⁻¹	Pribyl et al. (2003b)
k_{ec}^L	10^{-3} – 10^{-2} s ⁻¹	Pribyl et al. (2003b)
k_p	10^{-4} – 10^{-3} s ⁻¹	Pribyl et al. (2003b)
R	10^{10} – 10^{13} cm ⁻²	Pribyl et al. (2003b)
g_P	0.17×10^8 cm ⁻² s ⁻¹	Muratov et al. (2009)
g_L	0.54×10^{-2} s ⁻¹	Pribyl et al. (2003a)
C_A	10^9 cm ⁻²	Pribyl et al. (2003a)

and complex degradation is fast compared to protease degradation, e.g., $k_p \ll k_{ec}^S, k_{off}^L, k_{ec}^L$. In this limit, ε and δ are small and Eqs. (12) and (13) can be treated as a singular perturbation. On time scales of protease degradation the concentration of ligand-receptor and ROS-receptor complexes are approximately that of the surface concentration of free ligand and ROS, respectively. If we consider only the “outer” solutions of Eqs. (12) and (13), our full model reduces to the three equations:

$$\frac{\partial l(x, z, t)}{\partial t} = \frac{\partial^2 l(x, z, t)}{\partial x^2} + \frac{\partial^2 l(x, z, t)}{\partial z^2}, \quad (20)$$

$$\alpha l_z(x, 0, t) = l(x, 0, t) - p(x, t), \quad (21)$$

$$\frac{\partial s(x, z, t)}{\partial t} = \eta \left(\frac{\partial^2 s(x, z, t)}{\partial x^2} + \frac{\partial^2 s(x, z, t)}{\partial z^2} \right), \quad (22)$$

$$v s_z(x, 0, t) = s(x, 0, t) - \pi_s(p, x, t), \quad (23)$$

$$\frac{\partial p}{\partial t} = -p + \pi_p(l, s), \quad (24)$$

where all functions now represent “outer” solutions valid at times beyond initial transients in complex formation. In the Supplemental Information (SI), we show that the fast-binding approximation is accurate for the parameters used in our analysis. Explicit numerical solutions of the full set of dimensionless Eqs. (10)–(16) are compared with those derived from the fast-binding approximation (Eqs. (20)–(24)), leading to Table 1 in the SI.

4. Analysis and results

The spatio-temporal MAPK activation patterns can arise from different mechanisms. For example, one (or more) activation patterns could consist of a traveling front connecting two stable steady states, corresponding to high and low MAPK concentrations. In this section we describe the steady states of the system of Eqs. (20)–(24) and present an overview of the qualitative behavior of the solutions of the model. After establishing the dynamics of the mathematical model, we use the known model parameters to determine the nature of the MAPK patterns and some of their properties. Our analysis focuses on the model behavior at the cell layer level and neglects what happens to ROS and ligand concentrations in the bulk where molecules are simply diffusing.

4.1. Numerical solver

The complexity of our model requires analysis through numerical simulations. For this purpose we implemented a numerical solver in Fortran consisting of a centered finite difference scheme applied to a uniform discretization along the direction of the cell layer (e.g., x), and a staggered three-point

finite difference scheme applied to an optimal geometrical grid discretization (see Posta et al., 2008) along the direction normal to the cell layer (e.g., z). The equations are time-integrated using an explicit Euler discretization. This type of solver has been shown to maximize accuracy and minimize run-time when applied to problems that require an accurate resolution of the boundary data (e.g., species concentrations at cell layer level), but not of the data in the bulk (e.g., ROS and EGF concentrations in the extracellular matrix) (Posta et al., 2008). The boundary conditions of the problem consist of an homogeneous Dirichlet boundary condition at $z = \infty$, and homogeneous Neumann boundary conditions at $x = 0, \infty$. The instant of wound induction is taken as the zero of time, at which the ligand and protease concentrations are assumed to be zero throughout the domain. ROS concentration is also zero everywhere except around $x=0$, where $s(x=0, z=0, t=0) = \delta(0)$, with the Dirac delta-function approximated by a tall, thin Gaussian centered at $x=0$. The solver and its implementation are described in more detail in the SI.

4.2. Steady states & traveling fronts

Multistability is a common feature of reaction–diffusion systems similar to ours. For our model, all the steady states of the system, except the trivial one, will be spatially inhomogeneous because of the non-local nature of the problem. Therefore, by equilibrium solution of the system we mean solutions that are constant in space across the epithelial layer, but not along the normal direction. Within this context, the steady states of the model described by Eqs. (20)–(24) satisfy

$$\bar{l} = p, \quad \bar{s} = \pi_s(p, x, t), \quad p = \pi_p(\bar{l}, \bar{s}), \quad (25)$$

where the overbar indicates that the value of ligand or ROS concentration is taken at $z=0$ (e.g., $\bar{l} = l(x, z=0, t)$). In particular, the following condition needs to hold

$$\bar{l} = \pi_p(\bar{l}, \bar{s}). \quad (26)$$

This equation is always satisfied by the trivial solution ($\bar{l}=0, \bar{s}=0$), but under certain conditions it can have two more solutions, as highlighted in Fig. 4(a). In this case the three roots are two stable steady states, \bar{l}_0 and \bar{l}_2 , and an unstable one, \bar{l}_1 . The two stable steady states represent a state of no ligand signaling, $\bar{l}_0=0$, and a state of active ligand signaling, $\bar{l}_2 > 0$, respectively. From Fig. 4, we can also infer how ROS concentration \bar{s} and the activation threshold c_A control the steady states of the system. If we fix the activation threshold at a sufficiently high value, the

system attains only the trivial steady state, $\bar{l} = \bar{l}_0 = 0$, unless there is enough ROS to sustain the signaling pathway, as shown in Fig. 4(b). Conversely, if we fix \bar{s} , the system will be bistable only if the activation threshold c_A is sufficiently small (Fig. 4(c)).

Now that we have established the conditions for bistability, we wish to determine the existence of traveling waves connecting the two stable equilibria. Unfortunately, the model is too complicated to prove the existence of traveling waves unless we simplify it. We can prove the existence of traveling waves for a simpler scenario in which the concentration of ROS at cell layer level, \bar{s} , is constant and in the limit $n \rightarrow \infty$ for the Hill coefficient of π_p . In this limit, the sigmoidal protease production function π_p approaches a Heaviside function centered at $c_A - \bar{s}$:

$$\lim_{n \rightarrow \infty} \pi_p(\bar{l}, \bar{s}) = H(\bar{l} + \bar{s} - c_A) = \begin{cases} 1 & \bar{l} \geq c_A - \bar{s}, \\ 0 & \bar{l} < c_A - \bar{s}. \end{cases} \quad (27)$$

Under these conditions, the system is bistable for $0 < c_A - \bar{s} < 1$, and the roots of Eq. (26) are $\bar{l}_0 = 0$, $\bar{l}_1 = c_A - \bar{s}$, and $\bar{l}_2 = 1$. The existence of traveling fronts in the presence of bistability follows from the work of Cabre and Sola-Morales (2005). In addition, we can determine the velocity and direction of the traveling fronts by proceeding as in Muratov et al. (2009), obtaining

$$c_A - \bar{s} = \frac{1}{\pi} \int_{xv}^{\infty} \frac{\alpha \sqrt{q^2 - vq}}{q(1 + vq)(\alpha^2 q^2 - \alpha^2 vq + 1)} dq, \quad (28)$$

which gives an implicit relation for v , the velocity of the traveling wave for a fixed concentration of ROS. The integration variable q in the above integral arises from the Fourier transform used to derive Eq. (28). From Eq. (28), we find that the front velocity is a monotonically increasing function of the parameter α (see Eq. (17)). This result is expected since an increase in α corresponds to either an increase in ligand diffusivity or a decrease in ligand binding, and both changes result in the front reaching farther distances in a shorter amount of time. The direction of the front is determined by the threshold $c_A - \bar{s}$. If $c_A - \bar{s} < 1/2$, the front of active MAPK will move away from the wound, and deep into the cell layer. If $c_A - \bar{s} > 1/2$, MAPK activity will recede toward the wound. Regimes that delineate forward and backward traveling MAPK waves are indicated in Fig. 5. These results provide useful insight for the general case of $n < \infty$ and diffusing ROS. Numerical simulations show that for Hill coefficient as low as $n=6$, the instantaneous front velocity is within 10% of that obtained from Eq. (28). To summarize, we showed that the system can have two stable steady states and that traveling wave solutions connecting them are possible. In particular, ROS




Fig. 4. Steady state configurations. (a) Graphical solution of Eq. (26) for $\bar{s} = 0.1$, and $c_A = 0.7$. There are three solutions representing two stable steady states (\bar{l}_0 and \bar{l}_2) and one unstable equilibrium (\bar{l}_1). (b) If we fix the activation threshold ($c_A = 0.7$), different values of ROS concentration lead to different steady state configurations. Bistability is possible if there is enough ROS, otherwise the only stable steady state is the one with no MAPK activity. (c) We fix ROS concentration to $\bar{s} = 0.1$ and graphically solve Eq. (26) for different values of c_A . Bi-stability can arise only if the activation threshold is sufficiently small. The Hill coefficient $n=8$ was used in all plots.




Fig. 5. Traveling waves connecting the state of MAPK activation and the state of MAPK inactivity can move toward or away from the wound depending on ROS concentration at cell layer level (\bar{s}) and activation threshold (c_A) when $n \rightarrow \infty$. The thick black line indicates conditions under which the front is not moving. Regimes above the line lead to traveling waves moving toward the wound, while those below the line lead to fronts moving away from the wound.

concentration can determine the existence, velocity and direction of the fronts by effectively regulating the activation threshold of the MAPK cascade.

4.3. ROS/EGF regulation of MAPK activation

In this section we explore the parameter space of the wound healing assay to determine the nature of the MAPK fronts observed in Matsubayashi et al. (2004) and Nikolić et al. (2006). To avoid ambiguity, we divide the MAPK dynamics during wound healing into three wave-like events. The first event corresponds to the fast activation of MAPK initiated by the wound. It lasts until the activation front reaches its maximum depth in the cell layer. The second event is characterized by decrease of MAPK activity. It moves from deep into the epithelial monolayer toward the wound. These first two events qualitatively correspond to the first “rebounding wave” observed in the experiments (Matsubayashi et al., 2004; Nikolić et al., 2006). The last event consists of a slow activation front that starts near the wound edge and moves away from the wound. This last “wave” is initiated when the cells in the layer start moving to close the wound itself, and is sustained when the wound is large, preventing closure in finite time (Nikolić et al., 2006).

To reproduce the observed MAPK dynamics, we numerically integrated Eqs. (20)–(24) using the ligand related parameters given in Table 1. Although we could not find analogous references for physical parameters of ROS, we estimated parameter values from various sources. We used the self-diffusivity of water together with the Einstein relation to bound the value of η between 10 and 100. From the results in Reynolds et al. (2003), it seems reasonable to assume $k_{ec}^s \approx k_{ec}^l$. We assume that the initial concentration of ligand and protease is zero, while the concentration of ROS is represented by a narrow Gaussian with width equal to the size of a single cell; it represents the ROS initially released by cell rupture. Our numerical results are summarized in Figs. 6 and 7.




Fig. 6. Time evolution of the distance from the wound edge of the MAPK front. The circles and the intersecting vertical bars represent the average front position and one standard deviation error bars, respectively. Their values have been obtained from the experimental data in Nikolić et al. (2006). The solid trace has been obtained through a computer simulation of Eqs. (20)–(24) with dimensionless parameters: $\alpha = 0.4$, $\eta = 16$, $\nu = 5$, $c_A = 0.63$, $n = 9$, $\gamma = 2.2$, $p_A = 0.25$, $m = 9$, $\lambda = 0.25$, $\tau = 10$. These parameters are based on $k_p = 5.6 \times 10^{-3} \text{ s}^{-1}$, $D_L = 5.6 \times 10^{-7} \text{ cm}^2 \text{ s}^{-1}$, $k_{ec}^l = k_{off}^l = 10^{-2} \text{ s}^{-1}$, $g_L = 0.54 \times 10^{-2} \text{ s}^{-1}$, $g_p = 0.17 \times 10^8 \text{ cm}^{-2} \text{ s}^{-1}$, $R = 3.2 \times 10^{11} \text{ cm}^{-2}$, $k_{on}^l = 10^{-15} \text{ cm}^3 \text{ s}^{-1}$, $C_A = 10^9 \text{ cm}^{-2}$, $T_D = 1.8 \times 10^3 \text{ s}$.

Fig. 6 compares the time evolution of the distance of the front from the wound as predicted by Eqs. (20)–(24) with the experimental values observed in Nikolić et al. (2006) for scratch wounds. The position of the front is determined by evaluating the inflection point of protease concentration after each time step. From Fig. 6, we notice that the model does a good job to match the experimental data during increasing MAPK activity, but the model does not represent the decrease in MAPK activity as well. This result is due to the fact that the driving element of MAPK activation is the sharp functional π_p , and during MAPK activation the profile of $p(x,t)$ is also sharp (see Figs. 7(a) and (c)). Conversely, the decrease of MAPK activity corresponds to $\pi_p \sim 0$, and a dissipating concentration profile $p(x,t)$. For nearly flat density profiles (cf. the profile corresponding to 31 min in Fig. 7(b)), the inflection point is experimentally difficult to resolve and may appear to retract extremely fast even though the inflection point of $p(x,t)$ stays stationary as the gradients in $p(x,t)$ disappear. This effect may suggest a faster retraction in the wave front position than is predicted by the model (the ~ 5 –20 min regime in Fig. 6).

Another interesting feature of the MAPK front is the hump at ~ 40 –50 min. This feature is due to the time delay, τ , in Eq. (19) which retains a memory of the initial, fast wave (~ 0 –5 min). The experimental data is too sparse to reliably resolve this hump, if it actually occurs. More likely in reality, the delay in ROS generation, Eq. (19), should not be a sharp time translation, but rather a smoothed memory function described by an integral kernel.

Nonetheless, our overall model is able to qualitatively replicate the observed MAPK behavior. Fig. 7 shows the time evolution of the profiles of the three MAPK “waves”. The first wave (Fig. 7(a)) is driven by ROS production at the onset of wound and its fast diffusion. However, there is not enough ROS for either ligand or protease concentration to reach the signaling steady-state. As a result, the first “wave” can only propagate as far as $\sim 480 \mu\text{m}$ before receding. As ROS diffuses away, protease concentrations decrease (Fig. 7(b)) until the cells start to move (after about 30 min from injury). At that time, ROS is produced by the moving cells and fuels the positive feedback loop between ligand and protease. This feedback is strong enough to increase protease levels until they reach a signaling steady-state. At this point, the front moves like a traveling wave (Fig. 7(c)), with its speed and distance traveled regulated by the ROS source function π_s . We can also track the time-evolution of the variables in the model. Fig. 7(d) shows how protease




Fig. 7. Time evolution of protease. (a) After injury the fast diffusion of ROS drives MAPK activation. (b) As ROS diffuses away, the positive feedback loop between MAPK and ligand is not strong enough to sustain signaling and the depth of the front decreases. (c) Once the cells start moving, ROS production and the ligand–protease feedback loop fuel the activation front. (d) Time evolution of protease concentration at $x=0$. The parameters used to obtain these plots are the same as the ones used to generate Fig. 6.

concentration at the wound edge, $x=0$, changes in time. From this graph we observe that during the first MAPK event the protease concentration never reaches the “signaling” steady state and eventually decreases. During the third wave, protease concentration reaches the “signaling” steady state and remains there as shown by the flat part of the graph in Fig. 7(d). Note the dip around $T=60$ min due to the time delay in Π_S .

To summarize, only the third MAPK front behaves as a traveling wave, while the initial two events (corresponding to the first “rebounding wave” observed in experiments) are actually transient, diffusion-driven patterns.

5. Conclusions

We have formulated a mathematical model for the dynamics of intercellular signaling observed during wound healing experiments. From this model, we were able to replicate the signaling patterns observed in Matsubayashi et al. (2004) and Nikolić et al. (2006), and to provide mechanistic insight regarding their nature. From our results, the cooperative behavior between ROS and a diffusible ligand is necessary to explain the observed MAPK activation patterns during epithelial wound healing. Our choice of EGF as the signaling ligand is based upon our literature review, but lacks experimental evidence within the epithelial wound healing assay. However, we showed that the properties of EGF (and EGFR) fit the profile for the unidentified diffusible signals mentioned in Nikolić et al. (2006). Our model can be expanded to incorporate other diffusible signaling molecules. Although it may be possible to find physiologically realistic sets of parameters that lead to signaling patterns consisting of three separate traveling waves, the parameters associated with the EGF/ROS/MAPK system lead to only one final traveling wave. The first two notable events (initial MAPK wave and its “rebound”) are described by purely diffusive and decaying dynamics, respectively.

An aspect of our current model that needs improvement and further analysis is the determination of the ROS source function Π_S . From the experiments in Nikolić et al. (2006), this term seems to be negligible since extracellular ROS was detected only up to

~ 10 min after wounding. However, there is substantial evidence indicating that cell motility and EGFR-driven MAPK activation can induce ROS production. A plausible explanation for these conflicting results could be that ROS produced after wounding is fully recaptured by intracellular processes (including EGFR phosphorylation) and never crosses the cell membrane. Nonetheless, we performed many numerical tests and found that if $\Pi_S=0$ (data not shown), then all three MAPK events are diffusion driven and the signaling pattern is due to the different diffusion properties of ROS and ligand (EGF).

Currently, we represent the positive feedback between MAPK signaling and cell deformation through the molecular species responsible for MAPK activation. A more physically realistic approach would involve the modeling of the forces transmitted by adjoining cells that are contracting and extending while moving toward wound closure, then couple these forces with a biochemical signal (e.g., ROS) that can activate MAPK. There are many obstacles in implementing such model. Some of the cells at the wound edge develop lamellipodia and lead the migration, while the majority of the remaining cells seem to move through a tug-of-war mechanism (Trepate et al., 2009). It is also very difficult to experimentally quantify the mechanical forces responsible for cell migration. Recently, Trepate et al. (2009) were able to obtain direct measurements of cell traction and stress in a spreading, uninjured, epithelial layer. Additional complications arise from the multi-scale and three-dimensional nature of the problem, as well as stochastic effects due to the tug-of-war mechanism (Trepate et al., 2009). Nonetheless, our simple model could be extended by replacing π_S with a more realistic mechano–chemical functional.

Acknowledgments

We thank M. Gibbons, S. Shvartsman, and C. Muratov for useful discussion. This work was supported by NSF Grant DMS-0349195 and NIH Grant K25AI058672.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version at doi:[10.1016/j.jtbi.2010.05.029](https://doi.org/10.1016/j.jtbi.2010.05.029).

References

- Ali, M.H., Mungai, P.T., Schmacker, P.T., 2006. Stretch-induced phosphorylation of focal adhesion kinase in endothelial cells: role of mitochondrial oxidants. *Am. J. Physiol. Lung Cell Mol. Physiol.* 291, 38–45.
- Block, E.R., Matella, A.R., SundarRaj, N., Iszkula, E.R., Klarlund, J.K., 2004. Wounding induces motility in sheets of corneal epithelial cells through loss of spatial constraints. *J. Biol. Chem.* 279, 24307–24312.
- Cabre, X., Sola-Morales, J., 2005. Layer solutions on a half-space for boundary reactions. *Comm. Pure Appl. Math.* 1.VIII, 1678–1732.
- DiMilla, P.A., Barbee, K., Lauffenburger, D.A., 1991. Mathematical model for the effects of adhesion and mechanics on cell migration speed. *Biophys. J.* 84, 2907–2918.
- Ferrell, J.E., 1996. Tripping the switch fantastic: how a protein kinase cascade can convert graded inputs into switch-like outputs. *Trends Biochem. Sci.* 12, 460–466.
- Ferrell, J.E., 1997. How responses get more switch-like as you move down a protein kinase cascade. *Trends Biochem. Sci.* 8, 288–289.
- Hornberg, J.J., Binder, B., Bruggeman, F.J., Schoeberl, B., Heinrich, R., Westerhoff, H.V., 2005. Control of MAPK signalling: from complexity to what really matters. *Oncogene* 24, 5533–5542.
- Huang, C.F., Ferrell, J.E., 1996. Ultrasensitivity in the mitogen-activated protein kinase cascade. *PNAS* 93, 10078–10083.
- Joslin, E.J., Opreko, L.K., Wells, A., Wiley, H.S., Lauffenburger, D.A., 2007. EGF-receptor-mediated mammary epithelial cell migration is driven by sustained ERK signaling from autocrine stimulation. *J. Cell Sci.* 120, 3688–3699.
- Keener, J., Sneyd, J., 1998. *Mathematical Physiology*. Springer, Berlin.
- Kholodenko, B.N., 2006. Cell signalling dynamics in time and space. *Nat. Rev. Mol. Cell Biol.* 7, 165–176.
- Kholodenko, B.N., 2007. Untangling the signalling wires. *Nature Cell Biol.* 9, 247–249.
- Kiehart, D.P., 1999. Wound healing: the power of the purse string. *Curr. Biol.* 9, R602–R605.
- Maini, P.K., McElwain, D.L.S., Leavesley, D., 2004. Travelling waves in a wound healing assay. *Appl. Math. Lett.* 17, 575–580.
- Martin, P., Parkhurst, S.M., 2004. Parallels between tissue repair and embryo morphogenesis. *Development* 131, 3021–3034.
- Matsubayashi, Y., Ebisuya, M., Honjoh, S., Nishida, E., 2004. ERK activation propagates in epithelial cell sheets and regulates their migration during wound healing. *Curr. Biol.* 14, 731–735.
- McCubrey, J.A., LaHair, M.M., Franklin, R.A., 2006. Reactive oxygen species-induced activation of the MAP kinase signaling pathways. *Antioxid. Redox Sign.* 8, 1775–1793.
- Muratov, C.B., Posta, F., Shvartsman, S.Y., 2009. Autocrine signal transmission with extracellular ligand degradation. *Phys. Biol.* 6, 13.
- Murray, J.D., 2003. *Mathematical Biology II: Spatial Models and Biomedical Applications*. Springer, Berlin.
- Nikolić, D.L., Boettiger, A.N., Barbeck, J.D., Shvartsman, S.Y., 2006. The role of boundary conditions in an experimental model of epithelial wound healing. *AJP-Cell Physiol.* 291 (1), C68–C75.
- Orton, R.J., Sturm, O.E., Vyshemirky, V., Calder, M., Gilbert, D.R., Kolch, W., 2005. Computational modelling of the receptor-tyrosine-kinase-activated MAPK pathway. *Biochem. J.* 392, 249–261.
- Posta, F., Shvartsman, S.Y., Muratov, C.B., 2008. Compensated optimal grids for elliptic boundary-value problems. *J. Comput. Phys.* 227, 8622–8635.
- Poujade, M., Grasland-Mongrain, E., Hertzog, A., Jouanneau, J., Chavier, P., Ladoux, B., Buguin, A., Silberzan, P., 2007. Collective migration of an epithelial monolayer in response to a model wound. *PNAS* 104, 15988–15993.
- Pribyl, M., Muratov, C.B., Shvartsman, S.Y., 2003a. Discrete models of autocrine cell communication in epithelial layers. *Biophys. J.* 84, 3624–3635.
- Pribyl, M., Muratov, C.B., Shvartsman, S.Y., 2003b. Long-range signal transmission in autocrine relays. *Biophys. J.* 84, 883–896.
- Qiao, L., Nachbar, R.B., Kevrekidis, I.G., Shvartsman, S.Y., 2007. Bistability and oscillations in the Huang–Ferrell model of MAPK signaling. *PLoS Comput. Biol.* 3, 1819–1826.
- Reynolds, A.R., Tishcer, C., Vermeer, P.J., Rocks, O., Bastiaens, P.I.H., 2003. EGFR activation coupled to inhibition of tyrosine phosphatases causes lateral signal propagation. *Nature Cell Biol.* 5, 447–453.
- Rhee, S., Bae, Y., Lee, S., Kwon, J., 2000. Hydrogen peroxide: a key messenger that modulates protein phosphorylation through cysteine oxidation. *Sci. STKE* 53, pe1–pe4.
- Roy, S., Khanna, S., Nallu, K., Hunt, T.K., Sen, C.K., 2006. Dermal wound healing is subject to redox control. *Mol. Ther.* 13, 211–220.
- Santos, S.D.M., Vermeer, P.J., Bastiaens, P.I.H., 2007. Growth factor-induced MAPK network topology shapes erk response determining PC-12 cell fate. *Nature Cell Biol.* 9, 324–330.
- Sasagawa, S., Ozaki, Y., Fujita, K., Kuroda, S., 2005. Prediction and validation of the distinct dynamics of transient and sustained ERK activation. *Nat. Cell Biol.* 7, 365–373.
- Schoeberl, B., Eichler-Jonsson, C., Gilles, E.D., Müller, G., 2002. Computational modeling of the dynamics of the MAP kinase cascade activated by surface and internalized EGF receptors. *Nat. Biotechnol.* 20, 370–375.
- Sen, C.K., Roy, S., 2008. Redox signals in wound healing. *Biochim. Biophys. Acta* 1780, 1348–1361.
- Sherratt, J.A., Murray, J.D., 1990. Models of epidermal wound healing. *Proc. R. Soc. London* 241, 29–36.
- Shvartsman, S.Y., Muratov, C.B., Lauffenburger, D.A., 2002. Modeling and computational analysis of EGF receptor-mediated cell communication in *Drosophila* oogenesis. *Development* 129, 2577–2589.
- Torres, M., 2003. Mitogen-activated protein kinase pathways in redox signaling. *Front. Biosci.* 8, 369–391.
- Trepat, X., Wasserman, M.R., Angelini, T.E., Millet, E., Weitz, D.A., 2009. Physical forces during collective cell migration. *Nat. Phys.* 5, 426–430.
- Wiley, H.S., Shvartsman, S.Y., Lauffenburger, D.A., 2003. Computational modeling of the EGF-receptor system: a paradigm for systems biology. *TRENDS Cell Biol.* 13, 43–50.
- Xu, K., Ding, Y., Ling, J., Dong, Z., Yu, F.X., 2004. Wound-induced HB-EGF ectodomain shedding and EGFR activation in corneal epithelial cells. *Inv. Ophthalmol. Vis. Sci.* 45, 813–820.