

A GUIDE TO TENSOR-TRIANGULAR CLASSIFICATION

PAUL BALMER

ABSTRACT. This is a chapter of the Handbook of Homotopy Theory, that surveys the classifications of thick tensor-ideals.

1. INTRODUCTION

Stable homotopy theory shines across pure mathematics, from topology to analysis, from algebra to geometry. While its liturgy invokes Quillen model structures and ∞ -categories, profane users around the world often speak the vernacular of *triangulated categories*, as we shall do in this chapter.

Perhaps the first salient fact about stable homotopy categories is that in almost all cases they turn out to be *wild* categories – beyond the trivial examples of course. Dade famously began his paper [Dad78] with the admonition “There are just too many modules over p-groups!” and this truth resonates in all other fields as well: no hope to classify topological spaces up to stable homotopy equivalence; no more hope with complexes of sheaves, nor with equivariant C^* -algebras, nor with motives, etc, etc. One might dream that things improve with ‘small’ objects (compact, rigid, or else) but the problem persists even there: Stable homotopy theory is just too complicated!

Faced with the complexity of stable homotopy categories, we are led to the following paradigm shift. A classification *up to isomorphism* makes sense in any category, i.e. as soon as we can speak of *isomorphism*. But stable homotopy categories are more than mere categories: They carry additional structures, starting with the *triangulation*. In the case of a *tensor-triangulated category* (*tt-category* for short), as we consider in this chapter, we have two basic tools at hand: triangles and tensor. Instead of ignoring these additional structures, we should include them in the concept of

tt-classification

which is our nickname for *classification up to the tensor-triangular structure*.

More precisely, we want to decide when two objects X and Y can be obtained from one another by using tensor with anything, direct sums, summands, cones, suspension, etc. In mathematical terms, we ask when X and Y generate the same thick triangulated tensor-ideals. Heuristically, if you can build Y out of X by using the tt-structure then X contains at least as much information as Y . If you can go back and forth between X and Y , then they contain the same amount of information.

Date: 2019 December 17.

2010 *Mathematics Subject Classification.* 18F99, 55P42, 55U35.

Key words and phrases. tensor-triangulated category, spectrum, classification.

Research supported by NSF grant DMS-1600032.

The remarkable gain is that the tt-classification of an essentially small (rigid) tensor-triangulated category can *always* be achieved by means of a geometric object, more precisely a spectral topological space, called its *tensor-triangular spectrum*.⁽¹⁾ Let us highlight this starting point:

Fundamental fact: *Although almost every symmetric monoidal stable homotopy category \mathcal{K} is ‘wild’ as a category, we always have a tt-classification of its objects, via a topological space, $\mathrm{Spc}(\mathcal{K})$, called the spectrum of \mathcal{K} .*⁽²⁾

This chapter is dedicated to a survey of tt-classifications across different examples, as far as they are known to the author at this point in time.

The original idea of classifying objects up to the ambient structure was born in topology, around Ravenel’s conjectures [Rav84] and the ‘chromatic’ theorems of Devinatz-Hopkins-Smith [DHS88, HS98]; this relied on Morava’s work, among many other contributions. The ground-breaking insight of transposing from topology to other fields began with Hopkins [Hop87]. It is arguably Thomason [Tho97] who first understood how essential the tensor was in the global story. We recall in Remark 4.6 why such a *geometric* classification does not exist for mere triangulated categories, i.e. without the tensor.

The tt-spectrum was introduced in [Bal05] and is reviewed in Section 2. The survey begins in Section 3, with the initial example of topological stable homotopy theory. Section 4 touches commutative algebra and algebraic geometry. Section 5 is dedicated to stable module categories in modular representation theory and beyond. Section 6 discusses equivariant stable homotopy theory and Kasparov’s equivariant KK -theory. Section 7 pertains to motives and \mathbb{A}^1 -homotopy theory.

Everywhere, we have tried to give some idea of the actual tt-categories which come into play. When the amount of specialized definitions appears too high for this chapter, we simply point to the bibliographical references.

Finally, let us say a word about the bigger picture. In commutative algebra, the Zariski spectrum is not meant to be explicitly computed for every single commutative ring in the universe; instead, it serves as a stepping stone towards the geometric reasonings of algebraic geometry. In the same spirit, the tt-spectrum opens up a world of mathematical investigation, called *tensor-triangular geometry*, which reaches far beyond classical algebraic geometry into the broad kingdom of stable homotopy theory. The short final Section 8 points to further reading in that direction.

Acknowledgements: I would like to thank Tobias Barthel, Ivo Dell’Ambrogio, Martin Gallauer, Beren Sanders and Greg Stevenson for their help in assembling this survey. I apologize to anyone whose tt-geometric results are not mentioned here: For the sake of pithiness, I chose to restrict myself to the topic of tt-classification.

2. THE TT-SPECTRUM AND THE CLASSIFICATION OF TT-IDEALS

2.1. *Definition.* A *tt-category*, short for *tensor-triangulated category*, is a triangulated category \mathcal{K} together with a symmetric monoidal structure

$$\otimes: \mathcal{K} \times \mathcal{K} \longrightarrow \mathcal{K}$$

¹ Our use of the word ‘spectrum’ comes from commutative algebra, as in the ‘Zariski spectrum’, and should not be confused with the suspension-inverting ‘spectra’ of topology.

² $\mathrm{Spc}(\mathcal{K})$ is a space in the universe containing the ‘set’ of isomorphism classes of \mathcal{K} .

which is exact in each variable. See details in [HPS97, App. A] or [Nee01, KN02]. The \otimes -unit is denoted $\mathbb{1}$.

2.2. *Assumption.* Unless otherwise stated, we always assume that \mathcal{K} is *essentially small*, i.e. has a set of isomorphism classes of objects. Subcategories $\mathcal{J} \subseteq \mathcal{K}$ are always assumed full and replete (i.e. closed under isomorphisms).

2.3. *Definition.* A *triangulated* subcategory $\mathcal{J} \subseteq \mathcal{K}$ is a non-empty subcategory such that whenever $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$ is an exact triangle in \mathcal{K} and two out of X, Y and Z belong to \mathcal{J} then so does the third. A *thick* subcategory $\mathcal{J} \subseteq \mathcal{K}$ is a triangulated subcategory closed under direct summands: if $X \oplus Y \in \mathcal{J}$ then $X, Y \in \mathcal{J}$. A *tt-ideal* $\mathcal{J} \subseteq \mathcal{K}$, short for *thick tensor-ideal*, is a thick subcategory closed under tensoring with any object: $\mathcal{K} \otimes \mathcal{J} \subseteq \mathcal{J}$. A tt-ideal $\mathcal{J} \subseteq \mathcal{K}$ is called *radical* if $X^{\otimes n} \in \mathcal{J}$ for $n \geq 2$ forces $X \in \mathcal{J}$.

2.4. *Remark.* When every object of \mathcal{K} is rigid (i.e. admits a dual, a.k.a. *strongly dualizable* [HPS97, §2.1]), then we say that \mathcal{K} is *rigid* and we can show that every tt-ideal \mathcal{J} is automatically radical. See [Bal07, Prop. 2.4]. So, for simplicity, we assume that every tt-category \mathcal{K} that we discuss below is either rigid or that the phrase ‘tt-ideal’ means ‘radical tt-ideal’.

2.5. *Notation.* For a class $\mathcal{E} \subseteq \mathcal{K}$ of objects, the tt-ideal generated by \mathcal{E} is $\langle \mathcal{E} \rangle = \bigcap_{\mathcal{J} \supseteq \mathcal{E}} \mathcal{J}$, where \mathcal{J} runs through the tt-ideals containing \mathcal{E} .

2.6. *Definition.* A *prime* $\mathcal{P} \subset \mathcal{K}$ is a proper tt-ideal such that $X \otimes Y \in \mathcal{P}$ forces $X \in \mathcal{P}$ or $Y \in \mathcal{P}$. We denote the set of prime tt-ideals by

$$\mathrm{Spc}(\mathcal{K}) = \{ \mathcal{P} \subset \mathcal{K} \mid \mathcal{P} \text{ is prime} \}$$

and call it the *spectrum* of \mathcal{K} . The *support* of an object $X \in \mathcal{K}$ is the subset

$$\mathrm{supp}(X) = \{ \mathcal{P} \in \mathrm{Spc}(\mathcal{K}) \mid X \notin \mathcal{P} \}.$$

The topology of $\mathrm{Spc}(\mathcal{K})$ is defined to have $\{\mathrm{supp}(X)\}_{X \in \mathcal{K}}$ as basis of closed subsets. Explicitly, for each set of objects $\mathcal{E} \subseteq \mathcal{K}$ the subset $U(\mathcal{E}) = \{ \mathcal{P} \in \mathrm{Spc}(\mathcal{K}) \mid \mathcal{E} \cap \mathcal{P} \neq \emptyset \}$ is an open of $\mathrm{Spc}(\mathcal{K})$, and all open subsets are of this form, for some \mathcal{E} .

2.7. *Remark.* The above construction is introduced in [Bal05], where the pair $(\mathrm{Spc}(\mathcal{K}), \mathrm{supp})$ is characterized by a universal property: It is the *final support data*. See [Bal05, Thm. 3.2]. We shall not make this explicit but intuitively it means that the space $\mathrm{Spc}(\mathcal{K})$ is the best possible one carrying *closed* supports for objects of \mathcal{K} with the following rules for all X, Y, Z in \mathcal{K} :

- (1) $\mathrm{supp}(0)$ is empty and $\mathrm{supp}(\mathbb{1})$ is the whole space;
- (2) $\mathrm{supp}(X \oplus Y) = \mathrm{supp}(X) \cup \mathrm{supp}(Y)$;
- (3) $\mathrm{supp}(\Sigma X) = \mathrm{supp}(X)$;
- (4) $\mathrm{supp}(Z) \subseteq \mathrm{supp}(X) \cup \mathrm{supp}(Y)$ for each exact triangle $X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$;
- (5) $\mathrm{supp}(X \otimes Y) = \mathrm{supp}(X) \cap \mathrm{supp}(Y)$.

2.8. *Example.* Dually to the Zariski topology, in the tt-spectrum $\mathrm{Spc}(\mathcal{K})$ the closure of a point $\mathcal{P} \in \mathrm{Spc}(\mathcal{K})$ consists of all the primes *contained* in it: $\overline{\{\mathcal{P}\}} = \{ \mathcal{Q} \in \mathrm{Spc}(\mathcal{K}) \mid \mathcal{Q} \subseteq \mathcal{P} \}$. See [Bal05, Prop. 2.9].

2.9. *Remark.* The tt-spectrum $\mathrm{Spc}(\mathcal{K})$ is always a *spectral space* in the sense of Hochster [Hoc69]: it is quasi-compact, it admits a basis of quasi-compact open subsets, and each of its non-empty irreducible closed subsets has a unique generic point. See [Bal05, §2].

2.10. *Remark.* The construction $\mathcal{K} \mapsto \mathrm{Spc}(\mathcal{K})$ is a contravariant functor. Every exact \otimes -functor $F: \mathcal{K} \rightarrow \mathcal{K}'$ between tt-categories induces a continuous (spectral) map $\varphi = \mathrm{Spc}(F): \mathrm{Spc}(\mathcal{K}') \rightarrow \mathrm{Spc}(\mathcal{K})$ defined by $\mathcal{Q} \mapsto F^{-1}(\mathcal{Q})$. It satisfies $\varphi^{-1}(\mathrm{supp}(X)) = \mathrm{supp}(F(X))$ for all $X \in \mathcal{K}$. See [Bal05, § 3].

To express tt-classification via the spectrum, we need some preparation.

2.11. *Definition.* To every subset $V \subseteq \mathrm{Spc}(\mathcal{K})$ we can associate a tt-ideal

$$\mathcal{K}_V = \{ X \in \mathcal{K} \mid \mathrm{supp}(X) \subseteq V \}$$

of \mathcal{K} . (In fact, this tt-ideal is always radical. See Remark 2.4.)

2.12. *Definition.* A subset $V \subseteq \mathrm{Spc}(\mathcal{K})$ is called a *Thomason subset* if it is the union of the complements of a collection of quasi-compact open subsets: $V = \cup_{\alpha} V_{\alpha}$ where each V_{α} is closed with quasi-compact complement. In the terminology of Hochster [Hoc69], these are the *dual-open* subsets.

2.13. *Example.* If the space $\mathrm{Spc}(\mathcal{K})$ is topologically noetherian (i.e. all open subsets are quasi-compact), then V being Thomason is just being closed under specialization ($x \in V \Rightarrow \overline{\{x\}} \subseteq V$), i.e. being a union of closed subsets.

2.14. **Theorem** (Classification of tt-ideals, [Bal05, Thm. 4.10]). *The assignment $V \mapsto \mathcal{K}_V$ of Definition 2.11 defines an order-preserving bijection between the Thomason subsets $V \subseteq \mathrm{Spc}(\mathcal{K})$ and the (radical) tt-ideals $\mathcal{J} \subseteq \mathcal{K}$ of \mathcal{K} , whose inverse is given by $\mathcal{J} \mapsto \mathrm{supp}(\mathcal{J}) := \cup_{X \in \mathcal{J}} \mathrm{supp}(X) = \{ \mathcal{P} \mid \mathcal{J} \not\subseteq \mathcal{P} \}$.*

Specifically for the tt-classification of objects $X, Y \in \mathcal{K}$ (see Remark 2.4):

2.15. **Corollary.** *Two objects $X, Y \in \mathcal{K}$ generate the same tt-ideals $\langle X \rangle = \langle Y \rangle$ if and only if they have the same support $\mathrm{supp}(X) = \mathrm{supp}(Y)$. More precisely, Y belongs to $\langle X \rangle$ if and only if $\mathrm{supp}(Y) \subseteq \mathrm{supp}(X)$.*

The following converse to Theorem 2.14 holds. See [Bal05] for details.

2.16. **Theorem** (Balmer/Buan-Krause-Solberg). *Suppose that a spectral space \mathcal{S} carries a support data $\sigma(X) \subseteq \mathcal{S}$ for $X \in \mathcal{K}$ in the sense of [Bal05] and suppose that the assignment $\mathcal{S} \supseteq V \mapsto \{ X \in \mathcal{K} \mid \sigma(X) \subseteq V \}$ induces a bijection between Thomason subsets V of \mathcal{S} and (radical) tt-ideals of \mathcal{K} . Then the canonical map $\mathcal{S} \rightarrow \mathrm{Spc}(\mathcal{K})$ of Remark 2.7 is a homeomorphism.*

This result was established in [Bal05] under the additional assumption that \mathcal{S} be noetherian. It was proved in the above maximal generality in [BKS07]. (See Remark 2.9.)

2.17. *Remark.* Theorems 2.14 and 2.16 allow for a compact reformulation of tt-classifications, including the ones anterior to [Bal05]. Thus most classifications for the tt-categories \mathcal{K} discussed in Sections 3-7 are phrased in the simple form of a description of $\mathrm{Spc}(\mathcal{K})$. The tt-classification is then always the same, in terms of subsets of $\mathrm{Spc}(\mathcal{K})$, as in Theorems 2.14 and Corollary 2.15, and we shall not repeat these corollaries.

On the other hand, approaching tt-classification via $\mathrm{Spc}(\mathcal{K})$ buys us some flexibility, for partial results about $\mathrm{Spc}(\mathcal{K})$ can be interesting while a ‘partial classification’ is an odd concept. For instance, one can know $\mathrm{Spc}(\mathcal{K})$ as a set in some examples, with partial information on the topology. Or one can describe $\mathrm{Spc}(\mathcal{K}) = U \cup Z$ with

a complete description of the closed subset Z and its open complement U without knowing exactly how they attach. And so on.

In recent years, the geometric study of the tt-spectrum *per se* has led to new computations of $\mathrm{Spc}(\mathcal{K})$, from which the tt-classification can be deduced a posteriori. This will be illustrated in the later sections.

2.18. *Remark.* Some of the above results connect to lattice theory, see [BKS07, KP17]. It is a non-trivial property of a lattice, like that of tt-ideals in \mathcal{K} , to be *spatial*, i.e. in bijection with the open subsets of a topological space. In fact, without the tensor this fails in general (Remark 4.6).

The tt-classification of Theorem 2.14 tacitly assumes that \mathcal{K} consists of ‘small enough’ objects. Assumption 2.2 and Remark 2.4 belong to this logic too. Another indication of the smallness of \mathcal{K} is that we do not mention infinite coproducts in \mathcal{K} , and we only discuss thick subcategories, not localizing ones (i.e. those closed under arbitrary coproducts). When dealing with a ‘big’ tt-category \mathcal{T} , the natural candidate for a ‘small’ \mathcal{K} is the subcategory of rigid objects in \mathcal{T} , which may or may not coincide with compact ones.

There are also ‘big’ subcategories of ‘big’ tt-categories worth investigating, most famously *smashing* subcategories. It is an open problem whether the lattice of smashing \otimes -ideals is spatial or not. We prove in [BKS17] that it is a *frame*, thus it is at least ‘spatial’ in the quirky sense of pointless topology.

The connection between thick subcategories of compact objects and smashing subcategories is a topic in its own right, often dubbed the *Telescope Conjecture*. We shall not attempt to discuss it systematically here but will mention it in a few examples. See Krause [Kra00] for a beautiful abstract answer via ideals of morphisms.

3. TOPOLOGY

As already said, tt-classification (or at least ‘t-classification’) was born in topology, more precisely in *chromatic homotopy theory*, see [BB19]. The tt-category we consider here is the topological stable homotopy category SH , i.e. the homotopy category of topological spectra, and more specifically its subcategory SH^c of compact objects. See for instance [Rav92]. In other words, SH^c is the Spanier-Whitehead stable homotopy category of finite pointed CW-complexes.

The first operation one can do on SH is to p -localize it at a prime p , i.e. invert multiplication by every prime different from p . On compacts, this gives us $\mathrm{SH}_{(p)}^c$. Both SH^c and $\mathrm{SH}_{(p)}^c$ are essentially small rigid tt-categories.

3.1. *Remark.* Something special happens in SH^c and therefore in $\mathrm{SH}_{(p)}^c$ as well: The unit $\mathbb{1} = S^0$, a.k.a. the sphere spectrum, generates the category as a thick triangulated subcategory. Consequently, every thick subcategory is automatically a tt-ideal. In such situations, the tensor is not essential in the tt-classification and we are equivalently classifying thick subcategories.

3.2. *Remark.* A critical ingredient in chromatic theory is the countable family of so-called *Morava K-theories*, which are homology theories $K_{p,n}$, for $n \geq 1$, defined on $\mathrm{SH}_{(p)}^c$ and taking values in graded modules over the ‘graded field’ $\mathbb{F}_p[v_n, v_n^{-1}]$, with v_n in degree $2(p^n - 1)$. See [Rav92, § 1.5].

3.3. Theorem (Hopkins-Smith [HS98]). *The spectrum of the classical stable homotopy category SH^c is the following topological space:*

$$\mathrm{Spc}(\mathrm{SH}^c) = \begin{array}{ccccccc} & \mathcal{P}_{2,\infty} & \mathcal{P}_{3,\infty} & \cdots & \mathcal{P}_{p,\infty} & \cdots & \\ & \vdots & \vdots & & \vdots & & \\ & | & | & & | & & \\ \mathcal{P}_{2,n+1} & & \mathcal{P}_{3,n+1} & \cdots & \mathcal{P}_{p,n+1} & \cdots & \\ | & & | & & | & & \\ \mathcal{P}_{2,n} & & \mathcal{P}_{3,n} & \cdots & \mathcal{P}_{p,n} & \cdots & \\ & \vdots & \vdots & & \vdots & & \\ & | & | & & | & & \\ \mathcal{P}_{2,2} & & \mathcal{P}_{3,2} & \cdots & \mathcal{P}_{p,2} & \cdots & \\ & & \searrow & & \swarrow & & \\ & & & \mathcal{P}_{0,1} & & & \end{array}$$

in which every line indicates that the higher point belongs to the closure of the lower one (Example 2.8). More precisely:

- (a) The *tt*-prime $\mathcal{P}_{0,1}$ is the kernel of rationalization $\mathrm{SH}^c \rightarrow \mathrm{SH}_{\mathbb{Q}}^c \cong \mathrm{D}^b(\mathbb{Q})$, that is, the subcategory of torsion spectra. It is the dense point of $\mathrm{Spc}(\mathrm{SH}^c)$.
- (b) For each prime number p , the *tt*-prime $\mathcal{P}_{p,\infty}$ is the kernel of localization $\mathrm{SH}^c \rightarrow \mathrm{SH}_{(p)}^c$. These $\mathcal{P}_{p,\infty}$ are exactly the closed points of $\mathrm{Spc}(\mathrm{SH}^c)$.
- (c) For each prime number p and each integer $2 \leq n < \infty$, the *tt*-prime $\mathcal{P}_{p,n}$ is the kernel of the composite $\mathrm{SH}^c \rightarrow \mathrm{SH}_{(p)}^c \rightarrow \mathbb{F}_p[v_n^{\pm 1}]$ -grmod of localization at p and $(n-1)^{\mathrm{st}}$ Morava K -theory $K_{p,n-1}$ (Remark 3.2).
- (d) The support of an object X is either empty when $X = 0$, or the whole of $\mathrm{Spc}(\mathrm{SH}^c)$ when X is non-torsion, or a finite union of ‘columns’

$$\overline{\{\mathcal{P}_{p,m_p}\}} = \{\mathcal{P}_{p,n} \mid m_p \leq n \leq \infty\}$$

for integers $2 \leq m_p < \infty$ varying with the primes p .

- (e) A closed subset is either empty, or the whole $\mathrm{Spc}(\mathrm{SH}^c)$, or a finite union of closed points $\{\mathcal{P}_{p,\infty}\}$ and of columns $\overline{\{\mathcal{P}_{p,m_p}\}}$ with $m_p \geq 2$ as in (d).
- (f) A Thomason subset of $\mathrm{Spc}(\mathrm{SH}^c)$ is either empty, or the whole $\mathrm{Spc}(\mathrm{SH}^c)$, or an arbitrary union of columns $\overline{\{\mathcal{P}_{p,m_p}\}}$ with $m_p \geq 2$ as in (d).

The above Theorem 3.3 is not the way the chromatic filtration is expressed in the original literature; see the translation in [Bal10a, §9].

3.4. Example. An object $X \in \mathrm{SH}^c$ has support contained in the p -th column, $\mathrm{supp}(X) \subseteq \overline{\{\mathcal{P}_{p,2}\}}$, if and only if it is ‘ p -primary torsion’, i.e. it satisfies $p^\ell \cdot X = 0$ for some $\ell \geq 1$.

3.5. Example. The support of the *tt*-ideal $\mathcal{J} = \mathcal{P}_{0,1}$ of torsion spectra is exactly the Thomason subset $\mathrm{Spc}(\mathrm{SH}^c) \setminus \{\mathcal{P}_{0,1}\}$ and is therefore the disjoint union of all columns $\sqcup_p \overline{\{\mathcal{P}_{p,2}\}}$. This reflects the fact that a torsion object in SH^c is the direct sum of p -primary torsion objects as in Example 3.4.

3.6. Remark. The fact that the closed point $\{\mathcal{P}_{p,\infty}\}$ cannot be the support of an object reflects the fact that an object in $\mathrm{SH}_{(p)}^c$ which is killed by all Morava K -theories $K_{p,n}$ for $n \geq 1$ must be zero. It also shows that $\mathrm{Spc}(\mathcal{K})$ is not noetherian, already in this initial case of $\mathcal{K} = \mathrm{SH}^c$ (see Example 2.13).

3.7. *Remark.* In this setting, the Telescope Conjecture is open (again). See [Kra00] and further references therein.

4. COMMUTATIVE ALGEBRA AND ALGEBRAIC GEOMETRY

As already indicated, Hopkins [Hop87] initiated the transposition of the chromatic classification from topology to algebra. The correct statement for noetherian rings was proved by Neeman [Nee92] and the perfect version for general schemes, not necessarily noetherian, is due to Thomason in his last published paper [Tho97]. In terms of tt-spectra it becomes the following very beautiful result.

4.1. **Theorem** (Thomason [Tho97]). *Let \mathcal{X} be a scheme which is quasi-compact and quasi-separated. Then the spectrum of the derived category $D^{\text{perf}}(\mathcal{X})$ of perfect complexes (with $\otimes = {}^L\otimes_{\mathcal{O}_{\mathcal{X}}}$) is isomorphic to the underlying space $|\mathcal{X}|$ itself, via the homeomorphism*

$$\begin{array}{ccc} |\mathcal{X}| & \xrightarrow{\cong} & \text{Spc}(D^{\text{perf}}(\mathcal{X})) \\ x & \longmapsto & \mathcal{P}(x) \end{array}$$

where, for each point x of \mathcal{X} , the tt-prime $\mathcal{P}(x) = \{Y \in D^{\text{perf}}(\mathcal{X}) \mid Y_x \cong 0\}$ is the kernel of localization $D^{\text{perf}}(\mathcal{X}) \rightarrow D^{\text{perf}}(\mathcal{O}_{\mathcal{X},x})$ at x .

4.2. *Remark.* Equivalently, $\mathcal{P}(x)$ can be described as the kernel of the residue functor $D^{\text{perf}}(\mathcal{X}) \rightarrow D^{\text{b}}(\kappa(x))$ to the residue field $\kappa(x)$ of \mathcal{X} at x .

4.3. *Remark.* Recall that a scheme \mathcal{X} is quasi-compact and quasi-separated if the underlying space $|\mathcal{X}|$ admits a basis of quasi-compact open subsets (including $|\mathcal{X}|$ itself). This purely topological condition is equivalent to $|\mathcal{X}|$ being spectral. Hence this condition is the maximal generality in which the above result can hold in view of Remark 2.9. Noetherian schemes and affine schemes are quasi-compact and quasi-separated.

The affine case vindicates our use of the word ‘spectrum’⁽³⁾:

4.4. **Corollary.** *Let A be a commutative ring. Then the tt-spectrum of the homotopy category $K^{\text{b}}(A\text{-proj}) \cong D^{\text{perf}}(A)$ of bounded complexes of finitely generated projective A -modules is homeomorphic to the Zariski spectrum of A*

$$\text{Spec}(A) \xrightarrow{\sim} \text{Spc}(K^{\text{b}}(A\text{-proj})).$$

4.5. *Remark.* An error in [Hop87], corrected in [Nee92], was not to assume A noetherian. However we see that Thomason’s Corollary 4.4 does not assume A noetherian. The point is that the tt-classification (Theorem 2.14) which is equivalent to Corollary 4.4 involves actual Thomason subsets not mere specialization-closed subsets, whereas [Hop87] and [Nee92] are phrased in terms of specialization-closed subsets. The assumption that A is noetherian is only useful to replace ‘Thomason’ by ‘specialization-closed’ (Example 2.13).

4.6. *Remark.* As we saw in the topological example of Section 3, when the unit $\mathbb{1}$ generates the tt-category \mathcal{K} as a thick subcategory we do not really need the tensor. This is also the case for $\mathcal{K} = K^{\text{b}}(A\text{-proj})$ for instance.

³If this creates confusion with Example 2.8, note that the map of Corollary 4.4, $\mathfrak{p} \mapsto \mathcal{P}(\mathfrak{p}) = \text{Ker}(D^{\text{perf}}(A) \rightarrow D^{\text{perf}}(A_{\mathfrak{p}}))$, reverses inclusions: $\mathfrak{p} \subseteq \mathfrak{q} \Rightarrow \mathcal{P}(\mathfrak{p}) \supseteq \mathcal{P}(\mathfrak{q})$.

But in general the tensor is essential for classification by means of subsets of $\mathrm{Spc}(\mathcal{K})$. Indeed, the lattice of thick subcategories of a triangulated category \mathcal{K} cannot be classified in terms of the lattice of subsets of pretty much anything because it may not satisfy *distributivity*: $\mathcal{J}_1 \wedge (\mathcal{J}_2 \vee \mathcal{J}_3) = (\mathcal{J}_1 \wedge \mathcal{J}_2) \vee (\mathcal{J}_1 \wedge \mathcal{J}_3)$. Already for $\mathcal{K} = \mathrm{D}^{\mathrm{perf}}(\mathcal{X})$ over the projective line $\mathcal{X} = \mathbb{P}_k^1$ distributivity fails with \mathcal{J}_i the thick subcategory generated by $\mathcal{O}(i)$. See [BKS17, Rem. 5.10].

4.7. *Remark.* An application of Theorem 4.1 is the reconstruction of every quasi-compact and quasi-separated scheme \mathcal{X} from the data of the *tensor*-triangulated category $\mathrm{D}^{\mathrm{perf}}(\mathcal{X})$. Indeed, one can equip the tt-spectrum $\mathrm{Spc}(\mathcal{K})$ with a sheaf of commutative rings, which in the case of $\mathcal{K} = \mathrm{D}^{\mathrm{perf}}(\mathcal{X})$ recovers the structure sheaf $\mathcal{O}_{\mathcal{X}}$. See details in [Bal05, §6]. By contrast, Mukai [Muk81] proved earlier that such a reconstruction is impossible from the triangulated structure alone.

4.8. *Remark.* The above $\mathcal{K} = \mathrm{D}^{\mathrm{perf}}(R)$ are the compact and rigid objects in the big derived category $\mathcal{T} = \mathrm{D}(R)$. The Telescope Conjecture holds when R is noetherian by Neeman [Nee92] but fails in general by Keller [Kel94].

4.9. *Remark.* Other tt-categories can be associated to schemes, or commutative rings, for instance right-bounded derived categories. For first results in this direction, see work of Matsui and Takahashi [MT17, Mat18].

One can generalize Theorem 4.1 almost verbatim to reasonable stacks:

4.10. **Theorem** (Hall [Hal16, Thm. 1.2]). *Let \mathcal{X} be a quasi-compact algebraic stack with quasi-finite separated diagonal, whose stabilizer groups at geometric points are finite linearly reductive group schemes (\mathcal{X} is ‘tame’). Then $\mathrm{Spc}(\mathrm{D}^{\mathrm{perf}}(\mathcal{X})) \cong |\mathcal{X}|$.*

We refer to [Hal16] for terminology. Note earlier work of Krishna [Kri09] in characteristic 0, and of Dubey-Mallick [DM12] for finite groups acting on smooth schemes in characteristic prime to the order of the groups.

One can also consider the graded version of Corollary 4.4:

4.11. **Theorem** (Dell’Ambrogio-Stevenson [DS14, Thm. 4.7]). *Let A be a graded-commutative ring (graded over any abelian group), then there is a canonical isomorphism $\mathrm{Spc}(\mathrm{D}^{\mathrm{perf}}(A)) \cong \mathrm{Spec}^{\mathrm{h}}(A)$, between the tt-spectrum of $\mathrm{D}^{\mathrm{perf}}(A)$ and the spectrum of homogeneous prime ideals of A .*

Let us mention a variation relating to singularities.

4.12. **Theorem** (Stevenson [Ste14, Thm. 7.7]). *Let \mathcal{X} be a noetherian separated scheme with only hypersurface singularities. Then there is an order-preserving bijection between the specialization-closed subsets of the singular locus of \mathcal{X} and the thick $\mathrm{D}^{\mathrm{perf}}(\mathcal{X})$ -submodules of the singularity category $\mathrm{D}^{\mathrm{b}}(\mathrm{coh} \mathcal{X}) / \mathrm{D}^{\mathrm{perf}}(\mathcal{X})$.*

Here the singularity category is not itself a tt-category but a triangulated category with an action by the tt-category $\mathrm{D}^{\mathrm{perf}}(\mathcal{X})$. As such, this result is an application of Stevenson’s *relative tt-geometry* [Ste13]. Another application of Stevenson’s theory is the tt-classification for derived categories of matrix factorizations in Hirano [Hir19], which extends earlier result of Takahashi [Tak10].

5. MODULAR REPRESENTATION THEORY AND RELATED TOPICS

5.1. *Point.* Let G be a finite group and let k be a field. Maschke's Theorem says that the order of G is invertible in k if and only if kG is semisimple. In that case, all kG -modules are projective. *Modular* representation theory refers to the non-semisimple situation. Then the *stable module category* is the additive quotient [Hap88]

$$kG\text{-stmod} = \frac{kG\text{-mod}}{kG\text{-proj}}$$

which precisely measures how far kG is from being semisimple. It is a tt-category whose objects are all finitely generated kG -modules and whose groups of morphisms $\text{Hom}_{kG\text{-stmod}}(X, Y)$ are given by the quotient of the abelian group of kG -linear maps $\text{Hom}_{kG}(X, Y)$ modulo the subgroup of those maps factoring via a projective module. Tensor is over k with diagonal G -action: $g \cdot (x \otimes y) = (gx) \otimes (gy)$ in $X \otimes_k Y$. The \otimes -unit is $\mathbb{1} = k$ with trivial G -action.

5.2. *Point.* We can also consider the derived category $D^b(kG\text{-mod})$, with the 'same' tensor. Every non-zero tt-ideal $\mathcal{J} \subseteq D^b(kG\text{-mod})$ contains $D^{\text{perf}}(kG)$ because $kG \otimes - \cong \text{Ind}_1^G \text{Res}_1^G$ and $D^b(k\text{-mod}) = D^{\text{perf}}(k)$ is semisimple. Hence the tt-classification of $D^b(kG\text{-mod})$ and of its Verdier quotient by $D^{\text{perf}}(kG)$ are very close. (The former has just one more tt-ideal: zero.) By Rickard [Ric89], that quotient is equivalent to the stable module category:

$$\frac{D^b(kG\text{-mod})}{D^{\text{perf}}(kG)} \cong kG\text{-stmod}.$$

5.3. **Theorem** (Benson-Carlson-Rickard [BCR97]). *There is a homeomorphism between the spectrum of the stable module category and the so-called projective support variety*

$$\text{Spc}(kG\text{-stmod}) \cong \text{Proj}(\mathbf{H}^\bullet(G, k))$$

which can be extended (by adding one closed point) to a homeomorphism

$$\text{Spc}(D^b(kG\text{-mod})) \cong \text{Spec}^h(\mathbf{H}^\bullet(G, k)).$$

Explicitly, to every homogeneous prime $\mathfrak{p}^\bullet \subset \mathbf{H}^\bullet(G, k)$ corresponds the tt-prime

$$\mathcal{P}(\mathfrak{p}^\bullet) = \{ X \mid \text{there is a homogeneous } \zeta \notin \mathfrak{p}^\bullet \text{ such that } \zeta \cdot X = 0 \}.$$

Again, the above does not appear verbatim in the source. See details in [Bal05] or [Bal10a, Prop. 8.5]. A more recent proof can be found in [CI15].

5.4. *Remark.* The reader interested in the related derived category of cochains on the classifying space BG is referred to [BIK11] for finite groups and to the comprehensive recent work [BCHV19] for p -local compact groups; see comments and references therein about earlier work by Benson-Greenlees.

5.5. *Point.* For finite group schemes G , the following generalization of Theorem 5.3 would follow from claims made in [FP07] but a flaw was found in [FP07, Thm. 5.3], which was eventually fixed in the recent [BIKP18]; see in particular [BIKP18, Rem. 5.4 and Thm. 10.3].

5.6. **Theorem** (Benson-Friedlander-Iyengar-Krause-Pevtsova). *For a finite group scheme G over k , there is a homeomorphism $\text{Spc}(kG\text{-stmod}) \cong \text{Proj}(\mathbf{H}^\bullet(G, k))$.*

5.7. *Remark.* Stable module categories of finite group schemes over a field are very ‘noetherian’ and several other results are known about the ‘big’ stable module category as well, like the Telescope Conjecture. See details in [BIKP18]. The technique of *stratification* has led to the tt-classification (of small and large subcategories) in several ‘noetherian enough’ derived settings. See the survey in [BIK12] and further references in [BIKP18].

5.8. *Point.* Extending beyond field coefficients to other rings R , we can consider the *relative stable module category* $RG\text{-strel}$, obtained from the Frobenius exact structure on the exact category of finitely generated RG -modules with R -split exact sequences. Already in small Krull dimension, interesting phenomena can be observed, as in the following result.

5.9. **Theorem** (Baland-Chirvasitu-Stevenson [BCS19, Thm. 1.1]). *Let S be a discrete valuation ring having residue field k and uniformizing parameter t and let $R_n = S/t^n$. Let G be a finite group. Then the tt-spectrum of the relative stable module category*

$$\mathrm{Spc}(R_n G\text{-strel}) \cong \sqcup_{i=1}^n \mathrm{Spc}(kG\text{-stmod}),$$

is a coproduct of n copies of the projective support variety of Theorem 5.3.

On the topic of singularity categories, let us mention [Xu14] and its recent generalization (recall that a category is EI if any endomorphism is invertible):

5.10. **Theorem** (Wang [Wan19, Thm. 5.2]). *Let \mathcal{C} be a finite EI category, projective over a field k and $D_{\mathrm{sing}}(k\mathcal{C}) = D^{\mathrm{b}}(k\mathcal{C}\text{-mod})/D^{\mathrm{b}}(k\mathcal{C}\text{-proj})$ its singularity category. Then there is a homeomorphism*

$$\mathrm{Spc}(D_{\mathrm{sing}}(k\mathcal{C})) \cong \sqcup_{x \in \mathcal{C}} \mathrm{Spc}(kG_x\text{-stmod}),$$

where $G_x = \mathrm{Aut}_{\mathcal{C}}(x)$.

5.11. *Point.* Antieau-Stevenson [AS16] consider further derived categories of representations of small categories over commutative noetherian rings. They obtain several interesting classifications, including for localizing subcategories, and in particular for simply laced Dynkin quivers. See also the earlier [LS13].

5.12. *Point.* Let us now turn our attention to stable module categories related to Lie algebras. Boe-Kujawa-Nakano [BKN17b] prove several results about classical Lie superalgebras. In particular for the general linear Lie superalgebra $\mathfrak{g} = \mathfrak{g}(m|n) = \mathfrak{g}_{\bar{0}} \oplus \mathfrak{g}_{\bar{1}}$ and $\mathcal{K} = \underline{\mathcal{F}}$ the stable category of the category \mathcal{F} of finite dimensional \mathfrak{g} -modules which admit a compatible action by $G_{\bar{0}}$ and are completely reducible as $G_{\bar{0}}$ -modules (where $\mathrm{Lie} G_{\bar{0}} = \mathfrak{g}_{\bar{0}}$). They prove in [BKN17b, Thm. 5.2.2] that the spectrum $\mathrm{Spc}(\underline{\mathcal{F}})$ is homeomorphic to the N -homogenous spectrum $N\text{-Proj}(S^{\bullet}(\mathfrak{f}_{\bar{1}}))$ where \mathfrak{f} is the detecting subalgebra of \mathfrak{g} and $N = \mathrm{Norm}_{G_{\bar{0}}}(\mathfrak{f}_{\bar{1}})$.

The same authors more recently considered quantum groups:

5.13. **Theorem** (Boe-Kujawa-Nakano [BKN17a, Thm. 7.6.1]). *Let G be a complex simple algebraic group over \mathbb{C} with $\mathfrak{g} = \mathrm{Lie} G$. Assume that ζ is a primitive ℓ th root of unity where ℓ is greater than the Coxeter number for \mathfrak{g} . Then the tt-spectrum of the stable module category for the quantum group $U_{\zeta}(\mathfrak{g})$ is*

$$\mathrm{Spc}(U_{\zeta}(\mathfrak{g})\text{-stmod}) \cong G\text{-Proj}(\mathbb{C}[\mathcal{N}])$$

where \mathcal{N} is the nullcone, i.e. the set of nilpotent elements of \mathfrak{g} .

5.14. *Example.* Another example where $\mathrm{Spc}(A\text{-stmod})$ is isomorphic to the variety $\mathrm{Proj}(H^\bullet(A, k))$ is the algebra $A = k[X_1, \dots, X_n]/(X_1^\ell, \dots, X_n^\ell) \rtimes (\mathbb{Z}/\ell\mathbb{Z})^{\times n}$ which appears in Pevtsova-Witherspoon [PW15, Thm. 1.2].

6. EQUIVARIANT STABLE HOMOTOPY AND KK -THEORY

6.1. *Point.* Let G be a compact Lie group, e.g. a finite group, and let $\mathrm{SH}(G)$ be the equivariant stable homotopy category of genuine G -spectra. The tensor-triangulated category of compact (rigid) objects in $\mathrm{SH}(G)$ is denoted $\mathrm{SH}(G)^c$. In general, the spectrum of $\mathrm{SH}(G)^c$ is not quite known but significant progress occurred in recent years. It relies in an essential way on the non-equivariant case $G = 1$ of Section 3.

6.2. *Point.* For every chromatic tt-prime $\mathcal{P}_{p,n} \in \mathrm{Spc}(\mathrm{SH}^c)$ in the stable homotopy category SH^c (Theorem 3.3) and every closed subgroup $H \leq G$, let

$$\mathcal{P}(H, p, n) = (\Phi^H)^{-1}(\mathcal{P}_{p,n})$$

be its preimage under geometric H -fixed points $\Phi^H: \mathrm{SH}(G)^c \rightarrow \mathrm{SH}^c$, which is a tt-functor. This ‘equivariant’ tt-prime $\mathcal{P}(H, p, n)$ is the image of the chromatic $\mathcal{P}_{p,n}$ under $\mathrm{Spc}(\Phi^H): \mathrm{Spc}(\mathrm{SH}^c) \rightarrow \mathrm{Spc}(\mathrm{SH}(G)^c)$ as in Remark 2.10.

It is convenient to use the convention that $\mathcal{P}_{p,1}$ means $\mathcal{P}_{0,1}$ for all p . And similarly, to read $\mathcal{P}(H, p, 1)$ as $\mathcal{P}(H, 0, 1)$.

Let us first discuss the case where G is a finite group. Varying the subgroup $H \leq G$, the maps $\mathrm{Spc}(\Phi^H)$ cover $\mathrm{Spc}(\mathrm{SH}(G)^c)$ – a fact that is also true for general compact Lie groups, see Theorem 6.9.

6.3. **Theorem** (Balmer-Sanders [BS17]). *Let G be a finite group. Then every tt-prime in $\mathrm{SH}(G)^c$ is of the form $\mathcal{P}(H, p, n)$ for a unique subgroup $H \leq G$ up to conjugation and a unique chromatic tt-prime $\mathcal{P}_{p,n} \in \mathrm{Spc}(\mathrm{SH}^c)$. Understanding inclusions between tt-primes completely describes the topology on $\mathrm{Spc}(\mathrm{SH}(G)^c)$.*

If $K \triangleleft H$ is a normal subgroup of index $p > 0$, then $\mathcal{P}(K, p, n+1) \subset \mathcal{P}(H, p, n)$ for every $n \geq 1$. There is no inclusion $\mathcal{P}(K, q, n) \subseteq \mathcal{P}(H, p, m)$ unless the corresponding chromatic tt-primes are included $\mathcal{P}_{q,n} \subseteq \mathcal{P}_{p,m}$ (which forces $n \geq m$, and $p = q$ if $m > 1$) and K is conjugate to a q -subnormal subgroup of H (see 6.5).

6.4. *Point.* For finite groups of square-free order, like $G = C_p$ for instance, the above result completely describes $\mathrm{Spc}(\mathrm{SH}(G)^c)$, with its topology, and thus gives the tt-classification. This result was a first major example where $\mathrm{Spc}(\mathcal{K})$ was determined first and the tt-classification deduced as a corollary.

6.5. *Point.* For other groups, the question is to decide when $\mathcal{P}(K, p, n) \subset \mathcal{P}(H, p, m)$, in terms of $n - m$, for $K \leq H$ a p -subnormal subgroup of H (i.e. one such that there exists a tower of normal subgroups of index p from K to H). Theorem 6.3 implies that this inclusion holds when $n - m \geq \log_p([H : K])$.

The case of abelian groups (and a little more) was recently tackled in [BHN⁺19], showing that the above $\log_p([H : K])$ is not the sharpest bound.

6.6. **Theorem** (Barthel-Hausmann-Naumann-Nikolaus-Noel-Stapleton). *Let G be a finite abelian group, let $K \leq H \leq G$ be subgroups, let p be a prime and let $1 \leq n < \infty$ be an integer. Then the minimal i such that $\mathcal{P}(K, p, n) \subseteq \mathcal{P}(H, p, n - i)$ is $i = \mathrm{rk}_p(H/K)$ the p -rank of the quotient.*

See [BHN⁺19, Cor. 1.3]. The precise topology of $\mathrm{Spc}(\mathrm{SH}(G)^c)$ for general finite groups remains an open problem.

6.7. *Point.* Let us now consider the case of an arbitrary compact Lie group G but after rationalization $\mathrm{SH}(G)_{\mathbb{Q}}^c$. Of course, tensoring with \mathbb{Q} hides the ‘chromatic direction’ but this is an essential step in understanding the ‘equivariant direction’, by which we mean the role played by the subgroups of G .

A closed subgroup $K \leq H$ is *cotoral* if K is normal and H/K is a torus. Every closed subgroup $H \leq G$ defines a tt-prime \mathcal{P}_H in the spectrum $\mathrm{Spc}(\mathrm{SH}(G)_{\mathbb{Q}}^c)$, namely $\mathcal{P}_H := \mathrm{Ker}(\Phi^H)$ the kernel of geometric H -fixed points, i.e. the image of the unique prime (0) under the map $\mathrm{Spc}(\mathrm{D}^b(\mathbb{Q})) \rightarrow \mathrm{Spc}(\mathrm{SH}(G)^c)$ associated to the tt-functor $\Phi^H: \mathrm{SH}(G)_{\mathbb{Q}}^c \rightarrow \mathrm{SH}_{\mathbb{Q}}^c \cong \mathrm{D}^b(\mathbb{Q})$.

6.8. **Theorem** (Greenlees [Gre19, Thm. 1.3]). *Let G be a compact Lie group. Every tt-prime of the rational equivariant stable homotopy category $\mathrm{SH}(G)_{\mathbb{Q}}^c$ is equal to $\mathcal{P}_H = \mathrm{Ker}(\Phi^H)$ for a closed subgroup $H \leq G$, unique up to conjugation. Furthermore, specialization of tt-primes corresponds to cotoral inclusions: We have $\mathcal{P}_K \subseteq \mathcal{P}_H$ if and only if K is conjugate to a cotoral subgroup of H . The topology on $\mathrm{Spc}(\mathrm{SH}(G)_{\mathbb{Q}}^c)$ corresponds to the f -topology of [Gre98].*

Recently there has been further progress for arbitrary compact Lie group:

6.9. **Theorem** (Barthel-Greenlees-Hausmann [BGH18]). *Let G be a compact Lie group. Then every tt-prime of $\mathrm{SH}(G)^c$ is of the form $\mathcal{P}(H, p, n)$ as in 6.2. Moreover, the topology is completely understood in terms of inclusions of tt-primes.*

Barthel-Greenlees-Hausmann more precisely track the inclusion of primes, in terms of functions on the compact and totally-disconnected Hausdorff orbit space $\mathrm{Sub}(G)/G$ of G acting by conjugation on its closed subgroups. Furthermore, they give a complete description of the topology in the case of an *abelian* compact Lie group, extending Theorem 6.6; see [BGH18, Thm. 1.4].

* * *

6.10. *Point.* The closest to analysis that tt-geometry has gone so far is in the theory of C^* -algebras, via Kasparov’s KK -theory. Although this is not strictly speaking equivariant homotopy theory, we include it in this section as KK -theory belongs to the broad topic of noncommutative topology.

One begins with the ‘cellular’ subcategory, a. k. a. the ‘bootstrap’ category.

6.11. **Theorem** (Dell’Ambrogio [Del10, §6]). *Let G be a finite group and \mathcal{K}^G be the thick subcategory of the G -equivariant Kasparov category KK^G generated by the unit. Then the comparison map $\rho: \mathrm{Spc}(\mathcal{K}^G) \rightarrow \mathrm{Spec}(R(G))$ to the spectrum of the complex representation ring $R(G)$, as in Definition 8.2, is surjective and admits a continuous section. In the non-equivariant case, the above map induces a homeomorphism between the tt-spectrum of the so-called ‘bootstrap category’ $\mathcal{K}^1 = \mathrm{Boot}$ and $\mathrm{Spec}(\mathbb{Z})$.*

6.12. *Point.* Dell’Ambrogio conjectures that $\rho: \mathrm{Spc}(\mathcal{K}^G) \rightarrow \mathrm{Spec}(R(G))$ is a homeomorphism for all finite groups. The tt-classification for larger KK -categories, or for infinite groups, is another interesting open problem.

7. MOTIVES AND \mathbb{A}^1 -HOMOTOPY

7.1. *Point.* We consider here two classes of ‘motivic’ tensor-triangulated categories: First, we have the derived category of motives $DM(F; R)$, over a base field F and with coefficients in a ring R , as first introduced by Voevodsky [Voe00]; see also [Ayo14, CD09]. Secondly, we have $SH^{\mathbb{A}^1}(F)$ the stable \mathbb{A}^1 -homotopy category over the base F introduced by Morel and Voevodsky [MV99, Voe98, Mor06]. (Other base schemes can be considered.)


7.2. *Point.* Let us begin with $DM(F; R)$. Like in KK -theory (see 6.10), one can first consider the ‘cellular’ tt-subcategory of (mixed) Tate motives $DTM(F; R)$ generated as a localizing subcategory by the invertible Tate objects $R(i)$ for $i \in \mathbb{Z}$. Its subcategory of compact objects is the rigid tt-category $DTM(F; R)^c$ we shall discuss now.

7.3. *Point.* Peter [Pet13] established the tt-bridgehead into motivic territory, when he proved that the spectrum $\mathrm{Spc}(DTM(F; \mathbb{Q})^c) = *$ reduces to a point, for F a field satisfying the Beilinson-Soulé vanishing conjecture and a less standard restriction on rational motivic cohomology, namely $H_{\mathrm{mot}}^i(F; \mathbb{Q}(j)) = 0$ for $j \geq i \geq 2$. For instance, this applies to $F = \mathbb{Q}$. In fact, $\mathrm{Spc}(DTM(F; \mathbb{Q})^c) = *$ would follow from $\mathrm{Spc}(DM(F; \mathbb{Q})^c) = *$, a conjecture which is supported by:

7.4. **Theorem** (Kelly [Kel16, Thm. 36]). *Let F be a finite field such that every connected smooth projective variety X over F satisfies the Beilinson-Parshin conjecture and agreement of rational and numerical equivalence. Then $\mathrm{Spc}(DM(F; \mathbb{Q})^c) = *$ is a point.*

The above are rational results. Our understanding of the integral picture recently evolved thanks to the following breakthrough.

7.5. **Theorem** (Gallauer [Gal19, Thm. 8.6]). *Let F be an algebraically closed field of characteristic zero ⁽⁴⁾ whose rational motivic cohomology $H_{\mathrm{mot}}^i(F; \mathbb{Q}(j))$ vanishes for $i \leq 0 < j$ (Beilinson-Soulé) and for $j \geq i \geq 2$. Then the spectrum of $DTM(F; \mathbb{Z})^c$ is the following:*



where $\mathcal{P}_0 = \mathrm{Ker}(DTM(F; \mathbb{Z})^c \rightarrow DTM(F; \mathbb{Q})^c)$ consists of the torsion objects and, for every prime number p , the tt-primes $\mathcal{P}_{p,\mathrm{mot}}$ and $\mathcal{P}_{p,\mathrm{et}}$ are the kernels of motivic and étale cohomology with \mathbb{Z}/p coefficients, respectively.

7.6. *Example.* Peter’s conditions 7.3 on rational motivic cohomology are satisfied for $F = \mathbb{Q}$ for instance. This also provides an example for Theorem 7.5. Indeed, Gallauer’s result uses Peter’s theorem rationally.

7.7. *Remark.* Gallauer [Gal19] proves more general results about the derived category $DTM(F; \mathbb{Z}/p)^c$ of Tate motives with \mathbb{Z}/p coefficients without any condition about rational motivic cohomology. The latter theorem follows from the study of the derived categories of filtered modules in [GAdS18].

⁴In positive characteristic ℓ , replace the coefficients \mathbb{Z} by $\mathbb{Z}[1/\ell]$ and only allow $p \neq \ell$.

7.8. *Point.* In the case of Theorem 7.4, because -1 is a sum of squares in F , the rational derived category of motives coincides with the rational stable \mathbb{A}^1 -homotopy category $\mathrm{DM}(F; \mathbb{Q}) \cong \mathrm{SH}^{\mathbb{A}^1}(F; \mathbb{Q})$. Let us now mention some integral information about $\mathrm{SH}^{\mathbb{A}^1}(F)$. The first partial results about its spectrum were obtained in [Bal10a, §10]. The most advanced information is currently:

7.9. **Theorem** (Heller-Ormsby [HO18, Thm. 1.1]). *Let F be a field of characteristic different from 2. Then the comparison map (Definition 8.2) to the homogeneous spectrum of Milnor-Witt K -theory is surjective:*

$$\mathrm{Spc}(\mathrm{SH}^{\mathbb{A}^1}(F)^c) \rightarrow \mathrm{Spec}^h(K_*^{MW}(F)).$$

The exact computation of the tt-spectrum of $\mathrm{SH}(F)^c$ is a major open challenge, which involves understanding the fibers of the above map.

7.10. *Remark.* Partial results have also been obtained by Dell’Ambrogio and Tabuada [DT12] for non-commutative (dg-)motives.

8. POINTERS TO TT-GEOMETRY

8.1. *Point.* A snapshot of tensor-triangular geometry as of the year 2010 can be found in [Bal10b]. For a more recent survey, see [Ste18]. Beyond those references, let us simply highlight some aspects close to the author’s own research.

A very useful basic tool introduced in [Bal10a] is the following *comparison map* between tt-spectra and Zariski spectra of suitable graded rings:

8.2. *Definition.* Let $u \in \mathcal{K}$ be a \otimes -invertible object and $R_{\mathcal{K}, u}^\bullet = \bigoplus_{n \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{K}}(\mathbb{1}, u^{\otimes n})$ the associated graded-commutative graded ring. Then

$$\mathcal{P} \mapsto \{ f \in R_{\mathcal{K}, u}^\bullet \mid \mathrm{cone}(f) \notin \mathcal{P} \}$$

defines a continuous map $\rho^\bullet: \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spec}^h(R_{\mathcal{K}, u}^\bullet)$. Without grading, one can similarly define $\rho: \mathrm{Spc}(\mathcal{K}) \rightarrow \mathrm{Spec}(\mathrm{End}_{\mathcal{K}}(\mathbb{1}))$.

8.3. *Point.* Dell’Ambrogio-Stanley [DS16] give a class of cellular tt-categories for which ρ^\bullet is a homeomorphism, namely when the ring $R_{\mathcal{K}, \mathbb{1}}^\bullet$ is concentrated in even degrees and is ‘regular’ in a weak sense, which includes noetherian.

8.4. *Point.* The comparison map was generalized in two directions. First by Dell’Ambrogio-Stevenson [DS14], by allowing grading by a collection of invertible objects instead of a single one. Secondly, higher comparison maps were defined by Sanders [San13] in order to refine the analysis of the fibers of ‘lower’ comparison maps, through an inductive process.

8.5. *Point.* The above comparison map is still very much concerned with the computation of $\mathrm{Spc}(\mathcal{K})$. Moving away from this preoccupation, some first ‘geometric’ results were established in [Bal07], like the decomposition of an object associated to a decomposition of its support, and applications to filtrations of \mathcal{K} by (co)dimension of support. These ideas naturally led to *tensor-triangular Chow groups* in [Bal13] and further improvements by Klein [Kle16] and Belmans-Klein [BK17], using the already mentioned relative tt-geometry of [Ste13].

8.6. *Point.* In recent years, a great deal of progress followed from the development of the idea of *separable extensions* of tt-categories. The ubiquity of this notion through stable homotopy theory, in connection with equivariant ideas, can be seen in [BDS15]. As a slogan, this theory extends tt-geometry from the Zariski setting to the étale setting. Implications for the spectrum are discussed in [Bal16].

8.7. *Point.* Another area of tt-geometry which seems promising is the theory of *homological residue field*, which aims at abstractly understanding the various ‘fields’ which appear in examples: Morava K -theories, ordinary residue fields, π -points, etc. The reader can enter this ongoing project via [BKS19, Bal20].

REFERENCES

- [AS16] Benjamin Antieau and Greg Stevenson. Derived categories of representations of small categories over commutative Noetherian rings. *Pacific J. Math.*, 283(1):21–42, 2016.
- [Ayo14] Joseph Ayoub. A guide to (étale) motivic sheaves. In *Proc. of the International Congress of Mathematicians, Seoul (2014). Vol. II*, pages 1101–1124, 2014.
- [Bal05] Paul Balmer. The spectrum of prime ideals in tensor triangulated categories. *J. Reine Angew. Math.*, 588:149–168, 2005.
- [Bal07] Paul Balmer. Supports and filtrations in algebraic geometry and modular representation theory. *Amer. J. Math.*, 129(5):1227–1250, 2007.
- [Bal10a] Paul Balmer. Spectra, spectra, spectra – tensor triangular spectra versus Zariski spectra of endomorphism rings. *Algebr. Geom. Topol.*, 10(3):1521–1563, 2010.
- [Bal10b] Paul Balmer. Tensor triangular geometry. In *International Congress of Mathematicians, Hyderabad (2010), Vol. II*, pages 85–112. Hindustan Book Agency, 2010.
- [Bal13] Paul Balmer. Tensor triangular Chow groups. *J. Geom. Phys.*, 72:3–6, 2013.
- [Bal16] Paul Balmer. Separable extensions in tensor-triangular geometry and generalized Quillen stratification. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(4):907–925, 2016.
- [Bal20] Paul Balmer. Nilpotence theorems via homological residue fields. *Tunis. J. Math.*, 2(2):359–378, 2020.
- [BB19] Tobias Barthel and Agnès Beaudry. Chromatic structures in stable homotopy theory. In *Handbook of Homotopy Theory*. 2019. Preprint <https://arxiv.org/abs/1901.09004>.
- [BCHV19] Tobias Barthel, Natàlia Castellana, Drew Heard, and Gabriel Valenzuela. Stratification and duality for homotopical groups. *Adv. Math.*, 354:106733, 61, 2019.
- [BCR97] David J. Benson, Jon F. Carlson, and Jeremy Rickard. Thick subcategories of the stable module category. *Fund. Math.*, 153(1):59–80, 1997.
- [BCS19] Shawn Baland, Alexandru Chirvasitu, and Greg Stevenson. The prime spectra of relative stable module categories. *Trans. Amer. Math. Soc.*, 371(1):489–503, 2019.
- [BDS15] Paul Balmer, Ivo Dell’Ambrogio, and Beren Sanders. Restriction to finite-index subgroups as étale extensions in topology, KK-theory and geometry. *Algebr. Geom. Topol.*, 15(5):3025–3047, 2015.
- [BGH18] Tobias Barthel, John Greenlees, and Markus Hausmann. On the Balmer spectrum of compact Lie groups. Preprint <https://arxiv.org/abs/1810.04698>, 2018.
- [BHN⁺19] Tobias Barthel, Markus Hausmann, Niko Naumann, Thomas Nikolaus, Justin Noel, and Nathaniel Stapleton. The Balmer spectrum of the equivariant homotopy category of a finite abelian group. *Invent. Math.*, 216(1):215–240, 2019.
- [BIK11] Dave Benson, Srikanth B. Iyengar, and Henning Krause. Localising subcategories for cochains on the classifying space of a finite group. *C. R. Math. Acad. Sci. Paris*, 349(17-18):953–956, 2011.
- [BIK12] David J. Benson, Srikanth Iyengar, and Henning Krause. *Representations of finite groups: local cohomology and support*, volume 43 of *Oberwolfach Seminars*. Birkhäuser/Springer, Basel, 2012.
- [BIKP18] Dave Benson, Srikanth B. Iyengar, Henning Krause, and Julia Pevtsova. Stratification for module categories of finite group schemes. *J. Amer. Math. Soc.*, 31(1):265–302, 2018.
- [BK17] Pieter Belmans and Sebastian Klein. Relative tensor triangular Chow groups for coherent algebras. *J. Algebra*, 487:386–428, 2017.

- [BKN17a] Brian Boe, Jonathan Kujawa, and Daniel Nakano. Tensor triangular geometry for quantum groups. Preprint <https://arxiv.org/abs/1702.01289>, 2017.
- [BKN17b] Brian D. Boe, Jonathan R. Kujawa, and Daniel K. Nakano. Tensor triangular geometry for classical Lie superalgebras. *Adv. Math.*, 314:228–277, 2017.
- [BKS07] Aslak Bakke Buan, Henning Krause, and Øyvind Solberg. Support varieties: an ideal approach. *Homology, Homotopy Appl.*, 9(1):45–74, 2007.
- [BKS17] Paul Balmer, Henning Krause, and Greg Stevenson. The frame of smashing tensor-ideals. *Math. Proc. Cambridge Philos. Soc.*, to appear, preprint 2017. <https://arxiv.org/abs/1701.05937>.
- [BKS19] Paul Balmer, Henning Krause, and Greg Stevenson. Tensor-triangular fields: ruminations. *Selecta Math. (N.S.)*, 25(1):Art. 13, 36, 2019.
- [BS17] Paul Balmer and Beren Sanders. The spectrum of the equivariant stable homotopy category of a finite group. *Invent. Math.*, 208(1):283–326, 2017.
- [CD09] Denis-Charles Cisinski and Frédéric Déglise. Triangulated categories of mixed motives. Preprint <https://arxiv.org/abs/0912.2110>, 2009.
- [CI15] Jon F. Carlson and Srikanth B. Iyengar. Thick subcategories of the bounded derived category of a finite group. *Trans. Amer. Math. Soc.*, 367(4):2703–2717, 2015.
- [Dad78] Everett C. Dade. Endo-permutation modules over p -groups. I. *Ann. of Math. (2)*, 107(3):459–494, 1978.
- [Del10] Ivo Dell’Ambrogio. Tensor triangular geometry and KK -theory. *J. Homotopy Relat. Struct.*, 5(1):319–358, 2010.
- [DHS88] Ethan S. Devinatz, Michael J. Hopkins, and Jeffrey H. Smith. Nilpotence and stable homotopy theory. I. *Ann. of Math. (2)*, 128(2):207–241, 1988.
- [DM12] Umesh V. Dubey and Vivek M. Mallick. Spectrum of some triangulated categories. *J. Algebra*, 364:90–118, 2012.
- [DS14] Ivo Dell’Ambrogio and Greg Stevenson. Even more spectra: tensor triangular comparison maps via graded commutative 2-rings. *Appl. Categ. Structures*, 22(1):169–210, 2014.
- [DS16] Ivo Dell’Ambrogio and Donald Stanley. Affine weakly regular tensor triangulated categories. *Pacific J. Math.*, 285(1):93–109, 2016.
- [DT12] Ivo Dell’Ambrogio and Gonçalo Tabuada. Tensor triangular geometry of non-commutative motives. *Adv. Math.*, 229(2):1329–1357, 2012.
- [FP07] Eric M. Friedlander and Julia Pevtsova. Π -supports for modules for finite group schemes. *Duke Math. J.*, 139(2):317–368, 2007.
- [GAdS18] Martin Gallauer Alves de Souza. Tensor triangular geometry of filtered modules. *Algebra Number Theory*, 12(8):1975–2003, 2018.
- [Gal19] Martin Gallauer. tt -geometry of Tate motives over algebraically closed fields. *Compos. Math.*, 155(10):1888–1923, 2019.
- [Gre98] J. P. C. Greenlees. Rational Mackey functors for compact Lie groups. I. *Proc. London Math. Soc. (3)*, 76(3):549–578, 1998.
- [Gre19] J. P. C. Greenlees. The Balmer spectrum of rational equivariant cohomology theories. *J. Pure Appl. Algebra*, 223(7):2845–2871, 2019.
- [Hal16] Jack Hall. The Balmer spectrum of a tame stack. *Ann. K-Theory*, 1(3):259–274, 2016.
- [Hap88] Dieter Happel. *Triangulated categories in the representation theory of finite-dimensional algebras*, volume 119 of *LMS Lecture Note*. Cambr. Univ. Press, Cambridge, 1988.
- [Hir19] Yuki Hirano. Relative singular locus and Balmer spectrum of matrix factorizations. *Trans. Amer. Math. Soc.*, 371(7):4993–5021, 2019.
- [HO18] Jeremiah Heller and Kyle Ormsby. Primes and fields in stable motivic homotopy theory. *Geom. Topol.*, 22(4):2187–2218, 2018.
- [Hoc69] M. Hochster. Prime ideal structure in commutative rings. *Trans. Amer. Math. Soc.*, 142:43–60, 1969.
- [Hop87] Michael J. Hopkins. Global methods in homotopy theory. In *Homotopy theory (Durham, 1985)*, volume 117 of *LMS Lect. Note*, pages 73–96. Cambridge Univ. Press, 1987.
- [HPS97] Mark Hovey, John H. Palmieri, and Neil P. Strickland. Axiomatic stable homotopy theory. *Mem. Amer. Math. Soc.*, 128(610), 1997.

- [HS98] Michael J. Hopkins and Jeffrey H. Smith. Nilpotence and stable homotopy theory. II. *Ann. of Math. (2)*, 148(1):1–49, 1998.
- [Kel94] Bernhard Keller. A remark on the generalized smashing conjecture. *Manuscripta Math.*, 84(2):193–198, 1994.
- [Kel16] Shane Kelly. Some observations about motivic tensor triangulated geometry over a finite field. Preprint <https://arxiv.org/abs/1608.02913>, 2016.
- [Kle16] Sebastian Klein. Chow groups of tensor triangulated categories. *J. Pure Appl. Algebra*, 220(4):1343–1381, 2016.
- [KN02] B. Keller and A. Neeman. The connection between May’s axioms for a triangulated tensor product and Happel’s description of the derived category of the quiver D_4 . *Doc. Math.*, 7:535–560, 2002.
- [KP17] Joachim Kock and Wolfgang Pitsch. Hochster duality in derived categories and point-free reconstruction of schemes. *Trans. Amer. Math. Soc.*, 369(1):223–261, 2017.
- [Kra00] Henning Krause. Smashing subcategories and the telescope conjecture—an algebraic approach. *Invent. Math.*, 139(1):99–133, 2000.
- [Kri09] Amalendu Krishna. Perfect complexes on Deligne-Mumford stacks and applications. *J. K-Theory*, 4(3):559–603, 2009.
- [LS13] Yu-Han Liu and Susan J. Sierra. Recovering quivers from derived quiver representations. *Comm. Algebra*, 41(8):3013–3031, 2013.
- [Mat18] Hiroki Matsui. Connectedness of the Balmer spectrum of the right bounded derived category. *J. Pure Appl. Algebra*, 222(11):3733–3744, 2018.
- [Mor06] Fabien Morel. \mathbb{A}^1 -algebraic topology. In *International Congress of Mathematicians. Vol. II*, pages 1035–1059. Eur. Math. Soc., Zürich, 2006.
- [MT17] Hiroki Matsui and Ryo Takahashi. Thick tensor ideals of right bounded derived categories. *Algebra Number Theory*, 11(7):1677–1738, 2017.
- [Muk81] Shigeru Mukai. Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves. *Nagoya Math. J.*, 81:153–175, 1981.
- [MV99] Fabien Morel and Vladimir Voevodsky. \mathbb{A}^1 -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.*, (90):45–143 (2001), 1999.
- [Nee92] Amnon Neeman. The chromatic tower for $D(R)$. *Topology*, 31(3):519–532, 1992.
- [Nee01] Amnon Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, 2001.
- [Pet13] Tobias J. Peter. Prime ideals of mixed Artin-Tate motives. *Journal of K-theory*, 11(2):331–349, 2013.
- [PW15] Julia Pevtsova and Sarah Witherspoon. Tensor ideals and varieties for modules of quantum elementary abelian groups. *Proc. Amer. Math. Soc.*, 143(9):3727–3741, 2015.
- [Rav84] Douglas C. Ravenel. Localization with respect to certain periodic homology theories. *Amer. J. Math.*, 106(2):351–414, 1984.
- [Rav92] Douglas C. Ravenel. *Nilpotence and periodicity in stable homotopy theory*, volume 128 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1992.
- [Ric89] Jeremy Rickard. Derived categories and stable equivalence. *J. Pure Appl. Algebra*, 61(3):303–317, 1989.
- [San13] Beren Sanders. Higher comparison maps for the spectrum of a tensor triangulated category. *Adv. Math.*, 247:71–102, 2013.
- [Ste13] Greg Stevenson. Support theory via actions of tensor triangulated categories. *J. Reine Angew. Math.*, 681:219–254, 2013.
- [Ste14] Greg Stevenson. Subcategories of singularity categories via tensor actions. *Compos. Math.*, 150(2):229–272, 2014.
- [Ste18] Greg Stevenson. A tour of support theory for triangulated categories through tensor triangular geometry. In *Building bridges between algebra and topology*, Adv. Courses Math. CRM Barcelona, pages 63–101. Birkhäuser/Springer, Cham, 2018.
- [Tak10] Ryo Takahashi. Classifying thick subcategories of the stable category of Cohen-Macaulay modules. *Adv. Math.*, 225(4):2076–2116, 2010.
- [Tho97] R. W. Thomason. The classification of triangulated subcategories. *Compositio Math.*, 105(1):1–27, 1997.
- [Voe98] Vladimir Voevodsky. \mathbb{A}^1 -homotopy theory. In *Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998)*, pages 579–604, 1998.

- [Voe00] Vladimir Voevodsky. Triangulated categories of motives over a field. In *Cycles, transfers, and motivic homology theories*, volume 143 of *Ann. of Math. Stud.*, pages 188–238. Princeton Univ. Press, Princeton, NJ, 2000.
- [Wan19] Ren Wang. The spectrum of the singularity category of a category algebra. *Appl. Categ. Structures*, 27(4):427–433, 2019.
- [Xu14] Fei Xu. Spectra of tensor triangulated categories over category algebras. *Arch. Math. (Basel)*, 103(3):235–253, 2014.

PAUL BALMER, MATHEMATICS DEPARTMENT, UCLA, LOS ANGELES, CA 90095-1555, USA
E-mail address: balmer@math.ucla.edu
URL: <http://www.math.ucla.edu/~balmer>