

Derivative tests

Nicholas Hu · Last updated on 2026-02-02

Let $I \subseteq \mathbb{R}$ be a nonempty interval and $f : I \rightarrow \mathbb{R}$ be a function. We denote the interior of I by I° .

Monotonicity

We say that f is **increasing** if $f(x) \leq f(y)$ for all $x, y \in I$ with $x < y$ and that f is **strictly increasing** if the former inequality is strict. We define **decreasing** and **strictly decreasing** analogously.

Characterization of monotonicity

Suppose that f is continuous on I and differentiable on I° .¹ Then f is increasing on I if and only if $f' \geq 0$ on I° . In addition, f is strictly increasing on I if $f' > 0$ on I° .²

Extremality

We say that $x \in I$ is a **(global) minimizer** and that the value $f(x)$ is a **(global) minimum** of f if $f(x) \leq f(y)$ for all $y \in I$. We define **(global) maximizer** and **(global) maximum** analogously.

Extreme value theorem

If I is closed and bounded and f is continuous on I , then f has a minimizer and a maximizer in I .

We say that $x \in I$ is a **local minimizer** and that the value $f(x)$ is a **local minimum** of f if there exists an $\varepsilon > 0$ such that $f(x) \leq f(y)$ for all $y \in I \cap (x - \varepsilon, x + \varepsilon)$. We define **local maximizer** and **local maximum** analogously. Clearly, any global extremizer must also be a local extremizer.

Fermat's theorem

Suppose that $x \in I^\circ$ is a local extremizer of f . If f is differentiable at x , then $f'(x) = 0$ (that is, x is a **stationary point** of f).

Thus, under the hypotheses of the extreme value theorem, any extremizer of f will either be an interior point of I and hence a point at which f is stationary *or* not differentiable, or else will be a boundary point of I . We call a point of the former type (that is, a point at which f is stationary *or* not differentiable) a **critical point** of f .

First derivative test

If f is continuous at x and there exists an $\varepsilon > 0$ such that $f' \leq 0$ on $(x - \varepsilon, x)$ and $f' \geq 0$ on $(x, x + \varepsilon)$, then x is a local minimizer of f .

Second derivative test

If f is stationary at x and $f''(x) > 0$, then x is a local minimizer of f .

Convexity

We say that f is **convex** if $f((1 - \theta)x + \theta y) \leq (1 - \theta)f(x) + \theta f(y)$ for all *distinct* $x, y \in I$ and $\theta \in (0, 1)$ and that f is **strictly convex** if the inequality is strict. We define **concave** and **strictly concave** analogously.

Secant and tangent line characterizations of convexity

Given an $x \in I$, let $g(y; x) := \frac{f(y) - f(x)}{y - x}$ for all $y \in I \setminus \{x\}$. Then f is convex (resp., strictly convex) on I if and only if g is increasing (resp., strictly increasing) in y for all $x \in I$. In addition, if f is differentiable on I° , then f is convex (resp., strictly convex) on I if and only if $g(y; x) \geq f'(x)$ (resp., $g(y; x) > f'(x)$) for all $x \in I^\circ$ and $y \in I \setminus \{x\}$.

First-order characterization of convexity

Suppose that f is continuous on I and differentiable on I° . Then f is convex (resp., strictly convex) on I if and only if f' is increasing (resp., strictly increasing) on I° .

Second-order characterization of convexity

Suppose that f is continuous on I and twice differentiable on I° . Then f is convex on I if and only if $f'' \geq 0$ on I° . In addition, f is strictly convex on I if $f'' > 0$ on I° .

Minimizers of convex functions

If f is convex and x is a local minimizer of f , then x is a global minimizer of f . In addition, if f is strictly convex, then f has *at most one* local minimizer.

We say that f has an **inflection point** at x if it is continuous and changes from strictly convex to strictly concave or vice-versa at x .³

1. If I is open, then $I^\circ = I$ and the hypothesis reduces to f being differentiable on I . ↵

2. The converse is false in general: consider $f(x) = x^3$ on $I = \mathbb{R}$. ↵

3. Definitions vary - some require that the graph of f have a (possibly vertical) tangent line at $(x, f(x))$; others require that f be twice differentiable near (but not at) x . ↵